

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
4 April 2002 (04.04.2002)

PCT

(10) International Publication Number
WO 02/26982 A2

(51) International Patent Classification⁷: C12N 15/12,
C12Q 1/68, C07K 14/47, 16/18, A01K 67/027, A61K
38/17, 39/395, B01J 19/00, G01N 33/50, 33/53

(21) International Application Number: PCT/US01/30042

(22) International Filing Date:
25 September 2001 (25.09.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/236,869 29 September 2000 (29.09.2000) US
60/239,812 11 October 2000 (11.10.2000) US
60/240,108 12 October 2000 (12.10.2000) US
60/241,282 17 October 2000 (17.10.2000) US
60/242,218 20 October 2000 (20.10.2000) US

(71) Applicant (for all designated States except US): INCYTE
GENOMICS, INC. [US/US]; 3160 Porter Drive, Palo
Alto, CA 94087 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): YUE, Henry
[US/US]; 826 Lois Avenue, Sunnyvale, CA 94087 (US).
TANG, Y., Tom [US/US]; 4230 Ranwick Court, San
Jose, CA 95118 (US). NGUYEN, Daniel, B. [US/US];
1403 Ridgewood Drive, San Jose, CA 95118 (US). YAO,
Monique, G. [US/US]; 1189 Woodgate Drive, Carmel,
IN 46633 (US). XU, Yuming [US/US]; 1739 Walnut
Drive, Mountain View, CA 94040 (US). TRIBOULEY,
Catherine, M. [FR/US]; 1121 Tennessee Street #5, San
Francisco, CA 94107 (US). SANJANWALA, Madhu,
S. [US/US]; 210 Sylvia Court, Los Altos, CA 94024
(US). WALIA, Narinder, K. [US/US]; 890 Davis Street
#205, San Leandro, CA 94577 (US). BAUGHN, Mariah,
R. [US/US]; 14244 Santiago Road, San Leandro, CA
94577 (US). SAPPERSTEIN, Stephanie, K. [US/US];
3531 Highland Avenue, Redwood City, CA 94062 (US).
LAL, Preeti [IN/US]; P.O. Box 5142, Santa Clara, CA
95056 (US). THORNTON, Michael [US/US]; 9 Medway
Road, Woodside, CA 94062 (US). GANDHI, Ameena,
R. [US/US]; 837 Roble Avenue #1, Menlo Park, CA
94025 (US). RAMKUMAR, Jayalaxmi [US/US]; 34359
Maybird Circle, Fremont, CA 94555 (US). ELLIOTT,
Vicki, S. [US/US]; 3770 Polten Place Way, San Jose, CA
95121 (US). ARVIZU, Chandra [US/US]; 490 Sherwood

Way #1, Menlo Park, CA 94025 (US). THANGAVELU,
Kavitha [IN/US]; 1950 Montecito Avenue #23, Mountain
View, CA 94043 (US). GIETZEN, Kimberly, J. [US/US];
691 Los Huecos Drive, San Jose, CA 95123 (US). DING,
Li [CH/US]; 3353 Alma Street #146, Palo Alto, CA
94506 (US). AU-YOUNG, Janice [US/US]; 233 Golden
Eagle Lane, Brisbane, CA 94005 (US). TRAN, Bao
[US/US]; 750 Salberg Avenue, Santa Clara, CA 95051
(US). POLICKY, Jennifer, L. [US/US]; 1511 Jarvis
Court, San Jose, CA 95118 (US). LEE, Sally [US/US];
825 East Evelyn #425, Sunnyvale, CA 94086 (US). LU,
Dyung, Aina, M. [US/US]; 233 Coy Drive, San Jose, CA
95123 (US). BURFORD, Neil [GB/US]; 105 Wildwood
Circle, Durham, CT 06422 (US). WARREN, Bridget,
A. [US/US]; 10130 Parwood Drive #2, Cupertino, CA
95014 (US). GURURAJAN, Rajagopal [IN/US]; 5591
Dent Avenue, San Jose, CA 95118 (US). DUGGAN,
Brendan, M. [AU/US]; 243 Buena Vista Avenue #306,
Sunnyvale, CA 94086 (US). HONCHELL, Cynthia, D.
[US/US]; 400 Laurel Street #203, San Carlos, CA 94070
(US). HAFALIA, April, J., A. [US/US]; 2227 Calle de
Primavera, Santa Clara, CA 95054 (US).

(74) Agents: HAMLET-COX, Diana et al.; Incyte Genomics,
Inc., 3160 Porter Drive, Palo Alto, CA 94304 (US).

(81) Designated States (national): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK,
SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA,
ZW.

(84) Designated States (regional): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,
CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD,
TG).

Published:

— without international search report and to be republished
upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: SECRETED PROTEINS

(57) Abstract: The invention provides human secreted proteins (SECP) and polynucleotides which identify and encode SECP. The invention also provides expression vectors, host cells, antibodies, agonists, and antagonists. The invention also provides methods for diagnosing, treating, or preventing disorders associated with aberrant expression of SECP.



WO 02/26982 A2

SECRETED PROTEINS

TECHNICAL FIELD

This invention relates to nucleic acid and amino acid sequences of secreted proteins and to the use of these sequences in the diagnosis, treatment, and prevention of cell proliferative, autoimmune/inflammatory, cardiovascular, neurological, and developmental disorders, and in the assessment of the effects of exogenous compounds on the expression of nucleic acid and amino acid sequences of secreted proteins.

BACKGROUND OF THE INVENTION

Protein transport and secretion are essential for cellular function. Protein transport is mediated by a signal peptide located at the amino terminus of the protein to be transported or secreted. The signal peptide is comprised of about ten to twenty hydrophobic amino acids which target the nascent protein from the ribosome to a particular membrane bound compartment such as the endoplasmic reticulum (ER). Proteins targeted to the ER may either proceed through the secretory pathway or remain in any of the secretory organelles such as the ER, Golgi apparatus, or lysosomes. Proteins that transit through the secretory pathway are either secreted into the extracellular space or retained in the plasma membrane. Proteins that are retained in the plasma membrane contain one or more transmembrane domains, each comprised of about 20 hydrophobic amino acid residues. Secreted proteins are generally synthesized as inactive precursors that are activated by post-translational processing events during transit through the secretory pathway. Such events include glycosylation, proteolysis, and removal of the signal peptide by a signal peptidase. Other events that may occur during protein transport include chaperone-dependent unfolding and folding of the nascent protein and interaction of the protein with a receptor or pore complex. Examples of secreted proteins with amino terminal signal peptides are discussed below and include proteins with important roles in cell-to-cell signaling. Such proteins include transmembrane receptors and cell surface markers, extracellular matrix molecules, cytokines, hormones, growth and differentiation factors, enzymes, neuropeptides, vasomediators, cell surface markers, and antigen recognition molecules. (Reviewed in Alberts, B. et al. (1994) Molecular Biology of The Cell, Garland Publishing, New York, NY, pp. 557-560, 582-592.)

Cell surface markers include cell surface antigens identified on leukocytic cells of the immune system. These antigens have been identified using systematic, monoclonal antibody (mAb)-based "shot gun" techniques. These techniques have resulted in the production of hundreds of mAbs directed against unknown cell surface leukocytic antigens. These antigens have been grouped into "clusters of differentiation" based on common immunocytochemical localization patterns in various

differentiated and undifferentiated leukocytic cell types. Antigens in a given cluster are presumed to identify a single cell surface protein and are assigned a "cluster of differentiation" or "CD" designation. Some of the genes encoding proteins identified by CD antigens have been cloned and verified by standard molecular biology techniques. CD antigens have been characterized as both

5 transmembrane proteins and cell surface proteins anchored to the plasma membrane via covalent attachment to fatty acid-containing glycolipids such as glycosylphosphatidylinositol (GPI). (Reviewed in Barclay, A. N. et al. (1995) The Leucocyte Antigen Facts Book, Academic Press, San Diego, CA, pp. 17-20.)

Matrix proteins (MPs) are transmembrane and extracellular proteins which function in

10 formation, growth, remodeling, and maintenance of tissues and as important mediators and regulators of the inflammatory response. The expression and balance of MPs may be perturbed by biochemical changes that result from congenital, epigenetic, or infectious diseases. In addition, MPs affect leukocyte migration, proliferation, differentiation, and activation in the immune response. MPs are frequently characterized by the presence of one or more domains which may include collagen-like

15 domains, EGF-like domains, immunoglobulin-like domains, and fibronectin-like domains. In addition, MPs may be heavily glycosylated and may contain an Arginine-Glycine-Aspartate (RGD) tripeptide motif which may play a role in adhesive interactions. MPs include extracellular proteins such as fibronectin, collagen, galectin, vitronectin and its proteolytic derivative somatomedin B; and cell adhesion receptors such as cell adhesion molecules (CAMs), cadherins, and integrins. (Reviewed

20 in Ayad, S. et al. (1994) The Extracellular Matrix Facts Book, Academic Press, San Diego, CA, pp. 2-16; Ruoslahti, E. (1997) *Kidney Int.* 51:1413-1417; Sjaastad, M.D. and Nelson, W.J. (1997) *BioEssays* 19:47-55.)

Mucins are highly glycosylated glycoproteins that are the major structural component of the mucus gel. The physiological functions of mucins are cytoprotection, mechanical protection,

25 maintenance of viscosity in secretions, and cellular recognition. MUC6 is a human gastric mucin that is also found in gall bladder, pancreas, seminal vesicles, and female reproductive tract (Toribara, N.W. et al. (1997) *J. Biol. Chem.* 272:16398-16403). The MUC6 gene has been mapped to human chromosome 11 (Toribara, N.W. et al. (1993) *J. Biol. Chem.* 268:5879-5885). Hemomucin is a novel Drosophila surface mucin that may be involved in the induction of antibacterial effector molecules

30 (Theopold, U. et al. (1996) *J. Biol. Chem.* 271:12708-12715).

Tuftelins are one of four different enamel matrix proteins that have been identified so far. The other three known enamel matrix proteins are the amelogenins, enamelin and ameloblastin. Assembly of the enamel extracellular matrix from these component proteins is believed to be critical in producing a matrix competent to undergo mineral replacement. (Paine, C.T. et al. (1998) *Connect*

35 *Tissue Res.* 38:257-267). Tuftelin mRNA has been found to be expressed in human ameloblastoma

tumor, a non-mineralized odontogenic tumor (Deutsch, D. et al. (1998) Connect. Tissue Res. 39:177-184).

Olfactomedin-related proteins are extracellular matrix, secreted glycoproteins with conserved C-terminal motifs. They are expressed in a wide variety of tissues and in broad range of species, from *Caenorhabditis elegans* to *Homo sapiens*. Olfactomedin-related proteins comprise a gene family with at least 5 family members in humans. One of the five, TIGR/myocilin protein, is expressed in the eye and is associated with the pathogenesis of glaucoma (Kulkarni, N.H. et al. (2000) Genet. Res. 76:41-50). Research by Yokoyama et al. (1996) found a 135-amino acid protein, termed AMY, having 96% sequence identity with rat neuronal olfactomedin-related ER localized protein in a neuroblastoma cell line cDNA library, suggesting an essential role for AMY in nerve tissue (Yokoyama, M. et al. (1996) DNA Res. 3:311-320). Neuron-specific olfactomedin-related glycoproteins isolated from rat brain cDNA libraries show strong sequence similarity with olfactomedin. This similarity is suggestive of a matrix-related function of these glycoproteins in neurons and neurosecretory cells (Danielson, P.E. et al. (1994) J. Neurosci. Res. 38:468-478).

Mac-2 binding protein is a 90-kD serum protein (90K), a secreted glycoprotein isolated from both the human breast carcinoma cell line SK-BR-3, and human breast milk. It specifically binds to a human macrophage-associated lectin, Mac-2. Structurally, the mature protein is 567 amino acids in length and is preceded by an 18-amino acid leader. There are 16 cysteines and seven potential N-linked glycosylation sites. The first 106 amino acids represent a domain very similar to an ancient protein superfamily defined by a macrophage scavenger receptor cysteine-rich domain (Koths, K. et al. (1993) J. Biol. Chem. 268:14245-14249). 90K is elevated in the serum of subpopulations of AIDS patients and is expressed at varying levels in primary tumor samples and tumor cell lines. Ullrich et al. (1994) have demonstrated that 90K stimulates host defense systems and can induce interleukin-2 secretion. This immune stimulation is proposed to be a result of oncogenic transformation, viral infection or pathogenic invasion (Ullrich, A., et al. (1994) J. Biol. Chem. 269:18401-18407).

Semaphorins are a large group of axonal guidance molecules consisting of at least 30 different members and are found in vertebrates, invertebrates, and even certain viruses. All semaphorins contain the sema domain which is approximately 500 amino acids in length. Neuropilin, a semaphorin receptor, has been shown to promote neurite outgrowth in vitro. The extracellular region of neuropilins consists of three different domains: CUB, discoidin, and MAM domains. The CUB and the MAM motifs of neuropilin have been suggested to have roles in protein-protein interactions and are thought to be involved in the binding of semaphorins through the sema and the C-terminal domains (reviewed in Raper, J.A. (2000) Curr. Opin. Neurobiol. 10:88-94). Plexins are neuronal cell surface molecules that mediate cell adhesion via a homophilic binding mechanism in the presence of calcium ions. Plexins have been shown to be expressed in the receptors and neurons of

particular sensory systems (Ohta, K. et al. (1995) *Cell* 14:1189-1199). There is evidence that suggests that some plexins function to control motor and CNS axon guidance in the developing nervous system. Plexins, which themselves contain complete semaphorin domains, may be both the ancestors of classical semaphorins and binding partners for semaphorins (Winberg, M.L. et al (1998) *Cell* 95:903-916).

Human pregnancy-specific beta 1-glycoprotein (PSG) is a family of closely related glycoproteins of molecular weights of 72 KDa, 64KDa, 62KDa, and 54KDa. Together with the carcinoembryonic antigen, they comprise a subfamily within the immunoglobulin superfamily (Plouzek, C.A. and Chou, J.Y. (1991) *Endocrinology* 129:950-958) Different subpopulations of PSG have been found to be produced by the trophoblasts of the human placenta, and the amnionic and chorionic membranes (Plouzek, C.A. et al. (1993) *Placenta* 14:277-285).

Autocrine motility factor (AMF) is one of the motility cytokines regulating tumor cell migration; therefore identification of the signaling pathway coupled with it has critical importance. Autocrine motility factor receptor (AMFR) expression has been found to be associated with tumor progression in thymoma (Ohta Y. et al. (2000) *Int. J. Oncol.* 17:259-264). AMFR is a cell surface glycoprotein of molecular weight 78KDa.

Hormones are secreted molecules that travel through the circulation and bind to specific receptors on the surface of, or within, target cells. Although they have diverse biochemical compositions and mechanisms of action, hormones can be grouped into two categories. One category includes small lipophilic hormones that diffuse through the plasma membrane of target cells, bind to cytosolic or nuclear receptors, and form a complex that alters gene expression. Examples of these molecules include retinoic acid, thyroxine, and the cholesterol-derived steroid hormones such as progesterone, estrogen, testosterone, cortisol, and aldosterone. The second category includes hydrophilic hormones that function by binding to cell surface receptors that transduce signals across the plasma membrane. Examples of such hormones include amino acid derivatives such as catecholamines (epinephrine, norepinephrine) and histamine, and peptide hormones such as glucagon, insulin, gastrin, secretin, cholecystokinin, adrenocorticotrophic hormone, follicle stimulating hormone, luteinizing hormone, thyroid stimulating hormone, and vasopressin. (See, for example, Lodish et al. (1995) *Molecular Cell Biology*, Scientific American Books Inc., New York, NY, pp. 856-864.)

Pro-opiomelanocortin (POMC) is the precursor polypeptide of corticotropin (ACTH), a hormone synthesized by the anterior pituitary gland, which functions in the stimulation of the adrenal cortex. POMC is also the precursor polypeptide of the hormone beta-lipotropin (beta-LPH). Each hormone includes smaller peptides with distinct biological activities: alpha-melanotropin (alpha-MSH) and corticotropin-like intermediate lobe peptide (CLIP) are formed from ACTH; gamma-lipotropin (gamma-LPH) and beta-endorphin are peptide components of beta-LPH; while beta-MSH

is contained within gamma-LPH. Adrenal insufficiency due to ACTH deficiency, resulting from a genetic mutation in exons 2 and 3 of POMC results in an endocrine disorder characterized by early-onset obesity, adrenal insufficiency, and red hair pigmentation (Chretien, M. et al. (1979) *Canad. J. Biochem.* 57:1111-1121; Krude, H. et al. (1998) *Nature Genet.* 19:155-157; Online Mendelian

5 Inheritance in Man (OMIM) 176830).

Growth and differentiation factors are secreted proteins which function in intercellular communication. Some factors require oligomerization or association with membrane proteins for activity. Complex interactions among these factors and their receptors trigger intracellular signal transduction pathways that stimulate or inhibit cell division, cell differentiation, cell signaling, and

10 cell motility. Most growth and differentiation factors act on cells in their local environment (paracrine signaling). There are three broad classes of growth and differentiation factors. The first class includes the large polypeptide growth factors such as epidermal growth factor, fibroblast growth factor, transforming growth factor, insulin-like growth factor, and platelet-derived growth factor. The second class includes the hematopoietic growth factors such as the colony stimulating factors (CSFs).

15 Hematopoietic growth factors stimulate the proliferation and differentiation of blood cells such as B-lymphocytes, T-lymphocytes, erythrocytes, platelets, eosinophils, basophils, neutrophils, macrophages, and their stem cell precursors. The third class includes small peptide factors such as bombesin, vasopressin, oxytocin, endothelin, transferrin, angiotensin II, vasoactive intestinal peptide, and bradykinin, which function as hormones to regulate cellular functions other than proliferation.

20 Growth and differentiation factors play critical roles in neoplastic transformation of cells in vitro and in tumor progression in vivo. Inappropriate expression of growth factors by tumor cells may contribute to vascularization and metastasis of tumors. During hematopoiesis, growth factor misregulation can result in anemias, leukemias, and lymphomas. Certain growth factors such as interferon are cytotoxic to tumor cells both in vivo and in vitro. Moreover, some growth factors and

25 growth factor receptors are related both structurally and functionally to oncoproteins. In addition, growth factors affect transcriptional regulation of both proto-oncogenes and oncosuppressor genes. (Reviewed in Pimentel, E. (1994) Handbook of Growth Factors, CRC Press, Ann Arbor, MI, pp. 1-9.)

The Slit protein, first identified in *Drosophila*, is critical in central nervous system midline formation and potentially in nervous tissue histogenesis and axonal pathfinding. Itoh et al. have

30 identified mammalian homologues of the slit gene (human Slit-1, Slit-2, Slit-3 and rat Slit-1). The encoded proteins are putative secreted proteins containing EGF-like motifs and leucine-rich repeats, both of which are conserved protein-protein interaction domains. Slit-1, -2, and -3 mRNAs are expressed in the brain, spinal cord, and thyroid, respectively (Itoh, A. et al., (1998) *Brain Res. Mol. Brain Res.* 62:175-186). The Slit family of proteins are indicated to be functional ligands of

35 glypican-1 in nervous tissue and suggests that their interactions may be critical in certain stages

during central nervous system histogenesis (Liang, Y. et al., (1999) J. Biol. Chem. 274:17885-17892).

Neuropeptides and vasomediators (NP/VM) comprise a large family of endogenous signaling molecules. Included in this family are neuropeptides and neuropeptide hormones such as bombesin, neuropeptide Y, neurotensin, neuromedin N, melanocortins, opioids, galanin, somatostatin, tachykinins, urotensin II and related peptides involved in smooth muscle stimulation, vasopressin, vasoactive intestinal peptide, and circulatory system-borne signaling molecules such as angiotensin, complement, calcitonin, endothelins, formyl-methionyl peptides, glucagon, cholecystokinin and gastrin. NP/VMs can transduce signals directly, modulate the activity or release of other neurotransmitters and hormones, and act as catalytic enzymes in cascades. The effects of NP/VMs range from extremely brief to long-lasting. (Reviewed in Martin, C.R. et al. (1985) Endocrine Physiology, Oxford University Press, New York, NY, pp. 57-62.)

NP/VMs are involved in numerous neurological and cardiovascular disorders. For example, neuropeptide Y is involved in hypertension, congestive heart failure, affective disorders, and appetite regulation. Somatostatin inhibits secretion of growth hormone and prolactin in the anterior pituitary, as well as inhibiting secretion in intestine, pancreatic acinar cells, and pancreatic beta-cells. A reduction in somatostatin levels has been reported in Alzheimer's disease and Parkinson's disease. Vasopressin acts in the kidney to increase water and sodium absorption, and in higher concentrations stimulates contraction of vascular smooth muscle, platelet activation, and glycogen breakdown in the liver. Vasopressin and its analogues are used clinically to treat diabetes insipidus. Endothelin and angiotensin are involved in hypertension, and drugs, such as captopril, which reduce plasma levels of angiotensin, are used to reduce blood pressure (Watson, S. and S. Arkin (1994) The G-protein Linked Receptor Facts Book, Academic Press, San Diego CA, pp. 194; 252; 284; 55; 111).

Neuropeptides have also been shown to have roles in nociception (pain). Vasoactive intestinal peptide appears to play an important role in chronic neuropathic pain. Nociceptin, an endogenous ligand for the opioid receptor-like 1 receptor, is thought to have a predominantly anti-nociceptive effect, and has been shown to have analgesic properties in different animal models of tonic or chronic pain (Dickinson, T. and Fleetwood-Walker, S.M. (1998) Trends Pharmacol. Sci. 19:346-348).

Other proteins that contain signal peptides include secreted proteins with enzymatic activity, or enzyme inhibitory activity. Such activity includes, for example, oxidoreductase/dehydrogenase activity, transferase activity, hydrolase activity, lyase activity, isomerase activity, or ligase activity. For example, matrix metalloproteinases are secreted hydrolytic enzymes that degrade the extracellular matrix and thus play an important role in tumor metastasis, tissue morphogenesis, and arthritis (Reponen, P. et al. (1995) Dev. Dyn. 202:388-396; Firestein, G.S. (1992) Curr. Opin. Rheumatol. 4:348-354; Ray, J.M. and Stetler-Stevenson, W.G. (1994) Eur. Respir. J. 7:2062-2072; and Mignatti,

P. and Rifkin, D.B. (1993) *Physiol. Rev.* 73:161-195). Tissue Inhibitors of Metalloproteinase (TIMPs), on the other hand, are secreted proteins which bind to metalloproteinases and block their activity (Stetler-Stevenson, W.G. et al. (1989) *J. Biol. Chem.* 264:17374-17378). Additional examples are the acetyl-CoA synthetases which activate acetate for use in lipid synthesis or energy generation (Luong, A. et al. (2000) *J. Biol. Chem.* 275:26458-26466). The result of acetyl-CoA synthetase activity is the formation of acetyl-CoA from acetate and CoA. Acetyl-CoA synthetases share a region of sequence similarity identified as the AMP-binding domain signature. Acetyl-CoA synthetase has been shown to be associated with hypertension (H. Toh (1991) *Protein Seq. Data Anal.* 4:111-117; and Iwai, N. et al., (1994) *Hypertension* 23:375-380).

10 A number of isomerases catalyze steps in protein folding, phototransduction, and various anabolic and catabolic pathways. One class of isomerases is known as peptidyl-prolyl *cis-trans* isomerases (PPIases). PPIases catalyze the *cis* to *trans* isomerization of certain proline imidic bonds in proteins. Two families of PPIases are the FK506 binding proteins (FKBPs), and cyclophilins (CyPs). FKBPs bind the potent immunosuppressants FK506 and rapamycin, thereby inhibiting
15 signaling pathways in T-cells. Specifically, the PPIase activity of FKBPs is inhibited by binding of FK506 or rapamycin. There are five members of the FKBP family which are named according to their calculated molecular masses (FKBP12, FKBP13, FKBP25, FKBP52, and FKBP65), and localized to different regions of the cell where they associate with different protein complexes (Coss, M. et al. (1995) *J. Biol. Chem.* 270:29336 - 29341; Schreiber, S.L. (1991) *Science* 251:283 - 287).

20 The peptidyl-prolyl isomerase activity of CyP may be part of the signaling pathway that leads to T-cell activation. CyP isomerase activity is associated with protein folding and protein trafficking, and may also be involved in assembly/disassembly of protein complexes and regulation of protein activity. For example, in *Drosophila*, the CyP NinaA is required for correct localization of rhodopsins, while a mammalian CyP (Cyp40) is part of the Hsp90/Hsc70 complex that binds steroid
25 receptors. The mammalian CypA has been shown to bind the gag protein from human immunodeficiency virus 1 (HIV-1), an interaction that can be inhibited by cyclosporin. Since cyclosporin has potent anti-HIV-1 activity, CypA may play an essential function in HIV-1 replication. Finally, Cyp40 has been shown to bind and inactivate the transcription factor c-Myb, an effect that is reversed by cyclosporin. This effect implicates CyPs in the regulation of transcription,
30 transformation, and differentiation (Bergsma, D.J. et al (1991) *J. Biol. Chem.* 266:23204 - 23214; Hunter, T. (1998) *Cell* 92: 141-143; and Levenson, J.D. and Ness, S.A. (1998) *Mol. Cell.* 1:203-211).

Another protein that contains a signal peptide is encoded by the seizure-related gene, SEZ-6, a brain specific cDNA whose expression is increased by the convulsant drug pentylentetrazole. The SEZ-6 protein is expressed in the cerebrum and cerebellum. SEZ-6 contains five short consensus
35 repeats (SCR, or sushi domains) and two CUB (complement C1r/s-like repeat) domains in addition to

a signal peptide and a single transmembrane domain (Shimizu-Nishikawa, K. et al. (1995) Biochem. Biophys. Res. Commun. 216:382-389).

Gamma-carboxyglutamic acid (Gla) proteins rich in proline (PRGPs) are members of a family of vitamin K-dependent single-pass integral membrane proteins. These proteins are characterized by an extracellular amino terminal domain of approximately 45 amino acids rich in Gla. The intracellular carboxyl terminal region contains one or two copies of the sequence PPXY, a motif present in a variety of proteins involved in such diverse cellular functions as signal transduction, cell cycle progression, and protein turnover (Kulman, J.D. et al., (2001) Proc. Natl. Acad. Sci. U.S.A. 98:1370-1375). The process of post-translational modification of glutamic residues to form Gla is Vitamin K-dependent carboxylation. Proteins which contain Gla include plasma proteins involved in blood coagulation. These proteins are prothrombin, proteins C, S, and Z, and coagulation factors VII, IX, and X. Osteocalcin (bone-Gla protein, BGP) and matrix Gla-protein (MGP) also contain Gla (Friedman, P.A., and C.T. Przysiecki (1987) Int. J. Biochem. 19:1-7; C. Vermeer (1990) Biochem. J. 266:625-636).

The *Drosophila* sp. gene *crossveinless 2* is characterized as having a putative signal or transmembrane sequence, and a partial Von Willebrand Factor D domain similar to those domains known to regulate the formation of intramolecular and intermolecular bonds and five cysteine-rich domains, known to bind BMP-like (bone morphogenetic proteins) ligands. These features suggest that *crossveinless 2* may act extracellularly or in the secretory pathway to directly potentiate ligand signaling and hence, involvement in the BMP-like signaling pathway known to play a role in vein specification (Conley, C.A. et al., (2000) Development 127:3947-3959). The dorsal-ventral patterning in both vertebrate and *Drosophila* embryos requires a conserved system of extracellular proteins to generate a positional informational gradient.

Immunoglobulins

Antigen recognition molecules are key players in the sophisticated and complex immune systems which all vertebrates have developed to provide protection from viral, bacterial, fungal, and parasitic infections. A key feature of the immune system is its ability to distinguish foreign molecules, or antigens, from "self" molecules. Most cell surface and soluble molecules that mediate functions such as recognition, adhesion or binding have evolved from a common evolutionary precursor (i.e., these proteins have structural homology). A number of molecules outside the immune system that have similar functions are also derived from this same evolutionary precursor. This ability is mediated primarily by secreted and transmembrane proteins expressed by leukocytes (white blood cells) such as lymphocytes, granulocytes, and monocytes. Most of these proteins belong to the immunoglobulin (Ig) superfamily, members of which contain one or more repeats of a conserved structural domain. This Ig domain is comprised of antiparallel β sheets joined by a disulfide bond in

an arrangement called the Ig fold. The criteria for a protein to be a member of the Ig superfamily is to have one or more Ig domains, which are regions of 70-110 amino acid residues in length homologous to either Ig variable-like (V) or Ig constant-like (C) domains. Members of the Ig superfamily include antibodies (Ab), T cell receptors (TCRs), class I and II major histocompatibility (MHC) proteins and immune cell-specific surface markers such as the "cluster of differentiation" or CD antigens, CD2, CD3, CD4, CD8, poly-Ig receptors, Fc receptors, neural cell-adhesion molecule (NCAM) and platelet-derived growth factor receptor (PDGFR). These antigens have been identified using systematic, monoclonal antibody (mAb)-based "shot gun" techniques. These techniques have resulted in the production of hundreds of mAbs directed against unknown cell surface leukocytic antigens. These antigens have been grouped into "clusters of differentiation" based on common immunocytochemical localization patterns in various differentiated and undifferentiated leukocytic cell types. Antigens in a given cluster are presumed to identify a single cell surface protein and are assigned a "cluster of differentiation" or "CD" designation. Some of the genes encoding proteins identified by CD antigens have been cloned and verified by standard molecular biology techniques. CD antigens have been characterized as both transmembrane proteins and cell surface proteins anchored to the plasma membrane via covalent attachment to fatty acid-containing glycolipids such as glycosylphosphatidylinositol (GPI). (Reviewed in Barclay, A. N. et al. (1995) The Leucocyte Antigen Facts Book, Academic Press, San Diego, CA, pp. 17-20.)

Ig domains (V and C) are regions of conserved amino acid residues that give a polypeptide a globular tertiary structure called an immunoglobulin (or antibody) fold, which consists of two approximately parallel layers of β -sheets. Conserved cysteine residues form an intrachain disulfide-bonded loop, 55-75 amino acid residues in length, which connects the two layers of the β -sheets. Each β -sheet has three or four anti-parallel β -strands of 5-10 amino acid residues. Hydrophobic and hydrophilic interactions of amino acid residues within the β -strands stabilize the Ig fold (hydrophobic on inward facing amino acid residues and hydrophilic on the amino acid residues in the outward facing portion of the strands). A V domain consists of a longer polypeptide than a C domain, with an additional pair of β -strands in the Ig fold.

A consistent feature of Ig superfamily genes is that each sequence of an Ig domain is encoded by a single exon. It is possible that the superfamily evolved from a gene coding for a single Ig domain involved in mediating cell-cell interactions. New members of the superfamily then arose by exon and gene duplications. Modern Ig superfamily proteins contain different numbers of V and/or C domains. Another evolutionary feature of this superfamily is the ability to undergo DNA rearrangements, a unique feature retained by the antigen receptor members of the family.

Many members of the Ig superfamily are integral plasma membrane proteins with extracellular Ig domains. The hydrophobic amino acid residues of their transmembrane domains and

their cytoplasmic tails are very diverse, with little or no homology among Ig family members or to known signal-transducing structures. There are exceptions to this general superfamily description. For example, the cytoplasmic tail of PDGFR has tyrosine kinase activity. In addition Thy-1 is a glycoprotein found on thymocytes and T cells. This protein has no cytoplasmic tail, but is instead
5 attached to the plasma membrane by a covalent glycosphosphatidylinositol linkage.

Another common feature of many Ig superfamily proteins is the interactions between Ig domains which are essential for the function of these molecules. Interactions between Ig domains of a multimeric protein can be either homophilic or heterophilic (i.e., between the same or different Ig domains). Antibodies are multimeric proteins which have both homophilic and heterophilic
10 interactions between Ig domains. Pairing of constant regions of heavy chains forms the Fc region of an antibody and pairing of variable regions of light and heavy chains form the antigen binding site of an antibody. Heterophilic interactions also occur between Ig domains of different molecules. These interactions provide adhesion between cells for significant cell-cell interactions in the immune system and in the developing and mature nervous system. (Reviewed in Abbas, A.K. et al. (1991) Cellular
15 and Molecular Immunology, W.B. Saunders Company, Philadelphia, PA, pp.142-145.)

Antibodies

MHC proteins are cell surface markers that bind to and present foreign antigens to T cells. MHC molecules are classified as either class I or class II. Class I MHC molecules (MHC I) are expressed on the surface of almost all cells and are involved in the presentation of antigen to
20 cytotoxic T cells. For example, a cell infected with virus will degrade intracellular viral proteins and express the protein fragments bound to MHC I molecules on the cell surface. The MHC I/antigen complex is recognized by cytotoxic T-cells which destroy the infected cell and the virus within. Class II MHC molecules are expressed primarily on specialized antigen-presenting cells of the immune system, such as B-cells and macrophages. These cells ingest foreign proteins from the
25 extracellular fluid and express MHC II/antigen complex on the cell surface. This complex activates helper T-cells, which then secrete cytokines and other factors that stimulate the immune response. MHC molecules also play an important role in organ rejection following transplantation. Rejection occurs when the recipient's T-cells respond to foreign MHC molecules on the transplanted organ in the same way as to self MHC molecules bound to foreign antigen. (Reviewed in Alberts, B. et al.
30 (1994) Molecular Biology of the Cell, Garland Publishing, New York, NY, pp. 1229-1246

Antibodies are multimeric members of the Ig superfamily which are either expressed on the surface of B-cells or secreted by B-cells into the circulation. Antibodies bind and neutralize foreign antigens in the blood and other extracellular fluids. The prototypical antibody is a tetramer consisting of two identical heavy polypeptide chains (H-chains) and two identical light polypeptide chains (L-
35 chains) interlinked by disulfide bonds. This arrangement confers the characteristic Y-shape to

antibody molecules. Antibodies are classified based on their H-chain composition. The five antibody classes, IgA, IgD, IgE, IgG and IgM, are defined by the α , δ , ϵ , γ , and μ H-chain types. There are two types of L-chains, κ and λ , either of which may associate as a pair with any H-chain pair. IgG, the most common class of antibody found in the circulation, is tetrameric, while the other classes of antibodies are generally variants or multimers of this basic structure.

H-chains and L-chains each contain an N-terminal variable region and a C-terminal constant region. The constant region consists of about 110 amino acids in L-chains and about 330 or 440 amino acids in H-chains. The amino acid sequence of the constant region is nearly identical among H- or L-chains of a particular class. The variable region consists of about 110 amino acids in both H- and L-chains. However, the amino acid sequence of the variable region differs among H- or L-chains of a particular class. Within each H- or L-chain variable region are three hypervariable regions of extensive sequence diversity, each consisting of about 5 to 10 amino acids. In the antibody molecule, the H- and L-chain hypervariable regions come together to form the antigen recognition site. (Reviewed in Alberts, B. et al. *supra*, pp. 1206-1213 and 1216-1217.)

Both H-chains and L-chains contain the repeated Ig domains of members of the Ig superfamily. For example, a typical H-chain contains four Ig domains, three of which occur within the constant region and one of which occurs within the variable region and contributes to the formation of the antigen recognition site. Likewise, a typical L-chain contains two Ig domains, one of which occurs within the constant region and one of which occurs within the variable region.

The immune system is capable of recognizing and responding to any foreign molecule that enters the body. Therefore, the immune system must be armed with a full repertoire of antibodies against all potential antigens. Such antibody diversity is generated by somatic rearrangement of gene segments encoding variable and constant regions. These gene segments are joined together by site-specific recombination which occurs between highly conserved DNA sequences that flank each gene segment. Because there are hundreds of different gene segments, millions of unique genes can be generated combinatorially. In addition, imprecise joining of these segments and an unusually high rate of somatic mutation within these segments further contribute to the generation of a diverse antibody population.

The discovery of new secreted proteins, and the polynucleotides encoding them, satisfies a need in the art by providing new compositions which are useful in the diagnosis, prevention, and treatment of cell proliferative, autoimmune/inflammatory, cardiovascular, neurological, and developmental disorders, and in the assessment of the effects of exogenous compounds on the expression of nucleic acid and amino acid sequences of secreted proteins.

35

SUMMARY OF THE INVENTION

The invention features purified polypeptides, secreted proteins, referred to collectively as "SECP" and individually as "SECP-1," "SECP-2," "SECP-3," "SECP-4," "SECP-5," "SECP-6," "SECP-7," "SECP-8," "SECP-9," "SECP-10," "SECP-11," "SECP-12," "SECP-13," "SECP-14," "SECP-15," "SECP-16," "SECP-17," "SECP-18," "SECP-19," "SECP-20," "SECP-21," "SECP-22,"
5 "SECP-23," "SECP-24," "SECP-25," "SECP-26," "SECP-27," "SECP-28," "SECP-29," "SECP-30," "SECP-31," "SECP-32," "SECP-33," "SECP-34," "SECP-35," "SECP-36," "SECP-37," "SECP-38," "SECP-39," "SECP-40," "SECP-41," "SECP-42," "SECP-43," "SECP-44," "SECP-45," "SECP-46," "SECP-47," "SECP-48," "SECP-49," "SECP-50," "SECP-51," "SECP-52," "SECP-53," "SECP-54," "SECP-55," "SECP-56," "SECP-57," "SECP-58," "SECP-59," "SECP-60," "SECP-61," "SECP-62,"
10 "SECP-63," "SECP-64," "SECP-65," "SECP-66," and "SECP-67." In one aspect, the invention provides an isolated polypeptide selected from the group consisting of a) a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-67, b) a polypeptide comprising a naturally occurring amino acid sequence at least 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-67, c) a biologically active fragment of
15 a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-67, and d) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-67. In one alternative, the invention provides an isolated polypeptide comprising the amino acid sequence of SEQ ID NO:1-67.

The invention further provides an isolated polynucleotide encoding a polypeptide selected
20 from the group consisting of a) a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-67, b) a polypeptide comprising a naturally occurring amino acid sequence at least 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-67, c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-67, and d) an immunogenic fragment of a
25 polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-67. In one alternative, the polynucleotide encodes a polypeptide selected from the group consisting of SEQ ID NO:1-67. In another alternative, the polynucleotide is selected from the group consisting of SEQ ID NO:68-134.

Additionally, the invention provides a recombinant polynucleotide comprising a promoter
30 sequence operably linked to a polynucleotide encoding a polypeptide selected from the group consisting of a) a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-67, b) a polypeptide comprising a naturally occurring amino acid sequence at least 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-67, c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group
35 consisting of SEQ ID NO:1-67, and d) an immunogenic fragment of a polypeptide having an amino

acid sequence selected from the group consisting of SEQ ID NO:1-67. In one alternative, the invention provides a cell transformed with the recombinant polynucleotide. In another alternative, the invention provides a transgenic organism comprising the recombinant polynucleotide.

The invention also provides a method for producing a polypeptide selected from the group
5 consisting of a) a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-67, b) a polypeptide comprising a naturally occurring amino acid sequence at least 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-67, c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-67, and d) an immunogenic fragment of a polypeptide having an amino
10 acid sequence selected from the group consisting of SEQ ID NO:1-67. The method comprises a) culturing a cell under conditions suitable for expression of the polypeptide, wherein said cell is transformed with a recombinant polynucleotide comprising a promoter sequence operably linked to a polynucleotide encoding the polypeptide, and b) recovering the polypeptide so expressed.

Additionally, the invention provides an isolated antibody which specifically binds to a
15 polypeptide selected from the group consisting of a) a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-67, b) a polypeptide comprising a naturally occurring amino acid sequence at least 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-67, c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-67, and d) an
20 immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-67.

The invention further provides an isolated polynucleotide selected from the group consisting of a) a polynucleotide comprising a polynucleotide sequence selected from the group consisting of SEQ ID NO:68-134, b) a polynucleotide comprising a naturally occurring polynucleotide sequence at
25 least 90% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NO:68-134, c) a polynucleotide complementary to the polynucleotide of a), d) a polynucleotide complementary to the polynucleotide of b), and e) an RNA equivalent of a)-d). In one alternative, the polynucleotide comprises at least 60 contiguous nucleotides.

Additionally, the invention provides a method for detecting a target polynucleotide in a
30 sample, said target polynucleotide having a sequence of a polynucleotide selected from the group consisting of a) a polynucleotide comprising a polynucleotide sequence selected from the group consisting of SEQ ID NO:68-134, b) a polynucleotide comprising a naturally occurring polynucleotide sequence at least 90% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NO:68-134, c) a polynucleotide complementary to the polynucleotide of a), d) a
35 polynucleotide complementary to the polynucleotide of b), and e) an RNA equivalent of a)-d). The

method comprises a) hybridizing the sample with a probe comprising at least 20 contiguous nucleotides comprising a sequence complementary to said target polynucleotide in the sample, and which probe specifically hybridizes to said target polynucleotide, under conditions whereby a hybridization complex is formed between said probe and said target polynucleotide or fragments thereof, and b) detecting the presence or absence of said hybridization complex, and optionally, if present, the amount thereof. In one alternative, the probe comprises at least 60 contiguous nucleotides.

The invention further provides a method for detecting a target polynucleotide in a sample, said target polynucleotide having a sequence of a polynucleotide selected from the group consisting of a) a polynucleotide comprising a polynucleotide sequence selected from the group consisting of SEQ ID NO:68-134, b) a polynucleotide comprising a naturally occurring polynucleotide sequence at least 90% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NO:68-134, c) a polynucleotide complementary to the polynucleotide of a), d) a polynucleotide complementary to the polynucleotide of b), and e) an RNA equivalent of a)-d). The method comprises a) amplifying said target polynucleotide or fragment thereof using polymerase chain reaction amplification, and b) detecting the presence or absence of said amplified target polynucleotide or fragment thereof, and, optionally, if present, the amount thereof.

The invention further provides a composition comprising an effective amount of a polypeptide selected from the group consisting of a) a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-67, b) a polypeptide comprising a naturally occurring amino acid sequence at least 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-67, c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-67, and d) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-67, and a pharmaceutically acceptable excipient. In one embodiment, the composition comprises an amino acid sequence selected from the group consisting of SEQ ID NO:1-67. The invention additionally provides a method of treating a disease or condition associated with decreased expression of functional SECP, comprising administering to a patient in need of such treatment the composition.

The invention also provides a method for screening a compound for effectiveness as an agonist of a polypeptide selected from the group consisting of a) a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-67, b) a polypeptide comprising a naturally occurring amino acid sequence at least 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-67, c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-67, and d) an

immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-67. The method comprises a) exposing a sample comprising the polypeptide to a compound, and b) detecting agonist activity in the sample. In one alternative, the invention provides a composition comprising an agonist compound identified by the method and a pharmaceutically acceptable excipient. In another alternative, the invention provides a method of treating a disease or condition associated with decreased expression of functional SECP, comprising administering to a patient in need of such treatment the composition.

Additionally, the invention provides a method for screening a compound for effectiveness as an antagonist of a polypeptide selected from the group consisting of a) a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-67, b) a polypeptide comprising a naturally occurring amino acid sequence at least 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-67, c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-67, and d) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-67. The method comprises a) exposing a sample comprising the polypeptide to a compound, and b) detecting antagonist activity in the sample. In one alternative, the invention provides a composition comprising an antagonist compound identified by the method and a pharmaceutically acceptable excipient. In another alternative, the invention provides a method of treating a disease or condition associated with overexpression of functional SECP, comprising administering to a patient in need of such treatment the composition.

The invention further provides a method of screening for a compound that specifically binds to a polypeptide selected from the group consisting of a) a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-67, b) a polypeptide comprising a naturally occurring amino acid sequence at least 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-67, c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-67, and d) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-67. The method comprises a) combining the polypeptide with at least one test compound under suitable conditions, and b) detecting binding of the polypeptide to the test compound, thereby identifying a compound that specifically binds to the polypeptide.

The invention further provides a method of screening for a compound that modulates the activity of a polypeptide selected from the group consisting of a) a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-67, b) a polypeptide comprising a naturally occurring amino acid sequence at least 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-67, c) a biologically active fragment of a polypeptide

having an amino acid sequence selected from the group consisting of SEQ ID NO:1-67, and d) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-67. The method comprises a) combining the polypeptide with at least one test compound under conditions permissive for the activity of the polypeptide, b) assessing the activity of the polypeptide in the presence of the test compound, and c) comparing the activity of the polypeptide in the presence of the test compound with the activity of the polypeptide in the absence of the test compound, wherein a change in the activity of the polypeptide in the presence of the test compound is indicative of a compound that modulates the activity of the polypeptide.

The invention further provides a method for screening a compound for effectiveness in altering expression of a target polynucleotide, wherein said target polynucleotide comprises a polynucleotide sequence selected from the group consisting of SEQ ID NO:68-134, the method comprising a) exposing a sample comprising the target polynucleotide to a compound, and b) detecting altered expression of the target polynucleotide.

The invention further provides a method for assessing toxicity of a test compound, said method comprising a) treating a biological sample containing nucleic acids with the test compound; b) hybridizing the nucleic acids of the treated biological sample with a probe comprising at least 20 contiguous nucleotides of a polynucleotide selected from the group consisting of i) a polynucleotide comprising a polynucleotide sequence selected from the group consisting of SEQ ID NO:68-134, ii) a polynucleotide comprising a naturally occurring polynucleotide sequence at least 90% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NO:68-134, iii) a polynucleotide having a sequence complementary to i), iv) a polynucleotide complementary to the polynucleotide of ii), and v) an RNA equivalent of i)-iv). Hybridization occurs under conditions whereby a specific hybridization complex is formed between said probe and a target polynucleotide in the biological sample, said target polynucleotide selected from the group consisting of i) a polynucleotide comprising a polynucleotide sequence selected from the group consisting of SEQ ID NO:68-134, ii) a polynucleotide comprising a naturally occurring polynucleotide sequence at least 90% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NO:68-134, iii) a polynucleotide complementary to the polynucleotide of i), iv) a polynucleotide complementary to the polynucleotide of ii), and v) an RNA equivalent of i)-iv). Alternatively, the target polynucleotide comprises a fragment of a polynucleotide sequence selected from the group consisting of i)-v) above; c) quantifying the amount of hybridization complex; and d) comparing the amount of hybridization complex in the treated biological sample with the amount of hybridization complex in an untreated biological sample, wherein a difference in the amount of hybridization complex in the treated biological sample is indicative of toxicity of the test compound.

35

BRIEF DESCRIPTION OF THE TABLES

Table 1 summarizes the nomenclature for the full length polynucleotide and polypeptide sequences of the present invention.

Table 2 shows the GenBank identification number and annotation of the nearest GenBank homolog for polypeptides of the invention. The probability score for the match between each polypeptide and its GenBank homolog is also shown.

Table 3 shows structural features of polypeptide sequences of the invention, including predicted motifs and domains, along with the methods, algorithms, and searchable databases used for analysis of the polypeptides.

Table 4 lists the cDNA and/or genomic DNA fragments which were used to assemble polynucleotide sequences of the invention, along with selected fragments of the polynucleotide sequences.

Table 5 shows the representative cDNA library for polynucleotides of the invention.

Table 6 provides an appendix which describes the tissues and vectors used for construction of the cDNA libraries shown in Table 5.

Table 7 shows the tools, programs, and algorithms used to analyze the polynucleotides and polypeptides of the invention, along with applicable descriptions, references, and threshold parameters.

DESCRIPTION OF THE INVENTION

Before the present proteins, nucleotide sequences, and methods are described, it is understood that this invention is not limited to the particular machines, materials and methods described, as these may vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments only, and is not intended to limit the scope of the present invention which will be limited only by the appended claims.

It must be noted that as used herein and in the appended claims, the singular forms "a," "an," and "the" include plural reference unless the context clearly dictates otherwise. Thus, for example, a reference to "a host cell" includes a plurality of such host cells, and a reference to "an antibody" is a reference to one or more antibodies and equivalents thereof known to those skilled in the art, and so forth.

Unless defined otherwise, all technical and scientific terms used herein have the same meanings as commonly understood by one of ordinary skill in the art to which this invention belongs. Although any machines, materials, and methods similar or equivalent to those described herein can be used to practice or test the present invention, the preferred machines, materials and methods are now described. All publications mentioned herein are cited for the purpose of describing and disclosing

the cell lines, protocols, reagents and vectors which are reported in the publications and which might be used in connection with the invention. Nothing herein is to be construed as an admission that the invention is not entitled to antedate such disclosure by virtue of prior invention.

DEFINITIONS

5 “SECP” refers to the amino acid sequences of substantially purified SECP obtained from any species, particularly a mammalian species, including bovine, ovine, porcine, murine, equine, and human, and from any source, whether natural, synthetic, semi-synthetic, or recombinant.

 The term “agonist” refers to a molecule which intensifies or mimics the biological activity of SECP. Agonists may include proteins, nucleic acids, carbohydrates, small molecules, or any other
10 compound or composition which modulates the activity of SECP either by directly interacting with SECP or by acting on components of the biological pathway in which SECP participates.

 An “allelic variant” is an alternative form of the gene encoding SECP. Allelic variants may result from at least one mutation in the nucleic acid sequence and may result in altered mRNAs or in polypeptides whose structure or function may or may not be altered. A gene may have none, one, or
15 many allelic variants of its naturally occurring form. Common mutational changes which give rise to allelic variants are generally ascribed to natural deletions, additions, or substitutions of nucleotides. Each of these types of changes may occur alone, or in combination with the others, one or more times in a given sequence.

 “Altered” nucleic acid sequences encoding SECP include those sequences with deletions,
20 insertions, or substitutions of different nucleotides, resulting in a polypeptide the same as SECP or a polypeptide with at least one functional characteristic of SECP. Included within this definition are polymorphisms which may or may not be readily detectable using a particular oligonucleotide probe of the polynucleotide encoding SECP, and improper or unexpected hybridization to allelic variants, with a locus other than the normal chromosomal locus for the polynucleotide sequence encoding
25 SECP. The encoded protein may also be “altered,” and may contain deletions, insertions, or substitutions of amino acid residues which produce a silent change and result in a functionally equivalent SECP. Deliberate amino acid substitutions may be made on the basis of similarity in polarity, charge, solubility, hydrophobicity, hydrophilicity, and/or the amphipathic nature of the residues, as long as the biological or immunological activity of SECP is retained. For example,
30 negatively charged amino acids may include aspartic acid and glutamic acid, and positively charged amino acids may include lysine and arginine. Amino acids with uncharged polar side chains having similar hydrophilicity values may include: asparagine and glutamine; and serine and threonine. Amino acids with uncharged side chains having similar hydrophilicity values may include: leucine, isoleucine, and valine; glycine and alanine; and phenylalanine and tyrosine.

35 The terms “amino acid” and “amino acid sequence” refer to an oligopeptide, peptide,

polypeptide, or protein sequence, or a fragment of any of these, and to naturally occurring or synthetic molecules. Where "amino acid sequence" is recited to refer to a sequence of a naturally occurring protein molecule, "amino acid sequence" and like terms are not meant to limit the amino acid sequence to the complete native amino acid sequence associated with the recited protein molecule.

5 "Amplification" relates to the production of additional copies of a nucleic acid sequence. Amplification is generally carried out using polymerase chain reaction (PCR) technologies well known in the art.

The term "antagonist" refers to a molecule which inhibits or attenuates the biological activity of SECP. Antagonists may include proteins such as antibodies, nucleic acids, carbohydrates, small
10 molecules, or any other compound or composition which modulates the activity of SECP either by directly interacting with SECP or by acting on components of the biological pathway in which SECP participates.

The term "antibody" refers to intact immunoglobulin molecules as well as to fragments thereof, such as Fab, F(ab')₂, and Fv fragments, which are capable of binding an epitopic determinant.
15 Antibodies that bind SECP polypeptides can be prepared using intact polypeptides or using fragments containing small peptides of interest as the immunizing antigen. The polypeptide or oligopeptide used to immunize an animal (e.g., a mouse, a rat, or a rabbit) can be derived from the translation of RNA, or synthesized chemically, and can be conjugated to a carrier protein if desired. Commonly used carriers that are chemically coupled to peptides include bovine serum albumin, thyroglobulin,
20 and keyhole limpet hemocyanin (KLH). The coupled peptide is then used to immunize the animal.

The term "antigenic determinant" refers to that region of a molecule (i.e., an epitope) that makes contact with a particular antibody. When a protein or a fragment of a protein is used to immunize a host animal, numerous regions of the protein may induce the production of antibodies which bind specifically to antigenic determinants (particular regions or three-dimensional structures
25 on the protein). An antigenic determinant may compete with the intact antigen (i.e., the immunogen used to elicit the immune response) for binding to an antibody.

The term "aptamer" refers to a nucleic acid or oligonucleotide molecule that binds to a specific molecular target. Aptamers are derived from an in vitro evolutionary process (e.g., SELEX (Systematic Evolution of Ligands by EXponential Enrichment), described in U.S. Patent No.
30 5,270,163), which selects for target-specific aptamer sequences from large combinatorial libraries. Aptamer compositions may be double-stranded or single-stranded, and may include deoxyribonucleotides, ribonucleotides, nucleotide derivatives, or other nucleotide-like molecules. The nucleotide components of an aptamer may have modified sugar groups (e.g., the 2'-OH group of a ribonucleotide may be replaced by 2'-F or 2'-NH₂), which may improve a desired property, e.g.,
35 resistance to nucleases or longer lifetime in blood. Aptamers may be conjugated to other molecules,

e.g., a high molecular weight carrier to slow clearance of the aptamer from the circulatory system. Aptamers may be specifically cross-linked to their cognate ligands, e.g., by photo-activation of a cross-linker. (See, e.g., Brody, E.N. and L. Gold (2000) J. Biotechnol. 74:5-13.)

The term "intramer" refers to an aptamer which is expressed in vivo. For example, a vaccinia virus-based RNA expression system has been used to express specific RNA aptamers at high levels in the cytoplasm of leukocytes (Blind, M. et al. (1999) Proc. Natl Acad. Sci. USA 96:3606-3610).

The term "spiegelmer" refers to an aptamer which includes L-DNA, L-RNA, or other left-handed nucleotide derivatives or nucleotide-like molecules. Aptamers containing left-handed nucleotides are resistant to degradation by naturally occurring enzymes, which normally act on substrates containing right-handed nucleotides.

The term "antisense" refers to any composition capable of base-pairing with the "sense" (coding) strand of a specific nucleic acid sequence. Antisense compositions may include DNA; RNA; peptide nucleic acid (PNA); oligonucleotides having modified backbone linkages such as phosphorothioates, methylphosphonates, or benzylphosphonates; oligonucleotides having modified sugar groups such as 2'-methoxyethyl sugars or 2'-methoxyethoxy sugars; or oligonucleotides having modified bases such as 5-methyl cytosine, 2'-deoxyuracil, or 7-deaza-2'-deoxyguanosine. Antisense molecules may be produced by any method including chemical synthesis or transcription. Once introduced into a cell, the complementary antisense molecule base-pairs with a naturally occurring nucleic acid sequence produced by the cell to form duplexes which block either transcription or translation. The designation "negative" or "minus" can refer to the antisense strand, and the designation "positive" or "plus" can refer to the sense strand of a reference DNA molecule.

The term "biologically active" refers to a protein having structural, regulatory, or biochemical functions of a naturally occurring molecule. Likewise, "immunologically active" or "immunogenic" refers to the capability of the natural, recombinant, or synthetic SECP, or of any oligopeptide thereof, to induce a specific immune response in appropriate animals or cells and to bind with specific antibodies.

"Complementary" describes the relationship between two single-stranded nucleic acid sequences that anneal by base-pairing. For example, 5'-AGT-3' pairs with its complement, 3'-TCA-5'. In an alternative example, SEQ ID NO:135 and SEQ ID NO:136 pair with their complements, SEQ ID NO:114 and SEQ ID NO:116, respectively.

A "composition comprising a given polynucleotide sequence" and a "composition comprising a given amino acid sequence" refer broadly to any composition containing the given polynucleotide or amino acid sequence. The composition may comprise a dry formulation or an aqueous solution. Compositions comprising polynucleotide sequences encoding SECP or fragments of SECP may be employed as hybridization probes. The probes may be stored in freeze-dried form and may be

associated with a stabilizing agent such as a carbohydrate. In hybridizations, the probe may be deployed in an aqueous solution containing salts (e.g., NaCl), detergents (e.g., sodium dodecyl sulfate; SDS), and other components (e.g., Denhardt's solution, dry milk, salmon sperm DNA, etc.).

"Consensus sequence" refers to a nucleic acid sequence which has been subjected to repeated DNA sequence analysis to resolve uncalled bases, extended using the XL-PCR kit (Applied Biosystems, Foster City CA) in the 5' and/or the 3' direction, and resequenced, or which has been assembled from one or more overlapping cDNA, EST, or genomic DNA fragments using a computer program for fragment assembly, such as the GELVIEW fragment assembly system (GCG, Madison WI) or Phrap (University of Washington, Seattle WA). Some sequences have been both extended and assembled to produce the consensus sequence.

"Conservative amino acid substitutions" are those substitutions that are predicted to least interfere with the properties of the original protein, i.e., the structure and especially the function of the protein is conserved and not significantly changed by such substitutions. The table below shows amino acids which may be substituted for an original amino acid in a protein and which are regarded as conservative amino acid substitutions.

	Original Residue	Conservative Substitution
	Ala	Gly, Ser
	Arg	His, Lys
	Asn	Asp, Gln, His
20	Asp	Asn, Glu
	Cys	Ala, Ser
	Gln	Asn, Glu, His
	Glu	Asp, Gln, His
	Gly	Ala
25	His	Asn, Arg, Gln, Glu
	Ile	Leu, Val
	Leu	Ile, Val
	Lys	Arg, Gln, Glu
	Met	Leu, Ile
30	Phe	His, Met, Leu, Trp, Tyr
	Ser	Cys, Thr
	Thr	Ser, Val
	Trp	Phe, Tyr
	Tyr	His, Phe, Trp
35	Val	Ile, Leu, Thr

Conservative amino acid substitutions generally maintain (a) the structure of the polypeptide backbone in the area of the substitution, for example, as a beta sheet or alpha helical conformation, (b) the charge or hydrophobicity of the molecule at the site of the substitution, and/or (c) the bulk of the side chain.

A "deletion" refers to a change in the amino acid or nucleotide sequence that results in the absence of one or more amino acid residues or nucleotides.

The term "derivative" refers to a chemically modified polynucleotide or polypeptide. Chemical modifications of a polynucleotide can include, for example, replacement of hydrogen by an alkyl, acyl, hydroxyl, or amino group. A derivative polynucleotide encodes a polypeptide which retains at least one biological or immunological function of the natural molecule. A derivative
5 polypeptide is one modified by glycosylation, pegylation, or any similar process that retains at least one biological or immunological function of the polypeptide from which it was derived.

A "detectable label" refers to a reporter molecule or enzyme that is capable of generating a measurable signal and is covalently or noncovalently joined to a polynucleotide or polypeptide.

"Differential expression" refers to increased or upregulated; or decreased, downregulated, or
10 absent gene or protein expression, determined by comparing at least two different samples. Such comparisons may be carried out between, for example, a treated and an untreated sample, or a diseased and a normal sample.

"Exon shuffling" refers to the recombination of different coding regions (exons). Since an exon may represent a structural or functional domain of the encoded protein, new proteins may be
15 assembled through the novel reassortment of stable substructures, thus allowing acceleration of the evolution of new protein functions.

A "fragment" is a unique portion of SECP or the polynucleotide encoding SECP which is identical in sequence to but shorter in length than the parent sequence. A fragment may comprise up to the entire length of the defined sequence, minus one nucleotide/amino acid residue. For example, a
20 fragment may comprise from 5 to 1000 contiguous nucleotides or amino acid residues. A fragment used as a probe, primer, antigen, therapeutic molecule, or for other purposes, may be at least 5, 10, 15, 16, 20, 25, 30, 40, 50, 60, 75, 100, 150, 250 or at least 500 contiguous nucleotides or amino acid residues in length. Fragments may be preferentially selected from certain regions of a molecule. For example, a polypeptide fragment may comprise a certain length of contiguous amino acids selected
25 from the first 250 or 500 amino acids (or first 25% or 50%) of a polypeptide as shown in a certain defined sequence. Clearly these lengths are exemplary, and any length that is supported by the specification, including the Sequence Listing, tables, and figures, may be encompassed by the present embodiments.

A fragment of SEQ ID NO:68-134 comprises a region of unique polynucleotide sequence that
30 specifically identifies SEQ ID NO:68-134, for example, as distinct from any other sequence in the genome from which the fragment was obtained. A fragment of SEQ ID NO:68-134 is useful, for example, in hybridization and amplification technologies and in analogous methods that distinguish SEQ ID NO:68-134 from related polynucleotide sequences. The precise length of a fragment of SEQ ID NO:68-134 and the region of SEQ ID NO:68-134 to which the fragment corresponds are routinely
35 determinable by one of ordinary skill in the art based on the intended purpose for the fragment.

A fragment of SEQ ID NO:1-67 is encoded by a fragment of SEQ ID NO:68-134. A fragment of SEQ ID NO:1-67 comprises a region of unique amino acid sequence that specifically identifies SEQ ID NO:1-67. For example, a fragment of SEQ ID NO:1-67 is useful as an immunogenic peptide for the development of antibodies that specifically recognize SEQ ID NO:1-67.

5 The precise length of a fragment of SEQ ID NO:1-67 and the region of SEQ ID NO:1-67 to which the fragment corresponds are routinely determinable by one of ordinary skill in the art based on the intended purpose for the fragment.

A "full length" polynucleotide sequence is one containing at least a translation initiation codon (e.g., methionine) followed by an open reading frame and a translation termination codon. A

10 "full length" polynucleotide sequence encodes a "full length" polypeptide sequence.

"Homology" refers to sequence similarity or, interchangeably, sequence identity, between two or more polynucleotide sequences or two or more polypeptide sequences.

The terms "percent identity" and "% identity," as applied to polynucleotide sequences, refer to the percentage of residue matches between at least two polynucleotide sequences aligned using a

15 standardized algorithm. Such an algorithm may insert, in a standardized and reproducible way, gaps in the sequences being compared in order to optimize alignment between two sequences, and therefore achieve a more meaningful comparison of the two sequences.

Percent identity between polynucleotide sequences may be determined using the default parameters of the CLUSTAL V algorithm as incorporated into the MEGALIGN version 3.12e

20 sequence alignment program. This program is part of the LASERGENE software package, a suite of molecular biological analysis programs (DNASTAR, Madison WI). CLUSTAL V is described in Higgins, D.G. and P.M. Sharp (1989) CABIOS 5:151-153 and in Higgins, D.G. et al. (1992) CABIOS 8:189-191. For pairwise alignments of polynucleotide sequences, the default parameters are set as follows: Ktuple=2, gap penalty=5, window=4, and "diagonals saved"=4. The "weighted" residue

25 weight table is selected as the default. Percent identity is reported by CLUSTAL V as the "percent similarity" between aligned polynucleotide sequences.

Alternatively, a suite of commonly used and freely available sequence comparison algorithms is provided by the National Center for Biotechnology Information (NCBI) Basic Local Alignment Search Tool (BLAST) (Altschul, S.F. et al. (1990) J. Mol. Biol. 215:403-410), which is available

30 from several sources, including the NCBI, Bethesda, MD, and on the Internet at <http://www.ncbi.nlm.nih.gov/BLAST/>. The BLAST software suite includes various sequence analysis programs including "blastn," that is used to align a known polynucleotide sequence with other polynucleotide sequences from a variety of databases. Also available is a tool called "BLAST 2 Sequences" that is used for direct pairwise comparison of two nucleotide sequences. "BLAST 2

35 Sequences" can be accessed and used interactively at <http://www.ncbi.nlm.nih.gov/gorf/bl2.html>.

The "BLAST 2 Sequences" tool can be used for both blastn and blastp (discussed below). BLAST programs are commonly used with gap and other parameters set to default settings. For example, to compare two nucleotide sequences, one may use blastn with the "BLAST 2 Sequences" tool Version 2.0.12 (April-21-2000) set at default parameters. Such default parameters may be, for example:

5 *Matrix: BLOSUM62*
 Reward for match: 1
 Penalty for mismatch: -2
 Open Gap: 5 and Extension Gap: 2 penalties
 Gap x drop-off: 50
 10 *Expect: 10*
 Word Size: 11
 Filter: on

Percent identity may be measured over the length of an entire defined sequence, for example, as defined by a particular SEQ ID number, or may be measured over a shorter length, for example,
 15 over the length of a fragment taken from a larger, defined sequence, for instance, a fragment of at least 20, at least 30, at least 40, at least 50, at least 70, at least 100, or at least 200 contiguous nucleotides. Such lengths are exemplary only, and it is understood that any fragment length supported by the sequences shown herein, in the tables, figures, or Sequence Listing, may be used to describe a length over which percentage identity may be measured.

20 Nucleic acid sequences that do not show a high degree of identity may nevertheless encode similar amino acid sequences due to the degeneracy of the genetic code. It is understood that changes in a nucleic acid sequence can be made using this degeneracy to produce multiple nucleic acid sequences that all encode substantially the same protein.

 The phrases "percent identity" and "% identity," as applied to polypeptide sequences, refer to
 25 the percentage of residue matches between at least two polypeptide sequences aligned using a standardized algorithm. Methods of polypeptide sequence alignment are well-known. Some alignment methods take into account conservative amino acid substitutions. Such conservative substitutions, explained in more detail above, generally preserve the charge and hydrophobicity at the site of substitution, thus preserving the structure (and therefore function) of the polypeptide.

30 Percent identity between polypeptide sequences may be determined using the default parameters of the CLUSTAL V algorithm as incorporated into the MEGALIGN version 3.12e sequence alignment program (described and referenced above). For pairwise alignments of polypeptide sequences using CLUSTAL V, the default parameters are set as follows: Ktuple=1, gap penalty=3, window=5, and "diagonals saved"=5. The PAM250 matrix is selected as the default
 35 residue weight table. As with polynucleotide alignments, the percent identity is reported by

CLUSTAL V as the "percent similarity" between aligned polypeptide sequence pairs.

Alternatively the NCBI BLAST software suite may be used. For example, for a pairwise comparison of two polypeptide sequences, one may use the "BLAST 2 Sequences" tool Version 2.0.12 (April-21-2000) with blastp set at default parameters. Such default parameters may be, for example:

Matrix: BLOSUM62

Open Gap: 11 and Extension Gap: 1 penalties

Gap x drop-off: 50

Expect: 10

Word Size: 3

Filter: on

Percent identity may be measured over the length of an entire defined polypeptide sequence, for example, as defined by a particular SEQ ID number, or may be measured over a shorter length, for example, over the length of a fragment taken from a larger, defined polypeptide sequence, for instance, a fragment of at least 15, at least 20, at least 30, at least 40, at least 50, at least 70 or at least 150 contiguous residues. Such lengths are exemplary only, and it is understood that any fragment length supported by the sequences shown herein, in the tables, figures or Sequence Listing, may be used to describe a length over which percentage identity may be measured.

"Human artificial chromosomes" (HACs) are linear microchromosomes which may contain DNA sequences of about 6 kb to 10 Mb in size and which contain all of the elements required for chromosome replication, segregation and maintenance.

The term "humanized antibody" refers to an antibody molecule in which the amino acid sequence in the non-antigen binding regions has been altered so that the antibody more closely resembles a human antibody, and still retains its original binding ability.

"Hybridization" refers to the process by which a polynucleotide strand anneals with a complementary strand through base pairing under defined hybridization conditions. Specific hybridization is an indication that two nucleic acid sequences share a high degree of complementarity. Specific hybridization complexes form under permissive annealing conditions and remain hybridized after the "washing" step(s). The washing step(s) is particularly important in determining the stringency of the hybridization process, with more stringent conditions allowing less non-specific binding, i.e., binding between pairs of nucleic acid strands that are not perfectly matched. Permissive conditions for annealing of nucleic acid sequences are routinely determinable by one of ordinary skill in the art and may be consistent among hybridization experiments, whereas wash conditions may be varied among experiments to achieve the desired stringency, and therefore hybridization specificity.

Permissive annealing conditions occur, for example, at 68°C in the presence of about 6 x SSC, about

1% (w/v) SDS, and about 100 $\mu\text{g/ml}$ sheared, denatured salmon sperm DNA.

Generally, stringency of hybridization is expressed, in part, with reference to the temperature under which the wash step is carried out. Such wash temperatures are typically selected to be about 5°C to 20°C lower than the thermal melting point (T_m) for the specific sequence at a defined ionic strength and pH. The T_m is the temperature (under defined ionic strength and pH) at which 50% of the target sequence hybridizes to a perfectly matched probe. An equation for calculating T_m and conditions for nucleic acid hybridization are well known and can be found in Sambrook, J. et al. (1989) Molecular Cloning: A Laboratory Manual, 2nd ed., vol. 1-3, Cold Spring Harbor Press, Plainview NY; specifically see volume 2, chapter 9.

High stringency conditions for hybridization between polynucleotides of the present invention include wash conditions of 68°C in the presence of about 0.2 x SSC and about 0.1% SDS, for 1 hour. Alternatively, temperatures of about 65°C, 60°C, 55°C, or 42°C may be used. SSC concentration may be varied from about 0.1 to 2 x SSC, with SDS being present at about 0.1%. Typically, blocking reagents are used to block non-specific hybridization. Such blocking reagents include, for instance, sheared and denatured salmon sperm DNA at about 100-200 $\mu\text{g/ml}$. Organic solvent, such as formamide at a concentration of about 35-50% v/v, may also be used under particular circumstances, such as for RNA:DNA hybridizations. Useful variations on these wash conditions will be readily apparent to those of ordinary skill in the art. Hybridization, particularly under high stringency conditions, may be suggestive of evolutionary similarity between the nucleotides. Such similarity is strongly indicative of a similar role for the nucleotides and their encoded polypeptides.

The term "hybridization complex" refers to a complex formed between two nucleic acid sequences by virtue of the formation of hydrogen bonds between complementary bases. A hybridization complex may be formed in solution (e.g., C_0t or R_0t analysis) or formed between one nucleic acid sequence present in solution and another nucleic acid sequence immobilized on a solid support (e.g., paper, membranes, filters, chips, pins or glass slides, or any other appropriate substrate to which cells or their nucleic acids have been fixed).

The words "insertion" and "addition" refer to changes in an amino acid or nucleotide sequence resulting in the addition of one or more amino acid residues or nucleotides, respectively.

"Immune response" can refer to conditions associated with inflammation, trauma, immune disorders, or infectious or genetic disease, etc. These conditions can be characterized by expression of various factors, e.g., cytokines, chemokines, and other signaling molecules, which may affect cellular and systemic defense systems.

An "immunogenic fragment" is a polypeptide or oligopeptide fragment of SECP which is capable of eliciting an immune response when introduced into a living organism, for example, a mammal. The term "immunogenic fragment" also includes any polypeptide or oligopeptide fragment

of SECP which is useful in any of the antibody production methods disclosed herein or known in the art.

The term "microarray" refers to an arrangement of a plurality of polynucleotides, polypeptides, or other chemical compounds on a substrate.

5 The terms "element" and "array element" refer to a polynucleotide, polypeptide, or other chemical compound having a unique and defined position on a microarray.

The term "modulate" refers to a change in the activity of SECP. For example, modulation may cause an increase or a decrease in protein activity, binding characteristics, or any other biological, functional, or immunological properties of SECP.

10 The phrases "nucleic acid" and "nucleic acid sequence" refer to a nucleotide, oligonucleotide, polynucleotide, or any fragment thereof. These phrases also refer to DNA or RNA of genomic or synthetic origin which may be single-stranded or double-stranded and may represent the sense or the antisense strand, to peptide nucleic acid (PNA), or to any DNA-like or RNA-like material.

"Operably linked" refers to the situation in which a first nucleic acid sequence is placed in a
15 functional relationship with a second nucleic acid sequence. For instance, a promoter is operably linked to a coding sequence if the promoter affects the transcription or expression of the coding sequence. Operably linked DNA sequences may be in close proximity or contiguous and, where necessary to join two protein coding regions, in the same reading frame.

"Peptide nucleic acid" (PNA) refers to an antisense molecule or anti-gene agent which
20 comprises an oligonucleotide of at least about 5 nucleotides in length linked to a peptide backbone of amino acid residues ending in lysine. The terminal lysine confers solubility to the composition. PNAs preferentially bind complementary single stranded DNA or RNA and stop transcript elongation, and may be pegylated to extend their lifespan in the cell.

"Post-translational modification" of an SECP may involve lipidation, glycosylation,
25 phosphorylation, acetylation, racemization, proteolytic cleavage, and other modifications known in the art. These processes may occur synthetically or biochemically. Biochemical modifications will vary by cell type depending on the enzymatic milieu of SECP.

"Probe" refers to nucleic acid sequences encoding SECP, their complements, or fragments thereof, which are used to detect identical, allelic or related nucleic acid sequences. Probes are
30 isolated oligonucleotides or polynucleotides attached to a detectable label or reporter molecule. Typical labels include radioactive isotopes, ligands, chemiluminescent agents, and enzymes.

"Primers" are short nucleic acids, usually DNA oligonucleotides, which may be annealed to a target polynucleotide by complementary base-pairing. The primer may then be extended along the target DNA strand by a DNA polymerase enzyme. Primer pairs can be used for amplification (and
35 identification) of a nucleic acid sequence, e.g., by the polymerase chain reaction (PCR).

Probes and primers as used in the present invention typically comprise at least 15 contiguous nucleotides of a known sequence. In order to enhance specificity, longer probes and primers may also be employed, such as probes and primers that comprise at least 20, 25, 30, 40, 50, 60, 70, 80, 90, 100, or at least 150 consecutive nucleotides of the disclosed nucleic acid sequences. Probes and primers
5 may be considerably longer than these examples, and it is understood that any length supported by the specification, including the tables, figures, and Sequence Listing, may be used.

Methods for preparing and using probes and primers are described in the references, for example Sambrook, J. et al. (1989) Molecular Cloning: A Laboratory Manual, 2nd ed., vol. 1-3, Cold Spring Harbor Press, Plainview NY; Ausubel, F.M. et al. (1987) Current Protocols in Molecular
10 Biology, Greene Publ. Assoc. & Wiley-Intersciences, New York NY; Innis, M. et al. (1990) PCR Protocols, A Guide to Methods and Applications, Academic Press, San Diego CA. PCR primer pairs can be derived from a known sequence, for example, by using computer programs intended for that purpose such as Primer (Version 0.5, 1991, Whitehead Institute for Biomedical Research, Cambridge MA).

15 Oligonucleotides for use as primers are selected using software known in the art for such purpose. For example, OLIGO 4.06 software is useful for the selection of PCR primer pairs of up to 100 nucleotides each, and for the analysis of oligonucleotides and larger polynucleotides of up to 5,000 nucleotides from an input polynucleotide sequence of up to 32 kilobases. Similar primer selection programs have incorporated additional features for expanded capabilities. For example, the
20 PrimOU primer selection program (available to the public from the Genome Center at University of Texas South West Medical Center, Dallas TX) is capable of choosing specific primers from megabase sequences and is thus useful for designing primers on a genome-wide scope. The Primer3 primer selection program (available to the public from the Whitehead Institute/MIT Center for Genome Research, Cambridge MA) allows the user to input a "mispriming library," in which
25 sequences to avoid as primer binding sites are user-specified. Primer3 is useful, in particular, for the selection of oligonucleotides for microarrays. (The source code for the latter two primer selection programs may also be obtained from their respective sources and modified to meet the user's specific needs.) The PrimeGen program (available to the public from the UK Human Genome Mapping Project Resource Centre, Cambridge UK) designs primers based on multiple sequence alignments,
30 thereby allowing selection of primers that hybridize to either the most conserved or least conserved regions of aligned nucleic acid sequences. Hence, this program is useful for identification of both unique and conserved oligonucleotides and polynucleotide fragments. The oligonucleotides and polynucleotide fragments identified by any of the above selection methods are useful in hybridization technologies, for example, as PCR or sequencing primers, microarray elements, or specific probes to
35 identify fully or partially complementary polynucleotides in a sample of nucleic acids. Methods of

oligonucleotide selection are not limited to those described above.

A "recombinant nucleic acid" is a sequence that is not naturally occurring or has a sequence that is made by an artificial combination of two or more otherwise separated segments of sequence. This artificial combination is often accomplished by chemical synthesis or, more commonly, by the artificial manipulation of isolated segments of nucleic acids, e.g., by genetic engineering techniques such as those described in Sambrook, supra. The term recombinant includes nucleic acids that have been altered solely by addition, substitution, or deletion of a portion of the nucleic acid. Frequently, a recombinant nucleic acid may include a nucleic acid sequence operably linked to a promoter sequence. Such a recombinant nucleic acid may be part of a vector that is used, for example, to transform a cell.

Alternatively, such recombinant nucleic acids may be part of a viral vector, e.g., based on a vaccinia virus, that could be used to vaccinate a mammal wherein the recombinant nucleic acid is expressed, inducing a protective immunological response in the mammal.

A "regulatory element" refers to a nucleic acid sequence usually derived from untranslated regions of a gene and includes enhancers, promoters, introns, and 5' and 3' untranslated regions (UTRs). Regulatory elements interact with host or viral proteins which control transcription, translation, or RNA stability.

"Reporter molecules" are chemical or biochemical moieties used for labeling a nucleic acid, amino acid, or antibody. Reporter molecules include radionuclides; enzymes; fluorescent, chemiluminescent, or chromogenic agents; substrates; cofactors; inhibitors; magnetic particles; and other moieties known in the art.

An "RNA equivalent," in reference to a DNA sequence, is composed of the same linear sequence of nucleotides as the reference DNA sequence with the exception that all occurrences of the nitrogenous base thymine are replaced with uracil, and the sugar backbone is composed of ribose instead of deoxyribose.

The term "sample" is used in its broadest sense. A sample suspected of containing SECP, nucleic acids encoding SECP, or fragments thereof may comprise a bodily fluid; an extract from a cell, chromosome, organelle, or membrane isolated from a cell; a cell; genomic DNA, RNA, or cDNA, in solution or bound to a substrate; a tissue; a tissue print; etc.

The terms "specific binding" and "specifically binding" refer to that interaction between a protein or peptide and an agonist, an antibody, an antagonist, a small molecule, or any natural or synthetic binding composition. The interaction is dependent upon the presence of a particular structure of the protein, e.g., the antigenic determinant or epitope, recognized by the binding molecule. For example, if an antibody is specific for epitope "A," the presence of a polypeptide comprising the epitope A, or the presence of free unlabeled A, in a reaction containing free labeled A

and the antibody will reduce the amount of labeled A that binds to the antibody.

The term "substantially purified" refers to nucleic acid or amino acid sequences that are removed from their natural environment and are isolated or separated, and are at least 60% free, preferably at least 75% free, and most preferably at least 90% free from other components with which they are naturally associated.

A "substitution" refers to the replacement of one or more amino acid residues or nucleotides by different amino acid residues or nucleotides, respectively.

"Substrate" refers to any suitable rigid or semi-rigid support including membranes, filters, chips, slides, wafers, fibers, magnetic or nonmagnetic beads, gels, tubing, plates, polymers, microparticles and capillaries. The substrate can have a variety of surface forms, such as wells, trenches, pins, channels and pores, to which polynucleotides or polypeptides are bound.

A "transcript image" refers to the collective pattern of gene expression by a particular cell type or tissue under given conditions at a given time.

"Transformation" describes a process by which exogenous DNA is introduced into a recipient cell. Transformation may occur under natural or artificial conditions according to various methods well known in the art, and may rely on any known method for the insertion of foreign nucleic acid sequences into a prokaryotic or eukaryotic host cell. The method for transformation is selected based on the type of host cell being transformed and may include, but is not limited to, bacteriophage or viral infection, electroporation, heat shock, lipofection, and particle bombardment. The term "transformed cells" includes stably transformed cells in which the inserted DNA is capable of replication either as an autonomously replicating plasmid or as part of the host chromosome, as well as transiently transformed cells which express the inserted DNA or RNA for limited periods of time.

A "transgenic organism," as used herein, is any organism, including but not limited to animals and plants, in which one or more of the cells of the organism contains heterologous nucleic acid introduced by way of human intervention, such as by transgenic techniques well known in the art. The nucleic acid is introduced into the cell, directly or indirectly by introduction into a precursor of the cell, by way of deliberate genetic manipulation, such as by microinjection or by infection with a recombinant virus. The term genetic manipulation does not include classical cross-breeding, or in vitro fertilization, but rather is directed to the introduction of a recombinant DNA molecule. The transgenic organisms contemplated in accordance with the present invention include bacteria, cyanobacteria, fungi, plants and animals. The isolated DNA of the present invention can be introduced into the host by methods known in the art, for example infection, transfection, transformation or transconjugation. Techniques for transferring the DNA of the present invention into such organisms are widely known and provided in references such as Sambrook et al. (1989), supra.

A "variant" of a particular nucleic acid sequence is defined as a nucleic acid sequence having at least 40% sequence identity to the particular nucleic acid sequence over a certain length of one of the nucleic acid sequences using blastn with the "BLAST 2 Sequences" tool Version 2.0.9 (May-07-1999) set at default parameters. Such a pair of nucleic acids may show, for example, at least 50%, at least 60%, at least 70%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% or greater sequence identity over a certain defined length. A variant may be described as, for example, an "allelic" (as defined above), "splice," "species," or "polymorphic" variant. A splice variant may have significant identity to a reference molecule, but will generally have a greater or lesser number of polynucleotides due to alternate splicing of exons during mRNA processing. The corresponding polypeptide may possess additional functional domains or lack domains that are present in the reference molecule. Species variants are polynucleotide sequences that vary from one species to another. The resulting polypeptides will generally have significant amino acid identity relative to each other. A polymorphic variant is a variation in the polynucleotide sequence of a particular gene between individuals of a given species. Polymorphic variants also may encompass "single nucleotide polymorphisms" (SNPs) in which the polynucleotide sequence varies by one nucleotide base. The presence of SNPs may be indicative of, for example, a certain population, a disease state, or a propensity for a disease state.

A "variant" of a particular polypeptide sequence is defined as a polypeptide sequence having at least 40% sequence identity to the particular polypeptide sequence over a certain length of one of the polypeptide sequences using blastp with the "BLAST 2 Sequences" tool Version 2.0.9 (May-07-1999) set at default parameters. Such a pair of polypeptides may show, for example, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% or greater sequence identity over a certain defined length of one of the polypeptides.

THE INVENTION

The invention is based on the discovery of new human secreted proteins (SECP), the polynucleotides encoding SECP, and the use of these compositions for the diagnosis, treatment, or prevention of cell proliferative, autoimmune/inflammatory, cardiovascular, neurological, and developmental disorders.

Table 1 summarizes the nomenclature for the full length polynucleotide and polypeptide sequences of the invention. Each polynucleotide and its corresponding polypeptide are correlated to a single Incyte project identification number (Incyte Project ID). Each polypeptide sequence is denoted by both a polypeptide sequence identification number (Polypeptide SEQ ID NO:) and an Incyte

polypeptide sequence number (Incyte Polypeptide ID) as shown. Each polynucleotide sequence is denoted by both a polynucleotide sequence identification number (Polynucleotide SEQ ID NO:) and an Incyte polynucleotide consensus sequence number (Incyte Polynucleotide ID) as shown.

Table 2 shows sequences with homology to the polypeptides of SEQ ID NO:1-7, SEQ ID NO:34-35, and SEQ ID NO:57-58, as identified by BLAST analysis against the GenBank protein (genpept) database. Columns 1 and 2 show polypeptides of SEQ ID NO:1-7, SEQ ID NO:34-35, and SEQ ID NO:57-58 and their corresponding Incyte polypeptide sequence numbers (Incyte Polypeptide ID). Column 3 shows the GenBank identification number (Genbank ID NO:) of the nearest GenBank homolog. Column 4 shows the probability score for the match between SEQ ID NO:1-7, SEQ ID NO:34-35, and SEQ ID NO:57-58 and their GenBank homologs. Column 5 shows the annotation of the GenBank homolog along with relevant citations where applicable, all of which are expressly incorporated by reference herein.

Table 3 shows various structural features of each of the polypeptides of the invention. Columns 1 and 2 show the polypeptide sequence identification number (SEQ ID NO:) and the corresponding Incyte polypeptide sequence number (Incyte Polypeptide ID) for each polypeptide of the invention. Column 3 shows the number of amino acid residues in each polypeptide. Column 4 shows potential phosphorylation sites, and column 5 shows potential glycosylation sites, as determined by the MOTIFS program of the GCG sequence analysis software package (Genetics Computer Group, Madison WI). Column 6 shows amino acid residues comprising signature sequences, domains, and motifs, including the locations of signal peptides (as indicated by "Signal Peptide" and/or "signal_cleavage"). Column 7 shows analytical methods for protein structure/function analysis and in some cases, searchable databases to which the analytical methods were applied.

Together, Tables 2 and 3 summarize the properties of polypeptides of the invention, and these properties establish that the claimed polypeptides are secreted proteins. For example, SEQ ID NO:1 is 56% identical to a human cerebral cell adhesion molecule (GenBank ID g5764665) as determined by the Basic Local Alignment Search Tool (BLAST). (See Table 2.) The BLAST probability score is 3.8e-156, which indicates the probability of obtaining the observed polypeptide sequence alignment by chance. SEQ ID NO:1 also contains a lysyl hydrolase domain as determined by searching for statistically significant matches in the hidden Markov model (HMM)-based PFAM database of conserved protein family domains. (See Table 3.) Data from SPSCAN, HMMER, BLAST_PRODOM and BLAST_DOMO analyses using other sequence databases provide further corroborative evidence that SEQ ID NO:1 is a secreted hydrolase. In an alternative example, SEQ ID NO:2 is 32% identical to mouse seizure-related gene product 6 precursor (GenBank ID g693910) and is 67% identical from residue S22 to residue R527 to human CUB and sushi multiple domains 1

protein (GenBank ID g14794726) as determined by the Basic Local Alignment Search Tool (BLAST). (See Table 2.) The BLAST probability scores are 2.9×10^{-42} and 0.0 respectively, which indicate the probabilities of obtaining the observed polypeptide sequence alignments by chance. SEQ ID NO:2 also contains three sushi domains and two CUB domains as determined by searching for statistically significant matches in the hidden Markov model (HMM)-based PFAM database of conserved protein family domains. (See Table 3.) In addition, SEQ ID NO:2 contains a signal peptide as identified by HMMER analysis. Data from BLIMPS analysis provides further corroborative evidence that SEQ ID NO:2 is a secreted protein which contains sushi domains. In an alternative example, SEQ ID NO:3 shares 51% local identity to a mouse transmembrane protein (GenBank ID g7259265) as determined by the Basic Local Alignment Search Tool (BLAST). (See Table 2.) The BLAST probability score is 2.4×10^{-28} , which indicates the probability of obtaining the observed polypeptide sequence alignment by chance. SEQ ID NO:3 also contains a signal peptide as determined by searching for statistically significant matches in the hidden Markov model (HMM)-based PFAM database of conserved protein family domains. (See Table 3.) Data from SPSCAN analyses provide further corroborative evidence that SEQ ID NO:3 is a secreted protein. In an alternative example, SEQ ID NO:58 is 39% identical to ZOG, a rat zona glomerulosa specific protein (GenBank ID g3097285) as determined by the Basic Local Alignment Search Tool (BLAST). (See Table 2.) The BLAST probability score is 1.2×10^{-65} , which indicates the probability of obtaining the observed polypeptide sequence alignment by chance. SEQ ID NO:58 contains a signal peptide and single transmembrane domain. SEQ ID NO:58 also contains a number of EGF-like domains as determined by searching for statistically significant matches in the hidden Markov model (HMM)-based PFAM database of conserved protein family domains. (See Table 3.) The presence of this motif is confirmed by BLIMPS and MOTIFS analyses, providing further corroborative evidence that SEQ ID NO:58 is a secreted protein. SEQ ID NO:4-57 and SEQ ID NO:59-67 were analyzed and annotated in a similar manner. The algorithms and parameters for the analysis of SEQ ID NO:1-67 are described in Table 7.

As shown in Table 4, the full length polynucleotide sequences of the present invention were assembled using cDNA sequences or coding (exon) sequences derived from genomic DNA, or any combination of these two types of sequences. Columns 1 and 2 list the polynucleotide sequence identification number (Polynucleotide SEQ ID NO:) and the corresponding Incyte polynucleotide consensus sequence number (Incyte Polynucleotide ID) for each polynucleotide of the invention. Column 3 shows the length of each polynucleotide sequence in base pairs. Column 4 lists fragments of the polynucleotide sequences which are useful, for example, in hybridization or amplification technologies that identify SEQ ID NO:68-136 or that distinguish between SEQ ID NO:68-136 and related polynucleotide sequences. Column 5 shows identification numbers corresponding to cDNA

sequences, coding sequences (exons) predicted from genomic DNA, and/or sequence assemblages comprised of both cDNA and genomic DNA. These sequences were used to assemble the full length polynucleotide sequences of the invention. Columns 6 and 7 of Table 4 show the nucleotide start (5') and stop (3') positions of the cDNA and/or genomic sequences in column 5 relative to their respective full length sequences.

The identification numbers in Column 5 of Table 4 may refer specifically, for example, to Incyte cDNAs along with their corresponding cDNA libraries. For example, 8052177J1 is the identification number of an Incyte cDNA sequence, and FTUBTUE01 is the cDNA library from which it is derived. Incyte cDNAs for which cDNA libraries are not indicated were derived from pooled cDNA libraries (e.g., 71926854V1). Alternatively, the identification numbers in column 5 may refer to GenBank cDNAs or ESTs (e.g., g2204647) which contributed to the assembly of the full length polynucleotide sequences. In addition, the identification numbers in column 5 may identify sequences derived from the ENSEMBL (The Sanger Centre, Cambridge, UK) database (*i.e.*, those sequences including the designation "ENST"). Alternatively, the identification numbers in column 5 may be derived from the NCBI RefSeq Nucleotide Sequence Records Database (*i.e.*, those sequences including the designation "NM" or "NT") or the NCBI RefSeq Protein Sequence Records (*i.e.*, those sequences including the designation "NP"). Alternatively, the identification numbers in column 5 may refer to assemblages of both cDNA and Genscan-predicted exons brought together by an "exon stitching" algorithm. For example, FL_XXXXXX_N₁_N₂YYYYY_N₃_N₄ represents a "stitched" sequence in which XXXXXX is the identification number of the cluster of sequences to which the algorithm was applied, and YYYYYY is the number of the prediction generated by the algorithm, and N_{1,2,3,...}, if present, represent specific exons that may have been manually edited during analysis (See Example V). Alternatively, the identification numbers in column 5 may refer to assemblages of exons brought together by an "exon-stretching" algorithm. For example, FLXXXXXX_gAAAAA_gBBBBB_1_N is the identification number of a "stretched" sequence, with XXXXXX being the Incyte project identification number, gAAAAA being the GenBank identification number of the human genomic sequence to which the "exon-stretching" algorithm was applied, gBBBBB being the GenBank identification number or NCBI RefSeq identification number of the nearest GenBank protein homolog, and N referring to specific exons (See Example V). In instances where a RefSeq sequence was used as a protein homolog for the "exon-stretching" algorithm, a RefSeq identifier (denoted by "NM," "NP," or "NT") may be used in place of the GenBank identifier (*i.e.*, gBBBBB).

Alternatively, a prefix identifies component sequences that were hand-edited, predicted from genomic DNA sequences, or derived from a combination of sequence analysis methods. The following Table lists examples of component sequence prefixes and corresponding sequence analysis

methods associated with the prefixes (see Example IV and Example V).

Prefix	Type of analysis and/or examples of programs
GNN, GFG, ENST	Exon prediction from genomic sequences using, for example, GENSCAN (Stanford University, CA, USA) or FGENES (Computer Genomics Group, The Sanger Centre, Cambridge, UK).
GBI	Hand-edited analysis of genomic sequences.
FL	Stitched or stretched genomic sequences (see Example V).
INCY	Full length transcript and exon prediction from mapping of EST sequences to the genome. Genomic location and EST composition data are combined to predict the exons and resulting transcript.

In some cases, Incyte cDNA coverage redundant with the sequence coverage shown in column 5 was obtained to confirm the final consensus polynucleotide sequence, but the relevant Incyte cDNA identification numbers are not shown.

Table 5 shows the representative cDNA libraries for those full length polynucleotide sequences which were assembled using Incyte cDNA sequences. The representative cDNA library is the Incyte cDNA library which is most frequently represented by the Incyte cDNA sequences which were used to assemble and confirm the above polynucleotide sequences. The tissues and vectors which were used to construct the cDNA libraries shown in Table 5 are described in Table 6.

The invention also encompasses SECP variants. A preferred SECP variant is one which has at least about 80%, or alternatively at least about 90%, or even at least about 95% amino acid sequence identity to the SECP amino acid sequence, and which contains at least one functional or structural characteristic of SECP.

The invention also encompasses polynucleotides which encode SECP. In a particular embodiment, the invention encompasses a polynucleotide sequence comprising a sequence selected from the group consisting of SEQ ID NO:68-134, which encodes SECP. The polynucleotide sequences of SEQ ID NO:68-134, as presented in the Sequence Listing, embrace the equivalent RNA sequences, wherein occurrences of the nitrogenous base thymine are replaced with uracil, and the sugar backbone is composed of ribose instead of deoxyribose.

The invention also encompasses a variant of a polynucleotide sequence encoding SECP. In particular, such a variant polynucleotide sequence will have at least about 70%, or alternatively at least about 85%, or even at least about 95% polynucleotide sequence identity to the polynucleotide sequence encoding SECP. A particular aspect of the invention encompasses a variant of a polynucleotide sequence comprising a sequence selected from the group consisting of SEQ ID NO:68-134 which has at least about 70%, or alternatively at least about 85%, or even at least about

95% polynucleotide sequence identity to a nucleic acid sequence selected from the group consisting of SEQ ID NO:68-134. Any one of the polynucleotide variants described above can encode an amino acid sequence which contains at least one functional or structural characteristic of SECP.

It will be appreciated by those skilled in the art that as a result of the degeneracy of the genetic code, a multitude of polynucleotide sequences encoding SECP, some bearing minimal
5 similarity to the polynucleotide sequences of any known and naturally occurring gene, may be produced. Thus, the invention contemplates each and every possible variation of polynucleotide sequence that could be made by selecting combinations based on possible codon choices. These combinations are made in accordance with the standard triplet genetic code as applied to the
10 polynucleotide sequence of naturally occurring SECP, and all such variations are to be considered as being specifically disclosed.

Although nucleotide sequences which encode SECP and its variants are generally capable of hybridizing to the nucleotide sequence of the naturally occurring SECP under appropriately selected conditions of stringency, it may be advantageous to produce nucleotide sequences encoding SECP or
15 its derivatives possessing a substantially different codon usage, e.g., inclusion of non-naturally occurring codons. Codons may be selected to increase the rate at which expression of the peptide occurs in a particular prokaryotic or eukaryotic host in accordance with the frequency with which particular codons are utilized by the host. Other reasons for substantially altering the nucleotide sequence encoding SECP and its derivatives without altering the encoded amino acid sequences
20 include the production of RNA transcripts having more desirable properties, such as a greater half-life, than transcripts produced from the naturally occurring sequence.

The invention also encompasses production of DNA sequences which encode SECP and SECP derivatives, or fragments thereof, entirely by synthetic chemistry. After production, the synthetic sequence may be inserted into any of the many available expression vectors and cell
25 systems using reagents well known in the art. Moreover, synthetic chemistry may be used to introduce mutations into a sequence encoding SECP or any fragment thereof.

Also encompassed by the invention are polynucleotide sequences that are capable of hybridizing to the claimed polynucleotide sequences, and, in particular, to those shown in SEQ ID NO:68-134 and fragments thereof under various conditions of stringency. (See, e.g., Wahl, G.M. and
30 S.L. Berger (1987) *Methods Enzymol.* 152:399-407; Kimmel, A.R. (1987) *Methods Enzymol.* 152:507-511.) Hybridization conditions, including annealing and wash conditions, are described in "Definitions."

Methods for DNA sequencing are well known in the art and may be used to practice any of the embodiments of the invention. The methods may employ such enzymes as the Klenow fragment
35 of DNA polymerase I, SEQUENASE (US Biochemical, Cleveland OH), Taq polymerase (Applied

Biosystems), thermostable T7 polymerase (Amersham Pharmacia Biotech, Piscataway NJ), or combinations of polymerases and proofreading exonucleases such as those found in the ELONGASE amplification system (Life Technologies, Gaithersburg MD). Preferably, sequence preparation is automated with machines such as the MICROLAB 2200 liquid transfer system (Hamilton, Reno NV),
5 PTC200 thermal cycler (MJ Research, Watertown MA) and ABI CATALYST 800 thermal cycler (Applied Biosystems). Sequencing is then carried out using either the ABI 373 or 377 DNA sequencing system (Applied Biosystems), the MEGABACE 1000 DNA sequencing system (Molecular Dynamics, Sunnyvale CA), or other systems known in the art. The resulting sequences are analyzed using a variety of algorithms which are well known in the art. (See, e.g., Ausubel, F.M.
10 (1997) Short Protocols in Molecular Biology, John Wiley & Sons, New York NY, unit 7.7; Meyers, R.A. (1995) Molecular Biology and Biotechnology, Wiley VCH, New York NY, pp. 856-853.)

The nucleic acid sequences encoding SECP may be extended utilizing a partial nucleotide sequence and employing various PCR-based methods known in the art to detect upstream sequences, such as promoters and regulatory elements. For example, one method which may be employed,
15 restriction-site PCR, uses universal and nested primers to amplify unknown sequence from genomic DNA within a cloning vector. (See, e.g., Sarkar, G. (1993) PCR Methods Applic. 2:318-322.) Another method, inverse PCR, uses primers that extend in divergent directions to amplify unknown sequence from a circularized template. The template is derived from restriction fragments comprising a known genomic locus and surrounding sequences. (See, e.g., Triglia, T. et al. (1988) Nucleic Acids
20 Res. 16:8186.) A third method, capture PCR, involves PCR amplification of DNA fragments adjacent to known sequences in human and yeast artificial chromosome DNA. (See, e.g., Lagerstrom, M. et al. (1991) PCR Methods Applic. 1:111-119.) In this method, multiple restriction enzyme digestions and ligations may be used to insert an engineered double-stranded sequence into a region of unknown sequence before performing PCR. Other methods which may be used to retrieve
25 unknown sequences are known in the art. (See, e.g., Parker, J.D. et al. (1991) Nucleic Acids Res. 19:3055-3060). Additionally, one may use PCR, nested primers, and PROMOTERFINDER libraries (Clontech, Palo Alto CA) to walk genomic DNA. This procedure avoids the need to screen libraries and is useful in finding intron/exon junctions. For all PCR-based methods, primers may be designed using commercially available software, such as OLIGO 4.06 primer analysis software (National
30 Biosciences, Plymouth MN) or another appropriate program, to be about 22 to 30 nucleotides in length, to have a GC content of about 50% or more, and to anneal to the template at temperatures of about 68°C to 72°C.

When screening for full length cDNAs, it is preferable to use libraries that have been size-selected to include larger cDNAs. In addition, random-primed libraries, which often include
35 sequences containing the 5' regions of genes, are preferable for situations in which an oligo d(T)

library does not yield a full-length cDNA. Genomic libraries may be useful for extension of sequence into 5' non-transcribed regulatory regions.

Capillary electrophoresis systems which are commercially available may be used to analyze the size or confirm the nucleotide sequence of sequencing or PCR products. In particular, capillary sequencing may employ flowable polymers for electrophoretic separation, four different nucleotide-specific, laser-stimulated fluorescent dyes, and a charge coupled device camera for detection of the emitted wavelengths. Output/light intensity may be converted to electrical signal using appropriate software (e.g., GENOTYPER and SEQUENCE NAVIGATOR, Applied Biosystems), and the entire process from loading of samples to computer analysis and electronic data display may be computer controlled. Capillary electrophoresis is especially preferable for sequencing small DNA fragments which may be present in limited amounts in a particular sample.

In another embodiment of the invention, polynucleotide sequences or fragments thereof which encode SECP may be cloned in recombinant DNA molecules that direct expression of SECP, or fragments or functional equivalents thereof, in appropriate host cells. Due to the inherent degeneracy of the genetic code, other DNA sequences which encode substantially the same or a functionally equivalent amino acid sequence may be produced and used to express SECP.

The nucleotide sequences of the present invention can be engineered using methods generally known in the art in order to alter SECP-encoding sequences for a variety of purposes including, but not limited to, modification of the cloning, processing, and/or expression of the gene product. DNA shuffling by random fragmentation and PCR reassembly of gene fragments and synthetic oligonucleotides may be used to engineer the nucleotide sequences. For example, oligonucleotide-mediated site-directed mutagenesis may be used to introduce mutations that create new restriction sites, alter glycosylation patterns, change codon preference, produce splice variants, and so forth.

The nucleotides of the present invention may be subjected to DNA shuffling techniques such as MOLECULARBREEDING (Maxygen Inc., Santa Clara CA; described in U.S. Patent No. 5,837,458; Chang, C.-C. et al. (1999) Nat. Biotechnol. 17:793-797; Christians, F.C. et al. (1999) Nat. Biotechnol. 17:259-264; and Crameri, A. et al. (1996) Nat. Biotechnol. 14:315-319) to alter or improve the biological properties of SECP, such as its biological or enzymatic activity or its ability to bind to other molecules or compounds. DNA shuffling is a process by which a library of gene variants is produced using PCR-mediated recombination of gene fragments. The library is then subjected to selection or screening procedures that identify those gene variants with the desired properties. These preferred variants may then be pooled and further subjected to recursive rounds of DNA shuffling and selection/screening. Thus, genetic diversity is created through "artificial" breeding and rapid molecular evolution. For example, fragments of a single gene containing random point mutations may be recombined, screened, and then reshuffled until the desired properties are

optimized. Alternatively, fragments of a given gene may be recombined with fragments of homologous genes in the same gene family, either from the same or different species, thereby maximizing the genetic diversity of multiple naturally occurring genes in a directed and controllable manner.

- 5 In another embodiment, sequences encoding SECP may be synthesized, in whole or in part, using chemical methods well known in the art. (See, e.g., Caruthers, M.H. et al. (1980) *Nucleic Acids Symp. Ser.* 7:215-223; and Horn, T. et al. (1980) *Nucleic Acids Symp. Ser.* 7:225-232.) Alternatively, SECP itself or a fragment thereof may be synthesized using chemical methods. For example, peptide synthesis can be performed using various solution-phase or solid-phase techniques.
- 10 (See, e.g., Creighton, T. (1984) Proteins, Structures and Molecular Properties, WH Freeman, New York NY, pp. 55-60; and Roberge, J.Y. et al. (1995) *Science* 269:202-204.) Automated synthesis may be achieved using the ABI 431A peptide synthesizer (Applied Biosystems). Additionally, the amino acid sequence of SECP, or any part thereof, may be altered during direct synthesis and/or combined with sequences from other proteins, or any part thereof, to produce a variant polypeptide or
- 15 a polypeptide having a sequence of a naturally occurring polypeptide.

The peptide may be substantially purified by preparative high performance liquid chromatography. (See, e.g., Chiez, R.M. and F.Z. Regnier (1990) *Methods Enzymol.* 182:392-421.) The composition of the synthetic peptides may be confirmed by amino acid analysis or by sequencing. (See, e.g., Creighton, *supra*, pp. 28-53.)

- 20 In order to express a biologically active SECP, the nucleotide sequences encoding SECP or derivatives thereof may be inserted into an appropriate expression vector, i.e., a vector which contains the necessary elements for transcriptional and translational control of the inserted coding sequence in a suitable host. These elements include regulatory sequences, such as enhancers, constitutive and inducible promoters, and 5' and 3' untranslated regions in the vector and in polynucleotide sequences
- 25 encoding SECP. Such elements may vary in their strength and specificity. Specific initiation signals may also be used to achieve more efficient translation of sequences encoding SECP. Such signals include the ATG initiation codon and adjacent sequences, e.g. the Kozak sequence. In cases where sequences encoding SECP and its initiation codon and upstream regulatory sequences are inserted into the appropriate expression vector, no additional transcriptional or translational control signals
- 30 may be needed. However, in cases where only coding sequence, or a fragment thereof, is inserted, exogenous translational control signals including an in-frame ATG initiation codon should be provided by the vector. Exogenous translational elements and initiation codons may be of various origins, both natural and synthetic. The efficiency of expression may be enhanced by the inclusion of enhancers appropriate for the particular host cell system used. (See, e.g., Scharf, D. et al. (1994)
- 35 *Results Probl. Cell Differ.* 20:125-162.)

Methods which are well known to those skilled in the art may be used to construct expression vectors containing sequences encoding SECP and appropriate transcriptional and translational control elements. These methods include in vitro recombinant DNA techniques, synthetic techniques, and in vivo genetic recombination. (See, e.g., Sambrook, J. et al. (1989) Molecular Cloning, A Laboratory
 5 Manual, Cold Spring Harbor Press, Plainview NY, ch. 4, 8, and 16-17; Ausubel, F.M. et al. (1995) Current Protocols in Molecular Biology, John Wiley & Sons, New York NY, ch. 9, 13, and 16.)

A variety of expression vector/host systems may be utilized to contain and express sequences encoding SECP. These include, but are not limited to, microorganisms such as bacteria transformed with recombinant bacteriophage, plasmid, or cosmid DNA expression vectors; yeast transformed with
 10 yeast expression vectors; insect cell systems infected with viral expression vectors (e.g., baculovirus); plant cell systems transformed with viral expression vectors (e.g., cauliflower mosaic virus, CaMV, or tobacco mosaic virus, TMV) or with bacterial expression vectors (e.g., Ti or pBR322 plasmids); or animal cell systems. (See, e.g., Sambrook, supra; Ausubel, supra; Van Heeke, G. and S.M. Schuster (1989) J. Biol. Chem. 264:5503-5509; Engelhard, E.K. et al. (1994) Proc. Natl. Acad. Sci. USA
 15 91:3224-3227; Sandig, V. et al. (1996) Hum. Gene Ther. 7:1937-1945; Takamatsu, N. (1987) EMBO J. 6:307-311; The McGraw Hill Yearbook of Science and Technology (1992) McGraw Hill, New York NY, pp. 191-196; Logan, J. and T. Shenk (1984) Proc. Natl. Acad. Sci. USA 81:3655-3659; and Harrington, J.J. et al. (1997) Nat. Genet. 15:345-355.) Expression vectors derived from retroviruses, adenoviruses, or herpes or vaccinia viruses, or from various bacterial plasmids, may be used for
 20 delivery of nucleotide sequences to the targeted organ, tissue, or cell population. (See, e.g., Di Nicola, M. et al. (1998) Cancer Gen. Ther. 5(6):350-356; Yu, M. et al. (1993) Proc. Natl. Acad. Sci. USA 90(13):6340-6344; Buller, R.M. et al. (1985) Nature 317(6040):813-815; McGregor, D.P. et al. (1994) Mol. Immunol. 31(3):219-226; and Verma, I.M. and N. Somia (1997) Nature 389:239-242.)
 The invention is not limited by the host cell employed.

25 In bacterial systems, a number of cloning and expression vectors may be selected depending upon the use intended for polynucleotide sequences encoding SECP. For example, routine cloning, subcloning, and propagation of polynucleotide sequences encoding SECP can be achieved using a multifunctional E. coli vector such as PBLUESCRIPT (Stratagene, La Jolla CA) or PSPORT1 plasmid (Life Technologies). Ligation of sequences encoding SECP into the vector's multiple
 30 cloning site disrupts the *lacZ* gene, allowing a colorimetric screening procedure for identification of transformed bacteria containing recombinant molecules. In addition, these vectors may be useful for in vitro transcription, dideoxy sequencing, single strand rescue with helper phage, and creation of nested deletions in the cloned sequence. (See, e.g., Van Heeke, G. and S.M. Schuster (1989) J. Biol. Chem. 264:5503-5509.) When large quantities of SECP are needed, e.g. for the production of
 35 antibodies, vectors which direct high level expression of SECP may be used. For example, vectors

containing the strong, inducible SP6 or T7 bacteriophage promoter may be used.

Yeast expression systems may be used for production of SECP. A number of vectors containing constitutive or inducible promoters, such as alpha factor, alcohol oxidase, and PGH promoters, may be used in the yeast Saccharomyces cerevisiae or Pichia pastoris. In addition, such
5 vectors direct either the secretion or intracellular retention of expressed proteins and enable integration of foreign sequences into the host genome for stable propagation. (See, e.g., Ausubel, 1995, supra; Bitter, G.A. et al. (1987) *Methods Enzymol.* 153:516-544; and Scorer, C.A. et al. (1994) *Bio/Technology* 12:181-184.)

Plant systems may also be used for expression of SECP. Transcription of sequences
10 encoding SECP may be driven by viral promoters, e.g., the 35S and 19S promoters of CaMV used alone or in combination with the omega leader sequence from TMV (Takamatsu, N. (1987) *EMBO J.* 6:307-311). Alternatively, plant promoters such as the small subunit of RUBISCO or heat shock promoters may be used. (See, e.g., Coruzzi, G. et al. (1984) *EMBO J.* 3:1671-1680; Broglie, R. et al. (1984) *Science* 224:838-843; and Winter, J. et al. (1991) *Results Probl. Cell Differ.* 17:85-105.)
15 These constructs can be introduced into plant cells by direct DNA transformation or pathogen-mediated transfection. (See, e.g., The McGraw Hill Yearbook of Science and Technology (1992) McGraw Hill, New York NY, pp. 191-196.)

In mammalian cells, a number of viral-based expression systems may be utilized. In cases where an adenovirus is used as an expression vector, sequences encoding SECP may be ligated into
20 an adenovirus transcription/translation complex consisting of the late promoter and tripartite leader sequence. Insertion in a non-essential E1 or E3 region of the viral genome may be used to obtain infective virus which expresses SECP in host cells. (See, e.g., Logan, J. and T. Shenk (1984) *Proc. Natl. Acad. Sci. USA* 81:3655-3659.) In addition, transcription enhancers, such as the Rous sarcoma virus (RSV) enhancer, may be used to increase expression in mammalian host cells. SV40 or EBV-
25 based vectors may also be used for high-level protein expression.

Human artificial chromosomes (HACs) may also be employed to deliver larger fragments of DNA than can be contained in and expressed from a plasmid. HACs of about 6 kb to 10 Mb are constructed and delivered via conventional delivery methods (liposomes, polycationic amino polymers, or vesicles) for therapeutic purposes. (See, e.g., Harrington, J.J. et al. (1997) *Nat. Genet.*
30 15:345-355.)

For long term production of recombinant proteins in mammalian systems, stable expression of SECP in cell lines is preferred. For example, sequences encoding SECP can be transformed into cell lines using expression vectors which may contain viral origins of replication and/or endogenous expression elements and a selectable marker gene on the same or on a separate vector. Following the
35 introduction of the vector, cells may be allowed to grow for about 1 to 2 days in enriched media

before being switched to selective media. The purpose of the selectable marker is to confer resistance to a selective agent, and its presence allows growth and recovery of cells which successfully express the introduced sequences. Resistant clones of stably transformed cells may be propagated using tissue culture techniques appropriate to the cell type.

- 5 Any number of selection systems may be used to recover transformed cell lines. These include, but are not limited to, the herpes simplex virus thymidine kinase and adenine phosphoribosyltransferase genes, for use in *tk* and *apr* cells, respectively. (See, e.g., Wigler, M. et al. (1977) Cell 11:223-232; Lowy, I. et al. (1980) Cell 22:817-823.) Also, antimetabolite, antibiotic, or herbicide resistance can be used as the basis for selection. For example, *dhfr* confers resistance to
- 10 methotrexate; *neo* confers resistance to the aminoglycosides neomycin and G-418; and *als* and *pat* confer resistance to chlorsulfuron and phosphinotricin acetyltransferase, respectively. (See, e.g., Wigler, M. et al. (1980) Proc. Natl. Acad. Sci. USA 77:3567-3570; Colbere-Garapin, F. et al. (1981) J. Mol. Biol. 150:1-14.) Additional selectable genes have been described, e.g., *trpB* and *hisD*, which alter cellular requirements for metabolites. (See, e.g., Hartman, S.C. and R.C. Mulligan (1988) Proc.
- 15 Natl. Acad. Sci. USA 85:8047-8051.) Visible markers, e.g., anthocyanins, green fluorescent proteins (GFP; Clontech), β glucuronidase and its substrate β -glucuronide, or luciferase and its substrate luciferin may be used. These markers can be used not only to identify transformants, but also to quantify the amount of transient or stable protein expression attributable to a specific vector system. (See, e.g., Rhodes, C.A. (1995) Methods Mol. Biol. 55:121-131.)
- 20 Although the presence/absence of marker gene expression suggests that the gene of interest is also present, the presence and expression of the gene may need to be confirmed. For example, if the sequence encoding SECP is inserted within a marker gene sequence, transformed cells containing sequences encoding SECP can be identified by the absence of marker gene function. Alternatively, a marker gene can be placed in tandem with a sequence encoding SECP under the control of a single
- 25 promoter. Expression of the marker gene in response to induction or selection usually indicates expression of the tandem gene as well.

In general, host cells that contain the nucleic acid sequence encoding SECP and that express SECP may be identified by a variety of procedures known to those of skill in the art. These procedures include, but are not limited to, DNA-DNA or DNA-RNA hybridizations, PCR

30 amplification, and protein bioassay or immunoassay techniques which include membrane, solution, or chip based technologies for the detection and/or quantification of nucleic acid or protein sequences.

Immunological methods for detecting and measuring the expression of SECP using either specific polyclonal or monoclonal antibodies are known in the art. Examples of such techniques include enzyme-linked immunosorbent assays (ELISAs), radioimmunoassays (RIAs), and

35 fluorescence activated cell sorting (FACS). A two-site, monoclonal-based immunoassay utilizing

monoclonal antibodies reactive to two non-interfering epitopes on SECP is preferred, but a competitive binding assay may be employed. These and other assays are well known in the art. (See, e.g., Hampton, R. et al. (1990) Serological Methods, a Laboratory Manual, APS Press, St. Paul MN, Sect. IV; Coligan, J.E. et al. (1997) Current Protocols in Immunology, Greene Pub. Associates and Wiley-Interscience, New York NY; and Pound, J.D. (1998) Immunochemical Protocols, Humana Press, Totowa NJ.)

A wide variety of labels and conjugation techniques are known by those skilled in the art and may be used in various nucleic acid and amino acid assays. Means for producing labeled hybridization or PCR probes for detecting sequences related to polynucleotides encoding SECP include oligolabeling, nick translation, end-labeling, or PCR amplification using a labeled nucleotide. Alternatively, the sequences encoding SECP, or any fragments thereof, may be cloned into a vector for the production of an mRNA probe. Such vectors are known in the art, are commercially available, and may be used to synthesize RNA probes *in vitro* by addition of an appropriate RNA polymerase such as T7, T3, or SP6 and labeled nucleotides. These procedures may be conducted using a variety of commercially available kits, such as those provided by Amersham Pharmacia Biotech, Promega (Madison WI), and US Biochemical. Suitable reporter molecules or labels which may be used for ease of detection include radionuclides, enzymes, fluorescent, chemiluminescent, or chromogenic agents, as well as substrates, cofactors, inhibitors, magnetic particles, and the like.

Host cells transformed with nucleotide sequences encoding SECP may be cultured under conditions suitable for the expression and recovery of the protein from cell culture. The protein produced by a transformed cell may be secreted or retained intracellularly depending on the sequence and/or the vector used. As will be understood by those of skill in the art, expression vectors containing polynucleotides which encode SECP may be designed to contain signal sequences which direct secretion of SECP through a prokaryotic or eukaryotic cell membrane.

In addition, a host cell strain may be chosen for its ability to modulate expression of the inserted sequences or to process the expressed protein in the desired fashion. Such modifications of the polypeptide include, but are not limited to, acetylation, carboxylation, glycosylation, phosphorylation, lipidation, and acylation. Post-translational processing which cleaves a "prepro" or "pro" form of the protein may also be used to specify protein targeting, folding, and/or activity. Different host cells which have specific cellular machinery and characteristic mechanisms for post-translational activities (e.g., CHO, HeLa, MDCK, HEK293, and WI38) are available from the American Type Culture Collection (ATCC, Manassas VA) and may be chosen to ensure the correct modification and processing of the foreign protein.

In another embodiment of the invention, natural, modified, or recombinant nucleic acid sequences encoding SECP may be ligated to a heterologous sequence resulting in translation of a

fusion protein in any of the aforementioned host systems. For example, a chimeric SECP protein containing a heterologous moiety that can be recognized by a commercially available antibody may facilitate the screening of peptide libraries for inhibitors of SECP activity. Heterologous protein and peptide moieties may also facilitate purification of fusion proteins using commercially available
5 affinity matrices. Such moieties include, but are not limited to, glutathione S-transferase (GST), maltose binding protein (MBP), thioredoxin (Trx), calmodulin binding peptide (CBP), 6-His, FLAG, *c-myc*, and hemagglutinin (HA). GST, MBP, Trx, CBP, and 6-His enable purification of their cognate fusion proteins on immobilized glutathione, maltose, phenylarsine oxide, calmodulin, and metal-chelate resins, respectively. FLAG, *c-myc*, and hemagglutinin (HA) enable immunoaffinity
10 purification of fusion proteins using commercially available monoclonal and polyclonal antibodies that specifically recognize these epitope tags. A fusion protein may also be engineered to contain a proteolytic cleavage site located between the SECP encoding sequence and the heterologous protein sequence, so that SECP may be cleaved away from the heterologous moiety following purification. Methods for fusion protein expression and purification are discussed in Ausubel (1995, supra, ch. 10).
15 A variety of commercially available kits may also be used to facilitate expression and purification of fusion proteins.

In a further embodiment of the invention, synthesis of radiolabeled SECP may be achieved in vitro using the TNT rabbit reticulocyte lysate or wheat germ extract system (Promega). These systems couple transcription and translation of protein-coding sequences operably associated with the
20 T7, T3, or SP6 promoters. Translation takes place in the presence of a radiolabeled amino acid precursor, for example, ³⁵S-methionine.

SECP of the present invention or fragments thereof may be used to screen for compounds that specifically bind to SECP. At least one and up to a plurality of test compounds may be screened for specific binding to SECP. Examples of test compounds include antibodies, oligonucleotides,
25 proteins (e.g., receptors), or small molecules.

In one embodiment, the compound thus identified is closely related to the natural ligand of SECP, e.g., a ligand or fragment thereof, a natural substrate, a structural or functional mimetic, or a natural binding partner. (See, e.g., Coligan, J.E. et al. (1991) Current Protocols in Immunology 1(2): Chapter 5.) Similarly, the compound can be closely related to the natural receptor to which SECP
30 binds, or to at least a fragment of the receptor, e.g., the ligand binding site. In either case, the compound can be rationally designed using known techniques. In one embodiment, screening for these compounds involves producing appropriate cells which express SECP, either as a secreted protein or on the cell membrane. Preferred cells include cells from mammals, yeast, Drosophila, or E. coli. Cells expressing SECP or cell membrane fractions which contain SECP are then contacted
35 with a test compound and binding, stimulation, or inhibition of activity of either SECP or the

compound is analyzed.

An assay may simply test binding of a test compound to the polypeptide, wherein binding is detected by a fluorophore, radioisotope, enzyme conjugate, or other detectable label. For example, the assay may comprise the steps of combining at least one test compound with SECP, either in
5 solution or affixed to a solid support, and detecting the binding of SECP to the compound.

Alternatively, the assay may detect or measure binding of a test compound in the presence of a labeled competitor. Additionally, the assay may be carried out using cell-free preparations, chemical libraries, or natural product mixtures, and the test compound(s) may be free in solution or affixed to a solid support.

10 SECP of the present invention or fragments thereof may be used to screen for compounds that modulate the activity of SECP. Such compounds may include agonists, antagonists, or partial or inverse agonists. In one embodiment, an assay is performed under conditions permissive for SECP activity, wherein SECP is combined with at least one test compound, and the activity of SECP in the presence of a test compound is compared with the activity of SECP in the absence of the test
15 compound. A change in the activity of SECP in the presence of the test compound is indicative of a compound that modulates the activity of SECP. Alternatively, a test compound is combined with an in vitro or cell-free system comprising SECP under conditions suitable for SECP activity, and the assay is performed. In either of these assays, a test compound which modulates the activity of SECP may do so indirectly and need not come in direct contact with the test compound. At least one and up
20 to a plurality of test compounds may be screened.

In another embodiment, polynucleotides encoding SECP or their mammalian homologs may be "knocked out" in an animal model system using homologous recombination in embryonic stem (ES) cells. Such techniques are well known in the art and are useful for the generation of animal models of human disease. (See, e.g., U.S. Patent No. 5,175,383 and U.S. Patent No. 5,767,337.) For
25 example, mouse ES cells, such as the mouse 129/SvJ cell line, are derived from the early mouse embryo and grown in culture. The ES cells are transformed with a vector containing the gene of interest disrupted by a marker gene, e.g., the neomycin phosphotransferase gene (neo; Capecchi, M.R. (1989) Science 244:1288-1292). The vector integrates into the corresponding region of the host genome by homologous recombination. Alternatively, homologous recombination takes place using
30 the Cre-loxP system to knockout a gene of interest in a tissue- or developmental stage-specific manner (Marth, J.D. (1996) Clin. Invest. 97:1999-2002; Wagner, K.U. et al. (1997) Nucleic Acids Res. 25:4323-4330). Transformed ES cells are identified and microinjected into mouse cell blastocysts such as those from the C57BL/6 mouse strain. The blastocysts are surgically transferred to pseudopregnant dams, and the resulting chimeric progeny are genotyped and bred to produce
35 heterozygous or homozygous strains. Transgenic animals thus generated may be tested with potential

therapeutic or toxic agents.

Polynucleotides encoding SECP may also be manipulated in vitro in ES cells derived from human blastocysts. Human ES cells have the potential to differentiate into at least eight separate cell lineages including endoderm, mesoderm, and ectodermal cell types. These cell lineages differentiate into, for example, neural cells, hematopoietic lineages, and cardiomyocytes (Thomson, J.A. et al. (1998) Science 282:1145-1147).

Polynucleotides encoding SECP can also be used to create "knockin" humanized animals (pigs) or transgenic animals (mice or rats) to model human disease. With knockin technology, a region of a polynucleotide encoding SECP is injected into animal ES cells, and the injected sequence integrates into the animal cell genome. Transformed cells are injected into blastulae, and the blastulae are implanted as described above. Transgenic progeny or inbred lines are studied and treated with potential pharmaceutical agents to obtain information on treatment of a human disease. Alternatively, a mammal inbred to overexpress SECP, e.g., by secreting SECP in its milk, may also serve as a convenient source of that protein (Janne, J. et al. (1998) Biotechnol. Annu. Rev. 4:55-74).

15 THERAPEUTICS

Chemical and structural similarity, e.g., in the context of sequences and motifs, exists between regions of SECP and secreted proteins. In addition, the expression of SECP is closely associated with neurological, gastrointestinal, cardiovascular, reproductive, developmental, diseased, and tumorous tissues such as adrenal gland tumor tissue. Therefore, SECP appears to play a role in cell proliferative, autoimmune/inflammatory, cardiovascular, neurological, and developmental disorders. In the treatment of disorders associated with increased SECP expression or activity, it is desirable to decrease the expression or activity of SECP. In the treatment of disorders associated with decreased SECP expression or activity, it is desirable to increase the expression or activity of SECP.

Therefore, in one embodiment, SECP or a fragment or derivative thereof may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of SECP. Examples of such disorders include, but are not limited to, a cell proliferative disorder such as actinic keratosis, arteriosclerosis, atherosclerosis, bursitis, cirrhosis, hepatitis, mixed connective tissue disease (MCTD), myelofibrosis, paroxysmal nocturnal hemoglobinuria, polycythemia vera, psoriasis, primary thrombocythemia, and cancers including adenocarcinoma, leukemia, lymphoma, melanoma, myeloma, sarcoma, teratocarcinoma, and, in particular, a cancer of the adrenal gland, bladder, bone, bone marrow, brain, breast, cervix, gall bladder, ganglia, gastrointestinal tract, heart, kidney, liver, lung, muscle, ovary, pancreas, parathyroid, penis, prostate, salivary glands, skin, spleen, testis, thymus, thyroid, and uterus; an autoimmune/inflammatory disorder such as acquired immunodeficiency syndrome (AIDS), Addison's disease, adult respiratory

distress syndrome, allergies, ankylosing spondylitis, amyloidosis, anemia, asthma, atherosclerosis, autoimmune hemolytic anemia, autoimmune thyroiditis, autoimmune polyendocrinopathy-candidiasis-ectodermal dystrophy (APECED), bronchitis, cholecystitis, contact dermatitis, Crohn's disease, atopic dermatitis, dermatomyositis, diabetes mellitus, emphysema, episodic lymphopenia

5 with lymphocytotoxins, erythroblastosis fetalis, erythema nodosum, atrophic gastritis, glomerulonephritis, Goodpasture's syndrome, gout, Graves' disease, Hashimoto's thyroiditis, hypereosinophilia, irritable bowel syndrome, multiple sclerosis, myasthenia gravis, myocardial or pericardial inflammation, osteoarthritis, osteoporosis, pancreatitis, polymyositis, psoriasis, Reiter's syndrome, rheumatoid arthritis, scleroderma, Sjögren's syndrome, systemic anaphylaxis, systemic

10 lupus erythematosus, systemic sclerosis, thrombocytopenic purpura, ulcerative colitis, uveitis, Werner syndrome, complications of cancer, hemodialysis, and extracorporeal circulation, viral, bacterial, fungal, parasitic, protozoal, and helminthic infections, and trauma; a cardiovascular disorder such as congestive heart failure, ischemic heart disease, angina pectoris, myocardial infarction, hypertensive heart disease, degenerative valvular heart disease, calcific aortic valve stenosis, congenitally bicuspid

15 aortic valve, mitral annular calcification, mitral valve prolapse, rheumatic fever and rheumatic heart disease, infective endocarditis, nonbacterial thrombotic endocarditis, endocarditis of systemic lupus erythematosus, carcinoid heart disease, cardiomyopathy, myocarditis, pericarditis, neoplastic heart disease, congenital heart disease, complications of cardiac transplantation, arteriovenous fistula, atherosclerosis, hypertension, vasculitis, Raynaud's disease, aneurysms, arterial dissections, varicose

20 veins, thrombophlebitis and phlebothrombosis, vascular tumors, and complications of thrombolysis, balloon angioplasty, vascular replacement, and coronary artery bypass graft surgery; a neurological disorder such as epilepsy, ischemic cerebrovascular disease, stroke, cerebral neoplasms, Alzheimer's disease, Pick's disease, Huntington's disease, dementia, Parkinson's disease and other extrapyramidal disorders, amyotrophic lateral sclerosis and other motor neuron disorders, progressive neural

25 muscular atrophy, retinitis pigmentosa, hereditary ataxias, multiple sclerosis and other demyelinating diseases, bacterial and viral meningitis, brain abscess, subdural empyema, epidural abscess, suppurative intracranial thrombophlebitis, myelitis and radiculitis, viral central nervous system disease, prion diseases including kuru, Creutzfeldt-Jakob disease, and Gerstmann-

30 Straussler-Scheinker syndrome, fatal familial insomnia, nutritional and metabolic diseases of the nervous system, neurofibromatosis, tuberous sclerosis, cerebelloretinal hemangioblastomatosis, encephalotrigeminal syndrome, mental retardation and other developmental disorders of the central nervous system including Down syndrome, cerebral palsy, neuroskeletal disorders, autonomic nervous system disorders, cranial nerve disorders, spinal cord diseases, muscular dystrophy and other neuromuscular disorders, peripheral nervous system disorders, dermatomyositis and polymyositis,

35 inherited, metabolic, endocrine, and toxic myopathies, myasthenia gravis, periodic paralysis, mental

disorders including mood, anxiety, and schizophrenic disorders, seasonal affective disorder (SAD), akathisia, amnesia, catatonia, diabetic neuropathy, tardive dyskinesia, dystonias, paranoid psychoses, postherpetic neuralgia, Tourette's disorder, progressive supranuclear palsy, corticobasal degeneration, and familial frontotemporal dementia; and a developmental disorder such as renal tubular acidosis, anemia, Cushing's syndrome, achondroplastic dwarfism, Duchenne and Becker muscular dystrophy, epilepsy, gonadal dysgenesis, WAGR syndrome (Wilms' tumor, aniridia, genitourinary abnormalities, and mental retardation), Smith-Magenis syndrome, myelodysplastic syndrome, hereditary mucoepithelial dysplasia, hereditary keratodermas, hereditary neuropathies such as Charcot-Marie-Tooth disease and neurofibromatosis, hypothyroidism, hydrocephalus, seizure disorders such as Sydenham's chorea and cerebral palsy, spina bifida, anencephaly, craniorachischisis, congenital glaucoma, cataract, and sensorineural hearing loss.

In another embodiment, a vector capable of expressing SECP or a fragment or derivative thereof may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of SECP including, but not limited to, those described above.

In a further embodiment, a composition comprising a substantially purified SECP in conjunction with a suitable pharmaceutical carrier may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of SECP including, but not limited to, those provided above.

In still another embodiment, an agonist which modulates the activity of SECP may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of SECP including, but not limited to, those listed above.

In a further embodiment, an antagonist of SECP may be administered to a subject to treat or prevent a disorder associated with increased expression or activity of SECP. Examples of such disorders include, but are not limited to, those cell proliferative, autoimmune/inflammatory, cardiovascular, neurological, and developmental disorders described above. In one aspect, an antibody which specifically binds SECP may be used directly as an antagonist or indirectly as a targeting or delivery mechanism for bringing a pharmaceutical agent to cells or tissues which express SECP.

In an additional embodiment, a vector expressing the complement of the polynucleotide encoding SECP may be administered to a subject to treat or prevent a disorder associated with increased expression or activity of SECP including, but not limited to, those described above.

In other embodiments, any of the proteins, antagonists, antibodies, agonists, complementary sequences, or vectors of the invention may be administered in combination with other appropriate therapeutic agents. Selection of the appropriate agents for use in combination therapy may be made by one of ordinary skill in the art, according to conventional pharmaceutical principles. The

combination of therapeutic agents may act synergistically to effect the treatment or prevention of the various disorders described above. Using this approach, one may be able to achieve therapeutic efficacy with lower dosages of each agent, thus reducing the potential for adverse side effects.

An antagonist of SECP may be produced using methods which are generally known in the art. In particular, purified SECP may be used to produce antibodies or to screen libraries of pharmaceutical agents to identify those which specifically bind SECP. Antibodies to SECP may also be generated using methods that are well known in the art. Such antibodies may include, but are not limited to, polyclonal, monoclonal, chimeric, and single chain antibodies, Fab fragments, and fragments produced by a Fab expression library. Neutralizing antibodies (i.e., those which inhibit dimer formation) are generally preferred for therapeutic use.

For the production of antibodies, various hosts including goats, rabbits, rats, mice, humans, and others may be immunized by injection with SECP or with any fragment or oligopeptide thereof which has immunogenic properties. Depending on the host species, various adjuvants may be used to increase immunological response. Such adjuvants include, but are not limited to, Freund's, mineral gels such as aluminum hydroxide, and surface active substances such as lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, KLH, and dinitrophenol. Among adjuvants used in humans, BCG (bacilli Calmette-Guerin) and Corynebacterium parvum are especially preferable.

It is preferred that the oligopeptides, peptides, or fragments used to induce antibodies to SECP have an amino acid sequence consisting of at least about 5 amino acids, and generally will consist of at least about 10 amino acids. It is also preferable that these oligopeptides, peptides, or fragments are identical to a portion of the amino acid sequence of the natural protein. Short stretches of SECP amino acids may be fused with those of another protein, such as KLH, and antibodies to the chimeric molecule may be produced.

Monoclonal antibodies to SECP may be prepared using any technique which provides for the production of antibody molecules by continuous cell lines in culture. These include, but are not limited to, the hybridoma technique, the human B-cell hybridoma technique, and the EBV-hybridoma technique. (See, e.g., Kohler, G. et al. (1975) Nature 256:495-497; Kozbor, D. et al. (1985) J. Immunol. Methods 81:31-42; Cote, R.J. et al. (1983) Proc. Natl. Acad. Sci. USA 80:2026-2030; and Cole, S.P. et al. (1984) Mol. Cell Biol. 62:109-120.)

In addition, techniques developed for the production of "chimeric antibodies," such as the splicing of mouse antibody genes to human antibody genes to obtain a molecule with appropriate antigen specificity and biological activity, can be used. (See, e.g., Morrison, S.L. et al. (1984) Proc. Natl. Acad. Sci. USA 81:6851-6855; Neuberger, M.S. et al. (1984) Nature 312:604-608; and Takeda, S. et al. (1985) Nature 314:452-454.) Alternatively, techniques described for the production of single chain antibodies may be adapted, using methods known in the art, to produce SECP-specific single

chain antibodies. Antibodies with related specificity, but of distinct idiotypic composition, may be generated by chain shuffling from random combinatorial immunoglobulin libraries. (See, e.g., Burton, D.R. (1991) Proc. Natl. Acad. Sci. USA 88:10134-10137.)

Antibodies may also be produced by inducing in vivo production in the lymphocyte
5 population or by screening immunoglobulin libraries or panels of highly specific binding reagents as disclosed in the literature. (See, e.g., Orlandi, R. et al. (1989) Proc. Natl. Acad. Sci. USA 86:3833-3837; Winter, G. et al. (1991) Nature 349:293-299.)

Antibody fragments which contain specific binding sites for SECP may also be generated. For example, such fragments include, but are not limited to, F(ab')₂ fragments produced by pepsin
10 digestion of the antibody molecule and Fab fragments generated by reducing the disulfide bridges of the F(ab')₂ fragments. Alternatively, Fab expression libraries may be constructed to allow rapid and easy identification of monoclonal Fab fragments with the desired specificity. (See, e.g., Huse, W.D. et al. (1989) Science 246:1275-1281.)

Various immunoassays may be used for screening to identify antibodies having the desired
15 specificity. Numerous protocols for competitive binding or immunoradiometric assays using either polyclonal or monoclonal antibodies with established specificities are well known in the art. Such immunoassays typically involve the measurement of complex formation between SECP and its specific antibody. A two-site, monoclonal-based immunoassay utilizing monoclonal antibodies reactive to two non-interfering SECP epitopes is generally used, but a competitive binding assay may
20 also be employed (Pound, supra).

Various methods such as Scatchard analysis in conjunction with radioimmunoassay techniques may be used to assess the affinity of antibodies for SECP. Affinity is expressed as an association constant, K_a , which is defined as the molar concentration of SECP-antibody complex divided by the molar concentrations of free antigen and free antibody under equilibrium conditions.
25 The K_a determined for a preparation of polyclonal antibodies, which are heterogeneous in their affinities for multiple SECP epitopes, represents the average affinity, or avidity, of the antibodies for SECP. The K_a determined for a preparation of monoclonal antibodies, which are monospecific for a particular SECP epitope, represents a true measure of affinity. High-affinity antibody preparations with K_a ranging from about 10^9 to 10^{12} L/mole are preferred for use in immunoassays in which the
30 SECP-antibody complex must withstand rigorous manipulations. Low-affinity antibody preparations with K_a ranging from about 10^6 to 10^7 L/mole are preferred for use in immunopurification and similar procedures which ultimately require dissociation of SECP, preferably in active form, from the antibody (Catty, D. (1988) Antibodies, Volume I: A Practical Approach, IRL Press, Washington DC; Liddell, J.E. and A. Cryer (1991) A Practical Guide to Monoclonal Antibodies, John Wiley & Sons,
35 New York NY).

The titer and avidity of polyclonal antibody preparations may be further evaluated to determine the quality and suitability of such preparations for certain downstream applications. For example, a polyclonal antibody preparation containing at least 1-2 mg specific antibody/ml, preferably 5-10 mg specific antibody/ml, is generally employed in procedures requiring precipitation of SECP-antibody complexes. Procedures for evaluating antibody specificity, titer, and avidity, and guidelines for antibody quality and usage in various applications, are generally available. (See, e.g., Catty, supra, and Coligan et al. supra.)

In another embodiment of the invention, the polynucleotides encoding SECP, or any fragment or complement thereof, may be used for therapeutic purposes. In one aspect, modifications of gene expression can be achieved by designing complementary sequences or antisense molecules (DNA, RNA, PNA, or modified oligonucleotides) to the coding or regulatory regions of the gene encoding SECP. Such technology is well known in the art, and antisense oligonucleotides or larger fragments can be designed from various locations along the coding or control regions of sequences encoding SECP. (See, e.g., Agrawal, S., ed. (1996) Antisense Therapeutics, Humana Press Inc., Totawa NJ.)

In therapeutic use, any gene delivery system suitable for introduction of the antisense sequences into appropriate target cells can be used. Antisense sequences can be delivered intracellularly in the form of an expression plasmid which, upon transcription, produces a sequence complementary to at least a portion of the cellular sequence encoding the target protein. (See, e.g., Slater, J.E. et al. (1998) *J. Allergy Clin. Immunol.* 102(3):469-475; and Scanlon, K.J. et al. (1995) 9(13):1288-1296.) Antisense sequences can also be introduced intracellularly through the use of viral vectors, such as retrovirus and adeno-associated virus vectors. (See, e.g., Miller, A.D. (1990) *Blood* 76:271; Ausubel, supra; Uckert, W. and W. Walther (1994) *Pharmacol. Ther.* 63(3):323-347.) Other gene delivery mechanisms include liposome-derived systems, artificial viral envelopes, and other systems known in the art. (See, e.g., Rossi, J.J. (1995) *Br. Med. Bull.* 51(1):217-225; Boado, R.J. et al. (1998) *J. Pharm. Sci.* 87(11):1308-1315; and Morris, M.C. et al. (1997) *Nucleic Acids Res.* 25(14):2730-2736.)

In another embodiment of the invention, polynucleotides encoding SECP may be used for somatic or germline gene therapy. Gene therapy may be performed to (i) correct a genetic deficiency (e.g., in the cases of severe combined immunodeficiency (SCID)-X1 disease characterized by X-linked inheritance (Cavazzana-Calvo, M. et al. (2000) *Science* 288:669-672), severe combined immunodeficiency syndrome associated with an inherited adenosine deaminase (ADA) deficiency (Blaese, R.M. et al. (1995) *Science* 270:475-480; Bordignon, C. et al. (1995) *Science* 270:470-475), cystic fibrosis (Zabner, J. et al. (1993) *Cell* 75:207-216; Crystal, R.G. et al. (1995) *Hum. Gene Therapy* 6:643-666; Crystal, R.G. et al. (1995) *Hum. Gene Therapy* 6:667-703), thalassemias, familial hypercholesterolemia, and hemophilia resulting from Factor VIII or Factor IX deficiencies (Crystal,

R.G. (1995) *Science* 270:404-410; Verma, I.M. and N. Somia (1997) *Nature* 389:239-242)), (ii) express a conditionally lethal gene product (e.g., in the case of cancers which result from unregulated cell proliferation), or (iii) express a protein which affords protection against intracellular parasites (e.g., against human retroviruses, such as human immunodeficiency virus (HIV) (Baltimore, D. (1988) *Nature* 335:395-396; Poeschla, E. et al. (1996) *Proc. Natl. Acad. Sci. USA* 93:11395-11399), hepatitis B or C virus (HBV, HCV); fungal parasites, such as Candida albicans and Paracoccidioides brasiliensis; and protozoan parasites such as Plasmodium falciparum and Trypanosoma cruzi). In the case where a genetic deficiency in SECP expression or regulation causes disease, the expression of SECP from an appropriate population of transduced cells may alleviate the clinical manifestations caused by the genetic deficiency.

In a further embodiment of the invention, diseases or disorders caused by deficiencies in SECP are treated by constructing mammalian expression vectors encoding SECP and introducing these vectors by mechanical means into SECP-deficient cells. Mechanical transfer technologies for use with cells in vivo or ex vitro include (i) direct DNA microinjection into individual cells, (ii) ballistic gold particle delivery, (iii) liposome-mediated transfection, (iv) receptor-mediated gene transfer, and (v) the use of DNA transposons (Morgan, R.A. and W.F. Anderson (1993) *Annu. Rev. Biochem.* 62:191-217; Ivics, Z. (1997) *Cell* 91:501-510; Boulay, J-L. and H. R  c  pon (1998) *Curr. Opin. Biotechnol.* 9:445-450).

Expression vectors that may be effective for the expression of SECP include, but are not limited to, the PCDNA 3.1, EPITAG, PRCCMV2, PREP, PVAX, PCR2-TOPOTA vectors (Invitrogen, Carlsbad CA), PCMV-SCRIPT, PCMV-TAG, PEGSH/PERV (Stratagene, La Jolla CA), and PTET-OFF, PTET-ON, PTRE2, PTRE2-LUC, PTK-HYG (Clontech, Palo Alto CA). SECP may be expressed using (i) a constitutively active promoter, (e.g., from cytomegalovirus (CMV), Rous sarcoma virus (RSV), SV40 virus, thymidine kinase (TK), or β -actin genes), (ii) an inducible promoter (e.g., the tetracycline-regulated promoter (Gossen, M. and H. Bujard (1992) *Proc. Natl. Acad. Sci. USA* 89:5547-5551; Gossen, M. et al. (1995) *Science* 268:1766-1769; Rossi, F.M.V. and H.M. Blau (1998) *Curr. Opin. Biotechnol.* 9:451-456), commercially available in the T-REX plasmid (Invitrogen)); the ecdysone-inducible promoter (available in the plasmids PVGRXR and PIND; Invitrogen); the FK506/rapamycin inducible promoter; or the RU486/mifepristone inducible promoter (Rossi, F.M.V. and Blau, H.M. supra), or (iii) a tissue-specific promoter or the native promoter of the endogenous gene encoding SECP from a normal individual.

Commercially available liposome transformation kits (e.g., the PERFECT LIPID TRANSFECTION KIT, available from Invitrogen) allow one with ordinary skill in the art to deliver polynucleotides to target cells in culture and require minimal effort to optimize experimental parameters. In the alternative, transformation is performed using the calcium phosphate method

(Graham, F.L. and A.J. Eb (1973) *Virology* 52:456-467), or by electroporation (Neumann, E. et al. (1982) *EMBO J.* 1:841-845). The introduction of DNA to primary cells requires modification of these standardized mammalian transfection protocols.

In another embodiment of the invention, diseases or disorders caused by genetic defects with respect to SECP expression are treated by constructing a retrovirus vector consisting of (i) the polynucleotide encoding SECP under the control of an independent promoter or the retrovirus long terminal repeat (LTR) promoter, (ii) appropriate RNA packaging signals, and (iii) a Rev-responsive element (RRE) along with additional retrovirus *cis*-acting RNA sequences and coding sequences required for efficient vector propagation. Retrovirus vectors (e.g., PFB and PFBNEO) are commercially available (Stratagene) and are based on published data (Riviere, I. et al. (1995) *Proc. Natl. Acad. Sci. USA* 92:6733-6737), incorporated by reference herein. The vector is propagated in an appropriate vector producing cell line (VPCL) that expresses an envelope gene with a tropism for receptors on the target cells or a promiscuous envelope protein such as VSVg (Armentano, D. et al. (1987) *J. Virol.* 61:1647-1650; Bender, M.A. et al. (1987) *J. Virol.* 61:1639-1646; Adam, M.A. and A.D. Miller (1988) *J. Virol.* 62:3802-3806; Dull, T. et al. (1998) *J. Virol.* 72:8463-8471; Zufferey, R. et al. (1998) *J. Virol.* 72:9873-9880). U.S. Patent No. 5,910,434 to Rigg ("Method for obtaining retrovirus packaging cell lines producing high transducing efficiency retroviral supernatant") discloses a method for obtaining retrovirus packaging cell lines and is hereby incorporated by reference. Propagation of retrovirus vectors, transduction of a population of cells (e.g., CD4⁺ T-cells), and the return of transduced cells to a patient are procedures well known to persons skilled in the art of gene therapy and have been well documented (Ranga, U. et al. (1997) *J. Virol.* 71:7020-7029; Bauer, G. et al. (1997) *Blood* 89:2259-2267; Bonyhadi, M.L. (1997) *J. Virol.* 71:4707-4716; Ranga, U. et al. (1998) *Proc. Natl. Acad. Sci. USA* 95:1201-1206; Su, L. (1997) *Blood* 89:2283-2290).

In the alternative, an adenovirus-based gene therapy delivery system is used to deliver polynucleotides encoding SECP to cells which have one or more genetic abnormalities with respect to the expression of SECP. The construction and packaging of adenovirus-based vectors are well known to those with ordinary skill in the art. Replication defective adenovirus vectors have proven to be versatile for importing genes encoding immunoregulatory proteins into intact islets in the pancreas (Csete, M.E. et al. (1995) *Transplantation* 27:263-268). Potentially useful adenoviral vectors are described in U.S. Patent No. 5,707,618 to Armentano ("Adenovirus vectors for gene therapy"), hereby incorporated by reference. For adenoviral vectors, see also Antinozzi, P.A. et al. (1999) *Annu. Rev. Nutr.* 19:511-544 and Verma, I.M. and N. Somia (1997) *Nature* 18:389:239-242, both incorporated by reference herein.

In another alternative, a herpes-based, gene therapy delivery system is used to deliver

polynucleotides encoding SECP to target cells which have one or more genetic abnormalities with respect to the expression of SECP. The use of herpes simplex virus (HSV)-based vectors may be especially valuable for introducing SECP to cells of the central nervous system, for which HSV has a tropism. The construction and packaging of herpes-based vectors are well known to those with ordinary skill in the art. A replication-competent herpes simplex virus (HSV) type 1-based vector has been used to deliver a reporter gene to the eyes of primates (Liu, X. et al. (1999) Exp. Eye Res. 169:385-395). The construction of a HSV-1 virus vector has also been disclosed in detail in U.S. Patent No. 5,804,413 to DeLuca ("Herpes simplex virus strains for gene transfer"), which is hereby incorporated by reference. U.S. Patent No. 5,804,413 teaches the use of recombinant HSV d92 which consists of a genome containing at least one exogenous gene to be transferred to a cell under the control of the appropriate promoter for purposes including human gene therapy. Also taught by this patent are the construction and use of recombinant HSV strains deleted for ICP4, ICP27 and ICP22. For HSV vectors, see also Goins, W.F. et al. (1999) J. Virol. 73:519-532 and Xu, H. et al. (1994) Dev. Biol. 163:152-161, hereby incorporated by reference. The manipulation of cloned herpesvirus sequences, the generation of recombinant virus following the transfection of multiple plasmids containing different segments of the large herpesvirus genomes, the growth and propagation of herpesvirus, and the infection of cells with herpesvirus are techniques well known to those of ordinary skill in the art.

In another alternative, an alphavirus (positive, single-stranded RNA virus) vector is used to deliver polynucleotides encoding SECP to target cells. The biology of the prototypic alphavirus, Semliki Forest Virus (SFV), has been studied extensively and gene transfer vectors have been based on the SFV genome (Garoff, H. and K.-J. Li (1998) Curr. Opin. Biotechnol. 9:464-469). During alphavirus RNA replication, a subgenomic RNA is generated that normally encodes the viral capsid proteins. This subgenomic RNA replicates to higher levels than the full length genomic RNA, resulting in the overproduction of capsid proteins relative to the viral proteins with enzymatic activity (e.g., protease and polymerase). Similarly, inserting the coding sequence for SECP into the alphavirus genome in place of the capsid-coding region results in the production of a large number of SECP-coding RNAs and the synthesis of high levels of SECP in vector transduced cells. While alphavirus infection is typically associated with cell lysis within a few days, the ability to establish a persistent infection in hamster normal kidney cells (BHK-21) with a variant of Sindbis virus (SIN) indicates that the lytic replication of alphaviruses can be altered to suit the needs of the gene therapy application (Dryga, S.A. et al. (1997) Virology 228:74-83). The wide host range of alphaviruses will allow the introduction of SECP into a variety of cell types. The specific transduction of a subset of cells in a population may require the sorting of cells prior to transduction. The methods of manipulating infectious cDNA clones of alphaviruses, performing alphavirus cDNA and RNA

transfections, and performing alphavirus infections, are well known to those with ordinary skill in the art.

Oligonucleotides derived from the transcription initiation site, e.g., between about positions -10 and +10 from the start site, may also be employed to inhibit gene expression. Similarly, inhibition can be achieved using triple helix base-pairing methodology. Triple helix pairing is useful because it causes inhibition of the ability of the double helix to open sufficiently for the binding of polymerases, transcription factors, or regulatory molecules. Recent therapeutic advances using triplex DNA have been described in the literature. (See, e.g., Gee, J.E. et al. (1994) in Huber, B.E. and B.I. Carr, Molecular and Immunologic Approaches, Futura Publishing, Mt. Kisco NY, pp. 163-177.) A complementary sequence or antisense molecule may also be designed to block translation of mRNA by preventing the transcript from binding to ribosomes.

Ribozymes, enzymatic RNA molecules, may also be used to catalyze the specific cleavage of RNA. The mechanism of ribozyme action involves sequence-specific hybridization of the ribozyme molecule to complementary target RNA, followed by endonucleolytic cleavage. For example, engineered hammerhead motif ribozyme molecules may specifically and efficiently catalyze endonucleolytic cleavage of sequences encoding SECP.

Specific ribozyme cleavage sites within any potential RNA target are initially identified by scanning the target molecule for ribozyme cleavage sites, including the following sequences: GUA, GUU, and GUC. Once identified, short RNA sequences of between 15 and 20 ribonucleotides, corresponding to the region of the target gene containing the cleavage site, may be evaluated for secondary structural features which may render the oligonucleotide inoperable. The suitability of candidate targets may also be evaluated by testing accessibility to hybridization with complementary oligonucleotides using ribonuclease protection assays.

Complementary ribonucleic acid molecules and ribozymes of the invention may be prepared by any method known in the art for the synthesis of nucleic acid molecules. These include techniques for chemically synthesizing oligonucleotides such as solid phase phosphoramidite chemical synthesis. Alternatively, RNA molecules may be generated by in vitro and in vivo transcription of DNA sequences encoding SECP. Such DNA sequences may be incorporated into a wide variety of vectors with suitable RNA polymerase promoters such as T7 or SP6. Alternatively, these cDNA constructs that synthesize complementary RNA, constitutively or inducibly, can be introduced into cell lines, cells, or tissues.

RNA molecules may be modified to increase intracellular stability and half-life. Possible modifications include, but are not limited to, the addition of flanking sequences at the 5' and/or 3' ends of the molecule, or the use of phosphorothioate or 2' O-methyl rather than phosphodiesterase linkages within the backbone of the molecule. This concept is inherent in the production of PNAs

and can be extended in all of these molecules by the inclusion of nontraditional bases such as inosine, queosine, and wybutosine, as well as acetyl-, methyl-, thio-, and similarly modified forms of adenine, cytidine, guanine, thymine, and uridine which are not as easily recognized by endogenous endonucleases.

5 An additional embodiment of the invention encompasses a method for screening for a compound which is effective in altering expression of a polynucleotide encoding SECP. Compounds which may be effective in altering expression of a specific polynucleotide may include, but are not limited to, oligonucleotides, antisense oligonucleotides, triple helix-forming oligonucleotides, transcription factors and other polypeptide transcriptional regulators, and non-macromolecular
10 chemical entities which are capable of interacting with specific polynucleotide sequences. Effective compounds may alter polynucleotide expression by acting as either inhibitors or promoters of polynucleotide expression. Thus, in the treatment of disorders associated with increased SECP expression or activity, a compound which specifically inhibits expression of the polynucleotide encoding SECP may be therapeutically useful, and in the treatment of disorders associated with
15 decreased SECP expression or activity, a compound which specifically promotes expression of the polynucleotide encoding SECP may be therapeutically useful.

At least one, and up to a plurality, of test compounds may be screened for effectiveness in altering expression of a specific polynucleotide. A test compound may be obtained by any method commonly known in the art, including chemical modification of a compound known to be effective in
20 altering polynucleotide expression; selection from an existing, commercially-available or proprietary library of naturally-occurring or non-natural chemical compounds; rational design of a compound based on chemical and/or structural properties of the target polynucleotide; and selection from a library of chemical compounds created combinatorially or randomly. A sample comprising a polynucleotide encoding SECP is exposed to at least one test compound thus obtained. The sample
25 may comprise, for example, an intact or permeabilized cell, or an *in vitro* cell-free or reconstituted biochemical system. Alterations in the expression of a polynucleotide encoding SECP are assayed by any method commonly known in the art. Typically, the expression of a specific nucleotide is detected by hybridization with a probe having a nucleotide sequence complementary to the sequence of the polynucleotide encoding SECP. The amount of hybridization may be quantified, thus forming
30 the basis for a comparison of the expression of the polynucleotide both with and without exposure to one or more test compounds. Detection of a change in the expression of a polynucleotide exposed to a test compound indicates that the test compound is effective in altering the expression of the polynucleotide. A screen for a compound effective in altering expression of a specific polynucleotide can be carried out, for example, using a *Schizosaccharomyces pombe* gene expression system
35 (Atkins, D. et al. (1999) U.S. Patent No. 5,932,435; Arndt, G.M. et al. (2000) Nucleic Acids Res.

28:E15) or a human cell line such as HeLa cell (Clarke, M.L. et al. (2000) Biochem. Biophys. Res. Commun. 268:8-13). A particular embodiment of the present invention involves screening a combinatorial library of oligonucleotides (such as deoxyribonucleotides, ribonucleotides, peptide nucleic acids, and modified oligonucleotides) for antisense activity against a specific polynucleotide sequence (Bruce, T.W. et al. (1997) U.S. Patent No. 5,686,242; Bruce, T.W. et al. (2000) U.S. Patent No. 6,022,691).

Many methods for introducing vectors into cells or tissues are available and equally suitable for use in vivo, in vitro, and ex vivo. For ex vivo therapy, vectors may be introduced into stem cells taken from the patient and clonally propagated for autologous transplant back into that same patient. Delivery by transfection, by liposome injections, or by polycationic amino polymers may be achieved using methods which are well known in the art. (See, e.g., Goldman, C.K. et al. (1997) Nat. Biotechnol. 15:462-466.)

Any of the therapeutic methods described above may be applied to any subject in need of such therapy, including, for example, mammals such as humans, dogs, cats, cows, horses, rabbits, and monkeys.

An additional embodiment of the invention relates to the administration of a composition which generally comprises an active ingredient formulated with a pharmaceutically acceptable excipient. Excipients may include, for example, sugars, starches, celluloses, gums, and proteins. Various formulations are commonly known and are thoroughly discussed in the latest edition of Remington's Pharmaceutical Sciences (Maack Publishing, Easton PA). Such compositions may consist of SECP, antibodies to SECP, and mimetics, agonists, antagonists, or inhibitors of SECP.

The compositions utilized in this invention may be administered by any number of routes including, but not limited to, oral, intravenous, intramuscular, intra-arterial, intramedullary, intrathecal, intraventricular, pulmonary, transdermal, subcutaneous, intraperitoneal, intranasal, enteral, topical, sublingual, or rectal means.

Compositions for pulmonary administration may be prepared in liquid or dry powder form. These compositions are generally aerosolized immediately prior to inhalation by the patient. In the case of small molecules (e.g. traditional low molecular weight organic drugs), aerosol delivery of fast-acting formulations is well-known in the art. In the case of macromolecules (e.g. larger peptides and proteins), recent developments in the field of pulmonary delivery via the alveolar region of the lung have enabled the practical delivery of drugs such as insulin to blood circulation (see, e.g., Patton, J.S. et al., U.S. Patent No. 5,997,848). Pulmonary delivery has the advantage of administration without needle injection, and obviates the need for potentially toxic penetration enhancers.

Compositions suitable for use in the invention include compositions wherein the active ingredients are contained in an effective amount to achieve the intended purpose. The determination

of an effective dose is well within the capability of those skilled in the art.

Specialized forms of compositions may be prepared for direct intracellular delivery of macromolecules comprising SECP or fragments thereof. For example, liposome preparations containing a cell-impermeable macromolecule may promote cell fusion and intracellular delivery of the macromolecule. Alternatively, SECP or a fragment thereof may be joined to a short cationic N-terminal portion from the HIV Tat-1 protein. Fusion proteins thus generated have been found to transduce into the cells of all tissues, including the brain, in a mouse model system (Schwarze, S.R. et al. (1999) Science 285:1569-1572).

For any compound, the therapeutically effective dose can be estimated initially either in cell culture assays, e.g., of neoplastic cells, or in animal models such as mice, rats, rabbits, dogs, monkeys, or pigs. An animal model may also be used to determine the appropriate concentration range and route of administration. Such information can then be used to determine useful doses and routes for administration in humans.

A therapeutically effective dose refers to that amount of active ingredient, for example SECP or fragments thereof, antibodies of SECP, and agonists, antagonists or inhibitors of SECP, which ameliorates the symptoms or condition. Therapeutic efficacy and toxicity may be determined by standard pharmaceutical procedures in cell cultures or with experimental animals, such as by calculating the ED_{50} (the dose therapeutically effective in 50% of the population) or LD_{50} (the dose lethal to 50% of the population) statistics. The dose ratio of toxic to therapeutic effects is the therapeutic index, which can be expressed as the LD_{50}/ED_{50} ratio. Compositions which exhibit large therapeutic indices are preferred. The data obtained from cell culture assays and animal studies are used to formulate a range of dosage for human use. The dosage contained in such compositions is preferably within a range of circulating concentrations that includes the ED_{50} with little or no toxicity. The dosage varies within this range depending upon the dosage form employed, the sensitivity of the patient, and the route of administration.

The exact dosage will be determined by the practitioner, in light of factors related to the subject requiring treatment. Dosage and administration are adjusted to provide sufficient levels of the active moiety or to maintain the desired effect. Factors which may be taken into account include the severity of the disease state, the general health of the subject, the age, weight, and gender of the subject, time and frequency of administration, drug combination(s), reaction sensitivities, and response to therapy. Long-acting compositions may be administered every 3 to 4 days, every week, or biweekly depending on the half-life and clearance rate of the particular formulation.

Normal dosage amounts may vary from about 0.1 μg to 100,000 μg , up to a total dose of about 1 gram, depending upon the route of administration. Guidance as to particular dosages and methods of delivery is provided in the literature and generally available to practitioners in the art.

Those skilled in the art will employ different formulations for nucleotides than for proteins or their inhibitors. Similarly, delivery of polynucleotides or polypeptides will be specific to particular cells, conditions, locations, etc.

DIAGNOSTICS

5 In another embodiment, antibodies which specifically bind SECP may be used for the diagnosis of disorders characterized by expression of SECP, or in assays to monitor patients being treated with SECP or agonists, antagonists, or inhibitors of SECP. Antibodies useful for diagnostic purposes may be prepared in the same manner as described above for therapeutics. Diagnostic assays for SECP include methods which utilize the antibody and a label to detect SECP in human body
10 fluids or in extracts of cells or tissues. The antibodies may be used with or without modification, and may be labeled by covalent or non-covalent attachment of a reporter molecule. A wide variety of reporter molecules, several of which are described above, are known in the art and may be used.

A variety of protocols for measuring SECP, including ELISAs, RIAs, and FACS, are known in the art and provide a basis for diagnosing altered or abnormal levels of SECP expression. Normal
15 or standard values for SECP expression are established by combining body fluids or cell extracts taken from normal mammalian subjects, for example, human subjects, with antibodies to SECP under conditions suitable for complex formation. The amount of standard complex formation may be quantitated by various methods, such as photometric means. Quantities of SECP expressed in subject, control, and disease samples from biopsied tissues are compared with the standard values.
20 Deviation between standard and subject values establishes the parameters for diagnosing disease.

In another embodiment of the invention, the polynucleotides encoding SECP may be used for diagnostic purposes. The polynucleotides which may be used include oligonucleotide sequences, complementary RNA and DNA molecules, and PNAs. The polynucleotides may be used to detect and quantify gene expression in biopsied tissues in which expression of SECP may be correlated with
25 disease. The diagnostic assay may be used to determine absence, presence, and excess expression of SECP, and to monitor regulation of SECP levels during therapeutic intervention.

In one aspect, hybridization with PCR probes which are capable of detecting polynucleotide sequences, including genomic sequences, encoding SECP or closely related molecules may be used to identify nucleic acid sequences which encode SECP. The specificity of the probe, whether it is made
30 from a highly specific region, e.g., the 5' regulatory region, or from a less specific region, e.g., a conserved motif, and the stringency of the hybridization or amplification will determine whether the probe identifies only naturally occurring sequences encoding SECP, allelic variants, or related sequences.

Probes may also be used for the detection of related sequences, and may have at least 50%
35 sequence identity to any of the SECP encoding sequences. The hybridization probes of the subject

invention may be DNA or RNA and may be derived from the sequence of SEQ ID NO:68-134 or from genomic sequences including promoters, enhancers, and introns of the SECP gene.

Means for producing specific hybridization probes for DNAs encoding SECP include the cloning of polynucleotide sequences encoding SECP or SECP derivatives into vectors for the production of mRNA probes. Such vectors are known in the art, are commercially available, and may be used to synthesize RNA probes in vitro by means of the addition of the appropriate RNA polymerases and the appropriate labeled nucleotides. Hybridization probes may be labeled by a variety of reporter groups, for example, by radionuclides such as ^{32}P or ^{35}S , or by enzymatic labels, such as alkaline phosphatase coupled to the probe via avidin/biotin coupling systems, and the like.

Polynucleotide sequences encoding SECP may be used for the diagnosis of disorders associated with expression of SECP. Examples of such disorders include, but are not limited to, a cell proliferative disorder such as actinic keratosis, arteriosclerosis, atherosclerosis, bursitis, cirrhosis, hepatitis, mixed connective tissue disease (MCTD), myelofibrosis, paroxysmal nocturnal hemoglobinuria, polycythemia vera, psoriasis, primary thrombocythemia, and cancers including adenocarcinoma, leukemia, lymphoma, melanoma, myeloma, sarcoma, teratocarcinoma, and, in particular, a cancer of the adrenal gland, bladder, bone, bone marrow, brain, breast, cervix, gall bladder, ganglia, gastrointestinal tract, heart, kidney, liver, lung, muscle, ovary, pancreas, parathyroid, penis, prostate, salivary glands, skin, spleen, testis, thymus, thyroid, and uterus; an autoimmune/inflammatory disorder such as acquired immunodeficiency syndrome (AIDS), Addison's disease, adult respiratory distress syndrome, allergies, ankylosing spondylitis, amyloidosis, anemia, asthma, atherosclerosis, autoimmune hemolytic anemia, autoimmune thyroiditis, autoimmune polyendocrinopathy-candidiasis-ectodermal dystrophy (APECED), bronchitis, cholecystitis, contact dermatitis, Crohn's disease, atopic dermatitis, dermatomyositis, diabetes mellitus, emphysema, episodic lymphopenia with lymphocytotoxins, erythroblastosis fetalis, erythema nodosum, atrophic gastritis, glomerulonephritis, Goodpasture's syndrome, gout, Graves' disease, Hashimoto's thyroiditis, hypereosinophilia, irritable bowel syndrome, multiple sclerosis, myasthenia gravis, myocardial or pericardial inflammation, osteoarthritis, osteoporosis, pancreatitis, polymyositis, psoriasis, Reiter's syndrome, rheumatoid arthritis, scleroderma, Sjögren's syndrome, systemic anaphylaxis, systemic lupus erythematosus, systemic sclerosis, thrombocytopenic purpura, ulcerative colitis, uveitis, Werner syndrome, complications of cancer, hemodialysis, and extracorporeal circulation, viral, bacterial, fungal, parasitic, protozoal, and helminthic infections, and trauma; a cardiovascular disorder such as congestive heart failure, ischemic heart disease, angina pectoris, myocardial infarction, hypertensive heart disease, degenerative valvular heart disease, calcific aortic valve stenosis, congenitally bicuspid aortic valve, mitral annular calcification, mitral valve prolapse, rheumatic fever and rheumatic heart disease, infective endocarditis, nonbacterial thrombotic

endocarditis, endocarditis of systemic lupus erythematosus, carcinoid heart disease, cardiomyopathy, myocarditis, pericarditis, neoplastic heart disease, congenital heart disease, complications of cardiac transplantation, arteriovenous fistula, atherosclerosis, hypertension, vasculitis, Raynaud's disease, aneurysms, arterial dissections, varicose veins, thrombophlebitis and phlebothrombosis, vascular tumors, and complications of thrombolysis, balloon angioplasty, vascular replacement, and coronary artery bypass graft surgery; a neurological disorder such as epilepsy, ischemic cerebrovascular disease, stroke, cerebral neoplasms, Alzheimer's disease, Pick's disease, Huntington's disease, dementia, Parkinson's disease and other extrapyramidal disorders, amyotrophic lateral sclerosis and other motor neuron disorders, progressive neural muscular atrophy, retinitis pigmentosa, hereditary ataxias, multiple sclerosis and other demyelinating diseases, bacterial and viral meningitis, brain abscess, subdural empyema, epidural abscess, suppurative intracranial thrombophlebitis, myelitis and radiculitis, viral central nervous system disease, prion diseases including kuru, Creutzfeldt-Jakob disease, and Gerstmann-Straussler-Scheinker syndrome, fatal familial insomnia, nutritional and metabolic diseases of the nervous system, neurofibromatosis, tuberous sclerosis, cerebelloretinal hemangioblastomatosis, encephalotrigeminal syndrome, mental retardation and other developmental disorders of the central nervous system including Down syndrome, cerebral palsy, neuroskeletal disorders, autonomic nervous system disorders, cranial nerve disorders, spinal cord diseases, muscular dystrophy and other neuromuscular disorders, peripheral nervous system disorders, dermatomyositis and polymyositis, inherited, metabolic, endocrine, and toxic myopathies, myasthenia gravis, periodic paralysis, mental disorders including mood, anxiety, and schizophrenic disorders, seasonal affective disorder (SAD), akathisia, amnesia, catatonia, diabetic neuropathy, tardive dyskinesia, dystonias, paranoid psychoses, postherpetic neuralgia, Tourette's disorder, progressive supranuclear palsy, corticobasal degeneration, and familial frontotemporal dementia; and a developmental disorder such as renal tubular acidosis, anemia, Cushing's syndrome, achondroplastic dwarfism, Duchenne and Becker muscular dystrophy, epilepsy, gonadal dysgenesis, WAGR syndrome (Wilms' tumor, aniridia, genitourinary abnormalities, and mental retardation), Smith-Magenis syndrome, myelodysplastic syndrome, hereditary mucoepithelial dysplasia, hereditary keratodermas, hereditary neuropathies such as Charcot-Marie-Tooth disease and neurofibromatosis, hypothyroidism, hydrocephalus, seizure disorders such as Sydenham's chorea and cerebral palsy, spina bifida, anencephaly, craniorachischisis, congenital glaucoma, cataract, and sensorineural hearing loss. The polynucleotide sequences encoding SECP may be used in Southern or northern analysis, dot blot, or other membrane-based technologies; in PCR technologies; in dipstick, pin, and multiformat ELISA-like assays; and in microarrays utilizing fluids or tissues from patients to detect altered SECP expression. Such qualitative or quantitative methods are well known in the art.

In a particular aspect, the nucleotide sequences encoding SECP may be useful in assays that

detect the presence of associated disorders, particularly those mentioned above. The nucleotide sequences encoding SECP may be labeled by standard methods and added to a fluid or tissue sample from a patient under conditions suitable for the formation of hybridization complexes. After a suitable incubation period, the sample is washed and the signal is quantified and compared with a standard value. If the amount of signal in the patient sample is significantly altered in comparison to a control sample then the presence of altered levels of nucleotide sequences encoding SECP in the sample indicates the presence of the associated disorder. Such assays may also be used to evaluate the efficacy of a particular therapeutic treatment regimen in animal studies, in clinical trials, or to monitor the treatment of an individual patient.

10 In order to provide a basis for the diagnosis of a disorder associated with expression of SECP, a normal or standard profile for expression is established. This may be accomplished by combining body fluids or cell extracts taken from normal subjects, either animal or human, with a sequence, or a fragment thereof, encoding SECP, under conditions suitable for hybridization or amplification. Standard hybridization may be quantified by comparing the values obtained from normal subjects with values from an experiment in which a known amount of a substantially purified polynucleotide is used. Standard values obtained in this manner may be compared with values obtained from samples from patients who are symptomatic for a disorder. Deviation from standard values is used to establish the presence of a disorder.

Once the presence of a disorder is established and a treatment protocol is initiated, hybridization assays may be repeated on a regular basis to determine if the level of expression in the patient begins to approximate that which is observed in the normal subject. The results obtained from successive assays may be used to show the efficacy of treatment over a period ranging from several days to months.

With respect to cancer, the presence of an abnormal amount of transcript (either under- or overexpressed) in biopsied tissue from an individual may indicate a predisposition for the development of the disease, or may provide a means for detecting the disease prior to the appearance of actual clinical symptoms. A more definitive diagnosis of this type may allow health professionals to employ preventative measures or aggressive treatment earlier thereby preventing the development or further progression of the cancer.

30 Additional diagnostic uses for oligonucleotides designed from the sequences encoding SECP may involve the use of PCR. These oligomers may be chemically synthesized, generated enzymatically, or produced in vitro. Oligomers will preferably contain a fragment of a polynucleotide encoding SECP, or a fragment of a polynucleotide complementary to the polynucleotide encoding SECP, and will be employed under optimized conditions for identification of a specific gene or condition. Oligomers may also be employed under less stringent conditions for detection or

quantification of closely related DNA or RNA sequences.

In a particular aspect, oligonucleotide primers derived from the polynucleotide sequences encoding SECP may be used to detect single nucleotide polymorphisms (SNPs). SNPs are substitutions, insertions and deletions that are a frequent cause of inherited or acquired genetic disease in humans. Methods of SNP detection include, but are not limited to, single-stranded conformation polymorphism (SSCP) and fluorescent SSCP (fSSCP) methods. In SSCP, oligonucleotide primers derived from the polynucleotide sequences encoding SECP are used to amplify DNA using the polymerase chain reaction (PCR). The DNA may be derived, for example, from diseased or normal tissue, biopsy samples, bodily fluids, and the like. SNPs in the DNA cause differences in the secondary and tertiary structures of PCR products in single-stranded form, and these differences are detectable using gel electrophoresis in non-denaturing gels. In fSSCP, the oligonucleotide primers are fluorescently labeled, which allows detection of the amplimers in high-throughput equipment such as DNA sequencing machines. Additionally, sequence database analysis methods, termed *in silico* SNP (isSNP), are capable of identifying polymorphisms by comparing the sequence of individual overlapping DNA fragments which assemble into a common consensus sequence. These computer-based methods filter out sequence variations due to laboratory preparation of DNA and sequencing errors using statistical models and automated analyses of DNA sequence chromatograms. In the alternative, SNPs may be detected and characterized by mass spectrometry using, for example, the high throughput MASSARRAY system (Sequenom, Inc., San Diego CA).

Methods which may also be used to quantify the expression of SECP include radiolabeling or biotinylating nucleotides, coamplification of a control nucleic acid, and interpolating results from standard curves. (See, e.g., Melby, P.C. et al. (1993) *J. Immunol. Methods* 159:235-244; Duplaa, C. et al. (1993) *Anal. Biochem.* 212:229-236.) The speed of quantitation of multiple samples may be accelerated by running the assay in a high-throughput format where the oligomer or polynucleotide of interest is presented in various dilutions and a spectrophotometric or colorimetric response gives rapid quantitation.

In further embodiments, oligonucleotides or longer fragments derived from any of the polynucleotide sequences described herein may be used as elements on a microarray. The microarray can be used in transcript imaging techniques which monitor the relative expression levels of large numbers of genes simultaneously as described below. The microarray may also be used to identify genetic variants, mutations, and polymorphisms. This information may be used to determine gene function, to understand the genetic basis of a disorder, to diagnose a disorder, to monitor progression/regression of disease as a function of gene expression, and to develop and monitor the activities of therapeutic agents in the treatment of disease. In particular, this information may be used to develop a pharmacogenomic profile of a patient in order to select the most appropriate and

effective treatment regimen for that patient. For example, therapeutic agents which are highly effective and display the fewest side effects may be selected for a patient based on his/her pharmacogenomic profile.

In another embodiment, SECP, fragments of SECP, or antibodies specific for SECP may be used as elements on a microarray. The microarray may be used to monitor or measure protein-protein interactions, drug-target interactions, and gene expression profiles, as described above.

A particular embodiment relates to the use of the polynucleotides of the present invention to generate a transcript image of a tissue or cell type. A transcript image represents the global pattern of gene expression by a particular tissue or cell type. Global gene expression patterns are analyzed by quantifying the number of expressed genes and their relative abundance under given conditions and at a given time. (See Seilhamer et al., "Comparative Gene Transcript Analysis," U.S. Patent No. 5,840,484, expressly incorporated by reference herein.) Thus a transcript image may be generated by hybridizing the polynucleotides of the present invention or their complements to the totality of transcripts or reverse transcripts of a particular tissue or cell type. In one embodiment, the hybridization takes place in high-throughput format, wherein the polynucleotides of the present invention or their complements comprise a subset of a plurality of elements on a microarray. The resultant transcript image would provide a profile of gene activity.

Transcript images may be generated using transcripts isolated from tissues, cell lines, biopsies, or other biological samples. The transcript image may thus reflect gene expression *in vivo*, as in the case of a tissue or biopsy sample, or *in vitro*, as in the case of a cell line.

Transcript images which profile the expression of the polynucleotides of the present invention may also be used in conjunction with *in vitro* model systems and preclinical evaluation of pharmaceuticals, as well as toxicological testing of industrial and naturally-occurring environmental compounds. All compounds induce characteristic gene expression patterns, frequently termed molecular fingerprints or toxicant signatures, which are indicative of mechanisms of action and toxicity (Nuwaysir, E.F. et al. (1999) Mol. Carcinog. 24:153-159; Steiner, S. and N.L. Anderson (2000) Toxicol. Lett. 112-113:467-471, expressly incorporated by reference herein). If a test compound has a signature similar to that of a compound with known toxicity, it is likely to share those toxic properties. These fingerprints or signatures are most useful and refined when they contain expression information from a large number of genes and gene families. Ideally, a genome-wide measurement of expression provides the highest quality signature. Even genes whose expression is not altered by any tested compounds are important as well, as the levels of expression of these genes are used to normalize the rest of the expression data. The normalization procedure is useful for comparison of expression data after treatment with different compounds. While the assignment of gene function to elements of a toxicant signature aids in interpretation of toxicity mechanisms,

knowledge of gene function is not necessary for the statistical matching of signatures which leads to prediction of toxicity. (See, for example, Press Release 00-02 from the National Institute of Environmental Health Sciences, released February 29, 2000, available at <http://www.niehs.nih.gov/oc/news/toxchip.htm>.) Therefore, it is important and desirable in

5 toxicological screening using toxicant signatures to include all expressed gene sequences.

In one embodiment, the toxicity of a test compound is assessed by treating a biological sample containing nucleic acids with the test compound. Nucleic acids that are expressed in the treated biological sample are hybridized with one or more probes specific to the polynucleotides of the present invention, so that transcript levels corresponding to the polynucleotides of the present
10 invention may be quantified. The transcript levels in the treated biological sample are compared with levels in an untreated biological sample. Differences in the transcript levels between the two samples are indicative of a toxic response caused by the test compound in the treated sample.

Another particular embodiment relates to the use of the polypeptide sequences of the present invention to analyze the proteome of a tissue or cell type. The term proteome refers to the global
15 pattern of protein expression in a particular tissue or cell type. Each protein component of a proteome can be subjected individually to further analysis. Proteome expression patterns, or profiles, are analyzed by quantifying the number of expressed proteins and their relative abundance under given conditions and at a given time. A profile of a cell's proteome may thus be generated by separating and analyzing the polypeptides of a particular tissue or cell type. In one embodiment, the
20 separation is achieved using two-dimensional gel electrophoresis, in which proteins from a sample are separated by isoelectric focusing in the first dimension, and then according to molecular weight by sodium dodecyl sulfate slab gel electrophoresis in the second dimension (Steiner and Anderson, *supra*). The proteins are visualized in the gel as discrete and uniquely positioned spots, typically by staining the gel with an agent such as Coomassie Blue or silver or fluorescent stains. The optical
25 density of each protein spot is generally proportional to the level of the protein in the sample. The optical densities of equivalently positioned protein spots from different samples, for example, from biological samples either treated or untreated with a test compound or therapeutic agent, are compared to identify any changes in protein spot density related to the treatment. The proteins in the spots are partially sequenced using, for example, standard methods employing chemical or enzymatic
30 cleavage followed by mass spectrometry. The identity of the protein in a spot may be determined by comparing its partial sequence, preferably of at least 5 contiguous amino acid residues, to the polypeptide sequences of the present invention. In some cases, further sequence data may be obtained for definitive protein identification.

A proteomic profile may also be generated using antibodies specific for SECP to quantify the
35 levels of SECP expression. In one embodiment, the antibodies are used as elements on a microarray,

and protein expression levels are quantified by exposing the microarray to the sample and detecting the levels of protein bound to each array element (Lueking, A. et al. (1999) Anal. Biochem. 270:103-111; Mendoz, L.G. et al. (1999) Biotechniques 27:778-788). Detection may be performed by a variety of methods known in the art, for example, by reacting the proteins in the sample with a thiol-
5 or amino-reactive fluorescent compound and detecting the amount of fluorescence bound at each array element.

Toxicant signatures at the proteome level are also useful for toxicological screening, and should be analyzed in parallel with toxicant signatures at the transcript level. There is a poor correlation between transcript and protein abundances for some proteins in some tissues (Anderson,
10 N.L. and J. Seilhamer (1997) Electrophoresis 18:533-537), so proteome toxicant signatures may be useful in the analysis of compounds which do not significantly affect the transcript image, but which alter the proteomic profile. In addition, the analysis of transcripts in body fluids is difficult, due to rapid degradation of mRNA, so proteomic profiling may be more reliable and informative in such cases.

15 In another embodiment, the toxicity of a test compound is assessed by treating a biological sample containing proteins with the test compound. Proteins that are expressed in the treated biological sample are separated so that the amount of each protein can be quantified. The amount of each protein is compared to the amount of the corresponding protein in an untreated biological sample. A difference in the amount of protein between the two samples is indicative of a toxic
20 response to the test compound in the treated sample. Individual proteins are identified by sequencing the amino acid residues of the individual proteins and comparing these partial sequences to the polypeptides of the present invention.

In another embodiment, the toxicity of a test compound is assessed by treating a biological sample containing proteins with the test compound. Proteins from the biological sample are
25 incubated with antibodies specific to the polypeptides of the present invention. The amount of protein recognized by the antibodies is quantified. The amount of protein in the treated biological sample is compared with the amount in an untreated biological sample. A difference in the amount of protein between the two samples is indicative of a toxic response to the test compound in the treated sample.

30 Microarrays may be prepared, used, and analyzed using methods known in the art. (See, e.g., Brennan, T.M. et al. (1995) U.S. Patent No. 5,474,796; Schena, M. et al. (1996) Proc. Natl. Acad. Sci. USA 93:10614-10619; Baldeschweiler et al. (1995) PCT application WO95/251116; Shalon, D. et al. (1995) PCT application WO95/35505; Heller, R.A. et al. (1997) Proc. Natl. Acad. Sci. USA 94:2150-2155; and Heller, M.J. et al. (1997) U.S. Patent No. 5,605,662.) Various types of microarrays are
35 well known and thoroughly described in DNA Microarrays: A Practical Approach, M. Schena, ed.

(1999) Oxford University Press, London, hereby expressly incorporated by reference.

In another embodiment of the invention, nucleic acid sequences encoding SECP may be used to generate hybridization probes useful in mapping the naturally occurring genomic sequence. Either coding or noncoding sequences may be used, and in some instances, noncoding sequences may be preferable over coding sequences. For example, conservation of a coding sequence among members of a multi-gene family may potentially cause undesired cross hybridization during chromosomal mapping. The sequences may be mapped to a particular chromosome, to a specific region of a chromosome, or to artificial chromosome constructions, e.g., human artificial chromosomes (HACs), yeast artificial chromosomes (YACs), bacterial artificial chromosomes (BACs), bacterial P1 constructions, or single chromosome cDNA libraries. (See, e.g., Harrington, J.J. et al. (1997) *Nat. Genet.* 15:345-355; Price, C.M. (1993) *Blood Rev.* 7:127-134; and Trask, B.J. (1991) *Trends Genet.* 7:149-154.) Once mapped, the nucleic acid sequences of the invention may be used to develop genetic linkage maps, for example, which correlate the inheritance of a disease state with the inheritance of a particular chromosome region or restriction fragment length polymorphism (RFLP). (See, for example, Lander, E.S. and D. Botstein (1986) *Proc. Natl. Acad. Sci. USA* 83:7353-7357.)

Fluorescent in situ hybridization (FISH) may be correlated with other physical and genetic map data. (See, e.g., Heinz-Ulrich, et al. (1995) in Meyers, supra, pp. 965-968.) Examples of genetic map data can be found in various scientific journals or at the Online Mendelian Inheritance in Man (OMIM) World Wide Web site. Correlation between the location of the gene encoding SECP on a physical map and a specific disorder, or a predisposition to a specific disorder, may help define the region of DNA associated with that disorder and thus may further positional cloning efforts.

In situ hybridization of chromosomal preparations and physical mapping techniques, such as linkage analysis using established chromosomal markers, may be used for extending genetic maps. Often the placement of a gene on the chromosome of another mammalian species, such as mouse, may reveal associated markers even if the exact chromosomal locus is not known. This information is valuable to investigators searching for disease genes using positional cloning or other gene discovery techniques. Once the gene or genes responsible for a disease or syndrome have been crudely localized by genetic linkage to a particular genomic region, e.g., ataxia-telangiectasia to 11q22-23, any sequences mapping to that area may represent associated or regulatory genes for further investigation. (See, e.g., Gatti, R.A. et al. (1988) *Nature* 336:577-580.) The nucleotide sequence of the instant invention may also be used to detect differences in the chromosomal location due to translocation, inversion, etc., among normal, carrier, or affected individuals.

In another embodiment of the invention, SECP, its catalytic or immunogenic fragments, or oligopeptides thereof can be used for screening libraries of compounds in any of a variety of drug screening techniques. The fragment employed in such screening may be free in solution, affixed to a

solid support, borne on a cell surface, or located intracellularly. The formation of binding complexes between SECP and the agent being tested may be measured.

Another technique for drug screening provides for high throughput screening of compounds having suitable binding affinity to the protein of interest. (See, e.g., Geysen, et al. (1984) PCT application WO84/03564.) In this method, large numbers of different small test compounds are synthesized on a solid substrate. The test compounds are reacted with SECP, or fragments thereof, and washed. Bound SECP is then detected by methods well known in the art. Purified SECP can also be coated directly onto plates for use in the aforementioned drug screening techniques. Alternatively, non-neutralizing antibodies can be used to capture the peptide and immobilize it on a solid support.

In another embodiment, one may use competitive drug screening assays in which neutralizing antibodies capable of binding SECP specifically compete with a test compound for binding SECP. In this manner, antibodies can be used to detect the presence of any peptide which shares one or more antigenic determinants with SECP.

In additional embodiments, the nucleotide sequences which encode SECP may be used in any molecular biology techniques that have yet to be developed, provided the new techniques rely on properties of nucleotide sequences that are currently known, including, but not limited to, such properties as the triplet genetic code and specific base pair interactions.

Without further elaboration, it is believed that one skilled in the art can, using the preceding description, utilize the present invention to its fullest extent. The following embodiments are, therefore, to be construed as merely illustrative, and not limitative of the remainder of the disclosure in any way whatsoever.

The disclosures of all patents, applications, and publications mentioned above and below, including U.S. Ser. No. 60/236,869, U.S. Ser. No. 60/240,108, U.S. Ser. No. 60/239,812, U.S. Ser. No. 60/241,282, and U.S. Ser. No. 60/242,218, are hereby expressly incorporated by reference.

EXAMPLES

I. Construction of cDNA Libraries

Incyte cDNAs were derived from cDNA libraries described in the LIFESEQ GOLD database (Incyte Genomics, Palo Alto CA) and shown in Table 4, column 5. Some tissues were homogenized and lysed in guanidinium isothiocyanate, while others were homogenized and lysed in phenol or in a suitable mixture of denaturants, such as TRIZOL (Life Technologies), a monophasic solution of phenol and guanidine isothiocyanate. The resulting lysates were centrifuged over CsCl cushions or extracted with chloroform. RNA was precipitated from the lysates with either isopropanol or sodium acetate and ethanol, or by other routine methods.

Phenol extraction and precipitation of RNA were repeated as necessary to increase RNA purity. In some cases, RNA was treated with DNase. For most libraries, poly(A)+ RNA was isolated using oligo d(T)-coupled paramagnetic particles (Promega), OLIGOTEX latex particles (QIAGEN, Chatsworth CA), or an OLIGOTEX mRNA purification kit (QIAGEN). Alternatively, RNA was isolated directly from tissue lysates using other RNA isolation kits, e.g., the POLY(A)PURE mRNA purification kit (Ambion, Austin TX).

In some cases, Stratagene was provided with RNA and constructed the corresponding cDNA libraries. Otherwise, cDNA was synthesized and cDNA libraries were constructed with the UNIZAP vector system (Stratagene) or SUPERScript plasmid system (Life Technologies), using the recommended procedures or similar methods known in the art. (See, e.g., Ausubel, 1997, *supra*, units 5.1-6.6.) Reverse transcription was initiated using oligo d(T) or random primers. Synthetic oligonucleotide adapters were ligated to double stranded cDNA, and the cDNA was digested with the appropriate restriction enzyme or enzymes. For most libraries, the cDNA was size-selected (300-1000 bp) using SEPHACRYL S1000, SEPHAROSE CL2B, or SEPHAROSE CL4B column chromatography (Amersham Pharmacia Biotech) or preparative agarose gel electrophoresis. cDNAs were ligated into compatible restriction enzyme sites of the polylinker of a suitable plasmid, e.g., PBLUESCRIPT plasmid (Stratagene), PSORT1 plasmid (Life Technologies), PCDNA2.1 plasmid (Invitrogen, Carlsbad CA), PBK-CMV plasmid (Stratagene), PCR2-TOPOTA plasmid (Invitrogen), PCMV-ICIS plasmid (Stratagene), pIGEN (Incyte Genomics, Palo Alto CA), or pINCY (Incyte Genomics), or derivatives thereof. Recombinant plasmids were transformed into competent *E. coli* cells including XL1-Blue, XL1-BlueMRF, or SOLR from Stratagene or DH5 α , DH10B, or ElectroMAX DH10B from Life Technologies.

II. Isolation of cDNA Clones

Plasmids obtained as described in Example I were recovered from host cells by *in vivo* excision using the UNIZAP vector system (Stratagene) or by cell lysis. Plasmids were purified using at least one of the following: a Magic or WIZARD Minipreps DNA purification system (Promega); an AGTC Miniprep purification kit (Edge Biosystems, Gaithersburg MD); and QIAWELL 8 Plasmid, QIAWELL 8 Plus Plasmid, QIAWELL 8 Ultra Plasmid purification systems or the R.E.A.L. PREP 96 plasmid purification kit from QIAGEN. Following precipitation, plasmids were resuspended in 0.1 ml of distilled water and stored, with or without lyophilization, at 4°C.

Alternatively, plasmid DNA was amplified from host cell lysates using direct link PCR in a high-throughput format (Rao, V.B. (1994) Anal. Biochem. 216:1-14). Host cell lysis and thermal cycling steps were carried out in a single reaction mixture. Samples were processed and stored in 384-well plates, and the concentration of amplified plasmid DNA was quantified fluorometrically using PICOGREEN dye (Molecular Probes, Eugene OR) and a FLUOROSKAN II fluorescence

scanner (Labsystems Oy, Helsinki, Finland).

III. Sequencing and Analysis

Incyte cDNA recovered in plasmids as described in Example II were sequenced as follows. Sequencing reactions were processed using standard methods or high-throughput instrumentation such as the ABI CATALYST 800 (Applied Biosystems) thermal cycler or the PTC-200 thermal cycler (MJ Research) in conjunction with the HYDRA microdispenser (Robbins Scientific) or the MICROLAB 2200 (Hamilton) liquid transfer system. cDNA sequencing reactions were prepared using reagents provided by Amersham Pharmacia Biotech or supplied in ABI sequencing kits such as the ABI PRISM BIGDYE Terminator cycle sequencing ready reaction kit (Applied Biosystems). Electrophoretic separation of cDNA sequencing reactions and detection of labeled polynucleotides were carried out using the MEGABACE 1000 DNA sequencing system (Molecular Dynamics); the ABI PRISM 373 or 377 sequencing system (Applied Biosystems) in conjunction with standard ABI protocols and base calling software; or other sequence analysis systems known in the art. Reading frames within the cDNA sequences were identified using standard methods (reviewed in Ausubel, 1997, *supra*, unit 7.7). Some of the cDNA sequences were selected for extension using the techniques disclosed in Example VIII.

The polynucleotide sequences derived from Incyte cDNAs were validated by removing vector, linker, and poly(A) sequences and by masking ambiguous bases, using algorithms and programs based on BLAST, dynamic programming, and dinucleotide nearest neighbor analysis. The Incyte cDNA sequences or translations thereof were then queried against a selection of public databases such as the GenBank primate, rodent, mammalian, vertebrate, and eukaryote databases, and BLOCKS, PRINTS, DOMO, PRODOM, and hidden Markov model (HMM)-based protein family databases such as PFAM. (HMM is a probabilistic approach which analyzes consensus primary structures of gene families. See, for example, Eddy, S.R. (1996) *Curr. Opin. Struct. Biol.* 6:361-365.) The queries were performed using programs based on BLAST, FASTA, BLIMPS, and HMMER. The Incyte cDNA sequences were assembled to produce full length polynucleotide sequences. Alternatively, GenBank cDNAs, GenBank ESTs, stitched sequences, stretched sequences, or Genscan-predicted coding sequences (see Examples IV and V) were used to extend Incyte cDNA assemblages to full length. Assembly was performed using programs based on Phred, Phrap, and Consed, and cDNA assemblages were screened for open reading frames using programs based on GeneMark, BLAST, and FASTA. The full length polynucleotide sequences were translated to derive the corresponding full length polypeptide sequences. Alternatively, a polypeptide of the invention may begin at any of the methionine residues of the full length translated polypeptide. Full length polypeptide sequences were subsequently analyzed by querying against databases such as the GenBank protein databases (genpept), SwissProt, BLOCKS, PRINTS, DOMO, PRODOM, Prosite,

and hidden Markov model (HMM)-based protein family databases such as PFAM. Full length polynucleotide sequences are also analyzed using MACDNASIS PRO software (Hitachi Software Engineering, South San Francisco CA) and LASERGENE software (DNASTAR). Polynucleotide and polypeptide sequence alignments are generated using default parameters specified by the CLUSTAL algorithm as incorporated into the MEGALIGN multisequence alignment program (DNASTAR), which also calculates the percent identity between aligned sequences.

Table 7 summarizes the tools, programs, and algorithms used for the analysis and assembly of Incyte cDNA and full length sequences and provides applicable descriptions, references, and threshold parameters. The first column of Table 7 shows the tools, programs, and algorithms used, the second column provides brief descriptions thereof, the third column presents appropriate references, all of which are incorporated by reference herein in their entirety, and the fourth column presents, where applicable, the scores, probability values, and other parameters used to evaluate the strength of a match between two sequences (the higher the score or the lower the probability value, the greater the identity between two sequences).

The programs described above for the assembly and analysis of full length polynucleotide and polypeptide sequences were also used to identify polynucleotide sequence fragments from SEQ ID NO:68-136. Fragments from about 20 to about 4000 nucleotides which are useful in hybridization and amplification technologies are described in Table 4, column 2.

IV. Identification and Editing of Coding Sequences from Genomic DNA

Putative secreted proteins were initially identified by running the Genscan gene identification program against public genomic sequence databases (e.g., gbpri and gbhtg). Genscan is a general-purpose gene identification program which analyzes genomic DNA sequences from a variety of organisms (See Burge, C. and S. Karlin (1997) J. Mol. Biol. 268:78-94, and Burge, C. and S. Karlin (1998) Curr. Opin. Struct. Biol. 8:346-354). The program concatenates predicted exons to form an assembled cDNA sequence extending from a methionine to a stop codon. The output of Genscan is a FASTA database of polynucleotide and polypeptide sequences. The maximum range of sequence for Genscan to analyze at once was set to 30 kb. To determine which of these Genscan predicted cDNA sequences encode secreted proteins, the encoded polypeptides were analyzed by querying against PFAM models for secreted proteins. Potential secreted proteins were also identified by homology to Incyte cDNA sequences that had been annotated as secreted proteins. These selected Genscan-predicted sequences were then compared by BLAST analysis to the genpept and gbpri public databases. Where necessary, the Genscan-predicted sequences were then edited by comparison to the top BLAST hit from genpept to correct errors in the sequence predicted by Genscan, such as extra or omitted exons. BLAST analysis was also used to find any Incyte cDNA or public cDNA coverage of the Genscan-predicted sequences, thus providing evidence for transcription. When Incyte cDNA

coverage was available, this information was used to correct or confirm the Genscan predicted sequence. Full length polynucleotide sequences were obtained by assembling Genscan-predicted coding sequences with Incyte cDNA sequences and/or public cDNA sequences using the assembly process described in Example III. Alternatively, full length polynucleotide sequences were derived
5 entirely from edited or unedited Genscan-predicted coding sequences.

V. Assembly of Genomic Sequence Data with cDNA Sequence Data

"Stitched" Sequences

Partial cDNA sequences were extended with exons predicted by the Genscan gene identification program described in Example IV. Partial cDNAs assembled as described in Example
10 III were mapped to genomic DNA and parsed into clusters containing related cDNAs and Genscan exon predictions from one or more genomic sequences. Each cluster was analyzed using an algorithm based on graph theory and dynamic programming to integrate cDNA and genomic information, generating possible splice variants that were subsequently confirmed, edited, or extended to create a full length sequence. Sequence intervals in which the entire length of the interval was present on
15 more than one sequence in the cluster were identified, and intervals thus identified were considered to be equivalent by transitivity. For example, if an interval was present on a cDNA and two genomic sequences, then all three intervals were considered to be equivalent. This process allows unrelated but consecutive genomic sequences to be brought together, bridged by cDNA sequence. Intervals thus identified were then "stitched" together by the stitching algorithm in the order that they appear
20 along their parent sequences to generate the longest possible sequence, as well as sequence variants. Linkages between intervals which proceed along one type of parent sequence (cDNA to cDNA or genomic sequence to genomic sequence) were given preference over linkages which change parent type (cDNA to genomic sequence). The resultant stitched sequences were translated and compared by BLAST analysis to the genpept and gbpr public databases. Incorrect exons predicted by Genscan
25 were corrected by comparison to the top BLAST hit from genpept. Sequences were further extended with additional cDNA sequences, or by inspection of genomic DNA, when necessary.

"Stretched" Sequences

Partial DNA sequences were extended to full length with an algorithm based on BLAST analysis. First, partial cDNAs assembled as described in Example III were queried against public
30 databases such as the GenBank primate, rodent, mammalian, vertebrate, and eukaryote databases using the BLAST program. The nearest GenBank protein homolog was then compared by BLAST analysis to either Incyte cDNA sequences or GenScan exon predicted sequences described in Example IV. A chimeric protein was generated by using the resultant high-scoring segment pairs (HSPs) to map the translated sequences onto the GenBank protein homolog. Insertions or deletions
35 may occur in the chimeric protein with respect to the original GenBank protein homolog. The

GenBank protein homolog, the chimeric protein, or both were used as probes to search for homologous genomic sequences from the public human genome databases. Partial DNA sequences were therefore "stretched" or extended by the addition of homologous genomic sequences. The resultant stretched sequences were examined to determine whether it contained a complete gene.

5 VI. Chromosomal Mapping of SECP Encoding Polynucleotides

The sequences which were used to assemble SEQ ID NO:68-134 were compared with sequences from the Incyte LIFESEQ database and public domain databases using BLAST and other implementations of the Smith-Waterman algorithm. Sequences from these databases that matched SEQ ID NO:68-134 were assembled into clusters of contiguous and overlapping sequences using
10 assembly algorithms such as Phrap (Table 7). Radiation hybrid and genetic mapping data available from public resources such as the Stanford Human Genome Center (SHGC), Whitehead Institute for Genome Research (WIGR), and Généthon were used to determine if any of the clustered sequences had been previously mapped. Inclusion of a mapped sequence in a cluster resulted in the assignment of all sequences of that cluster, including its particular SEQ ID NO:, to that map location.

15 Map locations are represented by ranges, or intervals, of human chromosomes. The map position of an interval, in centiMorgans, is measured relative to the terminus of the chromosome's p-arm. (The centiMorgan (cM) is a unit of measurement based on recombination frequencies between chromosomal markers. On average, 1 cM is roughly equivalent to 1 megabase (Mb) of DNA in humans, although this can vary widely due to hot and cold spots of recombination.) The cM
20 distances are based on genetic markers mapped by Généthon which provide boundaries for radiation hybrid markers whose sequences were included in each of the clusters. Human genome maps and other resources available to the public, such as the NCBI "GeneMap'99" World Wide Web site (<http://www.ncbi.nlm.nih.gov/genemap/>), can be employed to determine if previously identified disease genes map within or in proximity to the intervals indicated above.

25 In this manner, SEQ ID NO:70 was mapped to chromosome 5 within the interval from 79.2 to 92.3 centiMorgans. SEQ ID NO:98 was mapped to chromosome 4 within the interval from 145.3 to 146.4 centiMorgans.

VII. Analysis of Polynucleotide Expression

Northern analysis is a laboratory technique used to detect the presence of a transcript of a
30 gene and involves the hybridization of a labeled nucleotide sequence to a membrane on which RNAs from a particular cell type or tissue have been bound. (See, e.g., Sambrook, *supra*, ch. 7; Ausubel (1995) *supra*, ch. 4 and 16.)

Analogous computer techniques applying BLAST were used to search for identical or related molecules in cDNA databases such as GenBank or LIFESEQ (Incyte Genomics). This analysis is
35 much faster than multiple membrane-based hybridizations. In addition, the sensitivity of the

computer search can be modified to determine whether any particular match is categorized as exact or similar. The basis of the search is the product score, which is defined as:

$$\frac{\text{BLAST Score} \times \text{Percent Identity}}{5 \times \text{minimum \{length(Seq. 1), length(Seq. 2)\}}}$$

The product score takes into account both the degree of similarity between two sequences and the length of the sequence match. The product score is a normalized value between 0 and 100, and is calculated as follows: the BLAST score is multiplied by the percent nucleotide identity and the product is divided by (5 times the length of the shorter of the two sequences). The BLAST score is calculated by assigning a score of +5 for every base that matches in a high-scoring segment pair (HSP), and -4 for every mismatch. Two sequences may share more than one HSP (separated by gaps). If there is more than one HSP, then the pair with the highest BLAST score is used to calculate the product score. The product score represents a balance between fractional overlap and quality in a BLAST alignment. For example, a product score of 100 is produced only for 100% identity over the entire length of the shorter of the two sequences being compared. A product score of 70 is produced either by 100% identity and 70% overlap at one end, or by 88% identity and 100% overlap at the other. A product score of 50 is produced either by 100% identity and 50% overlap at one end, or 79% identity and 100% overlap.

Alternatively, polynucleotide sequences encoding SECP are analyzed with respect to the tissue sources from which they were derived. For example, some full length sequences are assembled, at least in part, with overlapping Incyte cDNA sequences (see Example III). Each cDNA sequence is derived from a cDNA library constructed from a human tissue. Each human tissue is classified into one of the following organ/tissue categories: cardiovascular system; connective tissue; digestive system; embryonic structures; endocrine system; exocrine glands; genitalia, female; genitalia, male; germ cells; hemic and immune system; liver; musculoskeletal system; nervous system; pancreas; respiratory system; sense organs; skin; stomatognathic system; unclassified/mixed; or urinary tract. The number of libraries in each category is counted and divided by the total number of libraries across all categories. Similarly, each human tissue is classified into one of the following disease/condition categories: cancer, cell line, developmental, inflammation, neurological, trauma, cardiovascular, pooled, and other, and the number of libraries in each category is counted and divided by the total number of libraries across all categories. The resulting percentages reflect the tissue- and disease-specific expression of cDNA encoding SECP. cDNA sequences and cDNA library/tissue information are found in the LIFESEQ GOLD database (Incyte Genomics, Palo Alto CA).

VIII. Extension of SECP Encoding Polynucleotides

Full length polynucleotide sequences were also produced by extension of an appropriate fragment of the full length molecule using oligonucleotide primers designed from this fragment. One primer was synthesized to initiate 5' extension of the known fragment, and the other primer was synthesized to initiate 3' extension of the known fragment. The initial primers were designed using
5 OLIGO 4.06 software (National Biosciences), or another appropriate program, to be about 22 to 30 nucleotides in length, to have a GC content of about 50% or more, and to anneal to the target sequence at temperatures of about 68°C to about 72°C. Any stretch of nucleotides which would result in hairpin structures and primer-primer dimerizations was avoided.

Selected human cDNA libraries were used to extend the sequence. If more than one
10 extension was necessary or desired, additional or nested sets of primers were designed.

High fidelity amplification was obtained by PCR using methods well known in the art. PCR was performed in 96-well plates using the PTC-200 thermal cycler (MJ Research, Inc.). The reaction mix contained DNA template, 200 nmol of each primer, reaction buffer containing Mg^{2+} , $(NH_4)_2SO_4$, and 2-mercaptoethanol, Taq DNA polymerase (Amersham Pharmacia Biotech), ELONGASE enzyme
15 (Life Technologies), and Pfu DNA polymerase (Stratagene), with the following parameters for primer pair PCI A and PCI B: Step 1: 94°C, 3 min; Step 2: 94°C, 15 sec; Step 3: 60°C, 1 min; Step 4: 68°C, 2 min; Step 5: Steps 2, 3, and 4 repeated 20 times; Step 6: 68°C, 5 min; Step 7: storage at 4°C. In the alternative, the parameters for primer pair T7 and SK+ were as follows: Step 1: 94°C, 3 min; Step 2: 94°C, 15 sec; Step 3: 57°C, 1 min; Step 4: 68°C, 2 min; Step 5: Steps 2, 3, and 4 repeated 20 times;
20 Step 6: 68°C, 5 min; Step 7: storage at 4°C.

The concentration of DNA in each well was determined by dispensing 100 μ l PICOGREEN quantitation reagent (0.25% (v/v) PICOGREEN; Molecular Probes, Eugene OR) dissolved in 1X TE and 0.5 μ l of undiluted PCR product into each well of an opaque fluorimeter plate (Corning Costar, Acton MA), allowing the DNA to bind to the reagent. The plate was scanned in a Fluoroskan II
25 (Labsystems Oy, Helsinki, Finland) to measure the fluorescence of the sample and to quantify the concentration of DNA. A 5 μ l to 10 μ l aliquot of the reaction mixture was analyzed by electrophoresis on a 1 % agarose gel to determine which reactions were successful in extending the sequence.

The extended nucleotides were desalted and concentrated, transferred to 384-well plates,
30 digested with CviJI cholera virus endonuclease (Molecular Biology Research, Madison WI), and sonicated or sheared prior to religation into pUC 18 vector (Amersham Pharmacia Biotech). For shotgun sequencing, the digested nucleotides were separated on low concentration (0.6 to 0.8%) agarose gels, fragments were excised, and agar digested with Agar ACE (Promega). Extended clones were religated using T4 ligase (New England Biolabs, Beverly MA) into pUC 18 vector (Amersham
35 Pharmacia Biotech), treated with Pfu DNA polymerase (Stratagene) to fill-in restriction site

overhangs, and transfected into competent *E. coli* cells. Transformed cells were selected on antibiotic-containing media, and individual colonies were picked and cultured overnight at 37°C in 384-well plates in LB/2x carb liquid media.

The cells were lysed, and DNA was amplified by PCR using Taq DNA polymerase (Amersham Pharmacia Biotech) and Pfu DNA polymerase (Stratagene) with the following parameters: Step 1: 94°C, 3 min; Step 2: 94°C, 15 sec; Step 3: 60°C, 1 min; Step 4: 72°C, 2 min; Step 5: steps 2, 3, and 4 repeated 29 times; Step 6: 72°C, 5 min; Step 7: storage at 4°C. DNA was quantified by PICOGREEN reagent (Molecular Probes) as described above. Samples with low DNA recoveries were reamplified using the same conditions as described above. Samples were diluted with 20% dimethylsulfoxide (1:2, v/v), and sequenced using DYENAMIC energy transfer sequencing primers and the DYENAMIC DIRECT kit (Amersham Pharmacia Biotech) or the ABI PRISM BIGDYE Terminator cycle sequencing ready reaction kit (Applied Biosystems).

In like manner, full length polynucleotide sequences are verified using the above procedure or are used to obtain 5' regulatory sequences using the above procedure along with oligonucleotides designed for such extension, and an appropriate genomic library.

IX. Labeling and Use of Individual Hybridization Probes

Hybridization probes derived from SEQ ID NO:68-136 are employed to screen cDNAs, genomic DNAs, or mRNAs. Although the labeling of oligonucleotides, consisting of about 20 base pairs, is specifically described, essentially the same procedure is used with larger nucleotide fragments. Oligonucleotides are designed using state-of-the-art software such as OLIGO 4.06 software (National Biosciences) and labeled by combining 50 pmol of each oligomer, 250 μ Ci of [γ -³²P] adenosine triphosphate (Amersham Pharmacia Biotech), and T4 polynucleotide kinase (DuPont NEN, Boston MA). The labeled oligonucleotides are substantially purified using a SEPHADEX G-25 superfine size exclusion dextran bead column (Amersham Pharmacia Biotech). An aliquot containing 10⁷ counts per minute of the labeled probe is used in a typical membrane-based hybridization analysis of human genomic DNA digested with one of the following endonucleases: Ase I, Bgl II, Eco RI, Pst I, Xba I, or Pvu II (DuPont NEN).

The DNA from each digest is fractionated on a 0.7% agarose gel and transferred to nylon membranes (Nytran Plus, Schleicher & Schuell, Durham NH). Hybridization is carried out for 16 hours at 40°C. To remove nonspecific signals, blots are sequentially washed at room temperature under conditions of up to, for example, 0.1 x saline sodium citrate and 0.5% sodium dodecyl sulfate. Hybridization patterns are visualized using autoradiography or an alternative imaging means and compared.

X. Microarrays

The linkage or synthesis of array elements upon a microarray can be achieved utilizing

photolithography, piezoelectric printing (ink-jet printing, See, e.g., Baldeschweiler, supra), mechanical microspotting technologies, and derivatives thereof. The substrate in each of the aforementioned technologies should be uniform and solid with a non-porous surface (Skena (1999), supra). Suggested substrates include silicon, silica, glass slides, glass chips, and silicon wafers.

- 5 Alternatively, a procedure analogous to a dot or slot blot may also be used to arrange and link elements to the surface of a substrate using thermal, UV, chemical, or mechanical bonding procedures. A typical array may be produced using available methods and machines well known to those of ordinary skill in the art and may contain any appropriate number of elements. (See, e.g., Skena, M. et al. (1995) *Science* 270:467-470; Shalon, D. et al. (1996) *Genome Res.* 6:639-645; Marshall, A. and J. Hodgson (1998) *Nat. Biotechnol.* 16:27-31.)

- 10 Full length cDNAs, Expressed Sequence Tags (ESTs), or fragments or oligomers thereof may comprise the elements of the microarray. Fragments or oligomers suitable for hybridization can be selected using software well known in the art such as LASERGENE software (DNASTAR). The array elements are hybridized with polynucleotides in a biological sample. The polynucleotides in the biological sample are conjugated to a fluorescent label or other molecular tag for ease of detection.
- 15 After hybridization, nonhybridized nucleotides from the biological sample are removed, and a fluorescence scanner is used to detect hybridization at each array element. Alternatively, laser desorption and mass spectrometry may be used for detection of hybridization. The degree of complementarity and the relative abundance of each polynucleotide which hybridizes to an element on the microarray may be assessed. In one embodiment, microarray preparation and usage is described in detail below.

Tissue or Cell Sample Preparation

- Total RNA is isolated from tissue samples using the guanidinium thiocyanate method and poly(A)⁺ RNA is purified using the oligo-(dT) cellulose method. Each poly(A)⁺ RNA sample is reverse transcribed using MMLV reverse-transcriptase, 0.05 pg/ μ l oligo-(dT) primer (21mer), 1X first strand buffer, 0.03 units/ μ l RNase inhibitor, 500 μ M dATP, 500 μ M dGTP, 500 μ M dTTP, 40 μ M dCTP, 40 μ M dCTP-Cy3 (BDS) or dCTP-Cy5 (Amersham Pharmacia Biotech). The reverse transcription reaction is performed in a 25 ml volume containing 200 ng poly(A)⁺ RNA with GEMBRIGHT kits (Incyte). Specific control poly(A)⁺ RNAs are synthesized by in vitro transcription from non-coding yeast genomic DNA. After incubation at 37°C for 2 hr, each reaction sample (one with Cy3 and another with Cy5 labeling) is treated with 2.5 ml of 0.5M sodium hydroxide and incubated for 20 minutes at 85°C to stop the reaction and degrade the RNA. Samples are purified using two successive CHROMA SPIN 30 gel filtration spin columns (CLONTECH Laboratories, Inc. (CLONTECH), Palo Alto CA) and after combining, both reaction samples are ethanol precipitated using 1 ml of glycogen (1 mg/ml), 60 ml sodium acetate, and 300 ml of 100% ethanol. The sample is

then dried to completion using a SpeedVAC (Savant Instruments Inc., Holbrook NY) and resuspended in 14 μ l 5X SSC/0.2% SDS.

Microarray Preparation

Sequences of the present invention are used to generate array elements. Each array element
5 is amplified from bacterial cells containing vectors with cloned cDNA inserts. PCR amplification uses primers complementary to the vector sequences flanking the cDNA insert. Array elements are amplified in thirty cycles of PCR from an initial quantity of 1-2 ng to a final quantity greater than 5 μ g. Amplified array elements are then purified using SEPHACRYL-400 (Amersham Pharmacia Biotech).

10 Purified array elements are immobilized on polymer-coated glass slides. Glass microscope slides (Corning) are cleaned by ultrasound in 0.1% SDS and acetone, with extensive distilled water washes between and after treatments. Glass slides are etched in 4% hydrofluoric acid (VWR Scientific Products Corporation (VWR), West Chester PA), washed extensively in distilled water, and coated with 0.05% aminopropyl silane (Sigma) in 95% ethanol. Coated slides are cured in a
15 110°C oven.

Array elements are applied to the coated glass substrate using a procedure described in U.S. Patent No. 5,807,522, incorporated herein by reference. 1 μ l of the array element DNA, at an average concentration of 100 ng/ μ l, is loaded into the open capillary printing element by a high-speed robotic apparatus. The apparatus then deposits about 5 nl of array element sample per slide.

20 Microarrays are UV-crosslinked using a STRATALINKER UV-crosslinker (Stratagene). Microarrays are washed at room temperature once in 0.2% SDS and three times in distilled water. Non-specific binding sites are blocked by incubation of microarrays in 0.2% casein in phosphate buffered saline (PBS) (Tropix, Inc., Bedford MA) for 30 minutes at 60°C followed by washes in 0.2% SDS and distilled water as before.

25 Hybridization

Hybridization reactions contain 9 μ l of sample mixture consisting of 0.2 μ g each of Cy3 and Cy5 labeled cDNA synthesis products in 5X SSC, 0.2% SDS hybridization buffer. The sample mixture is heated to 65°C for 5 minutes and is aliquoted onto the microarray surface and covered with an 1.8 cm² coverslip. The arrays are transferred to a waterproof chamber having a cavity just
30 slightly larger than a microscope slide. The chamber is kept at 100% humidity internally by the addition of 140 μ l of 5X SSC in a corner of the chamber. The chamber containing the arrays is incubated for about 6.5 hours at 60°C. The arrays are washed for 10 min at 45°C in a first wash buffer (1X SSC, 0.1% SDS), three times for 10 minutes each at 45°C in a second wash buffer (0.1X SSC), and dried.

35 Detection

Reporter-labeled hybridization complexes are detected with a microscope equipped with an Innova 70 mixed gas 10 W laser (Coherent, Inc., Santa Clara CA) capable of generating spectral lines at 488 nm for excitation of Cy3 and at 632 nm for excitation of Cy5. The excitation laser light is focused on the array using a 20X microscope objective (Nikon, Inc., Melville NY). The slide
5 containing the array is placed on a computer-controlled X-Y stage on the microscope and raster-scanned past the objective. The 1.8 cm x 1.8 cm array used in the present example is scanned with a resolution of 20 micrometers.

In two separate scans, a mixed gas multiline laser excites the two fluorophores sequentially. Emitted light is split, based on wavelength, into two photomultiplier tube detectors (PMT R1477,
10 Hamamatsu Photonics Systems, Bridgewater NJ) corresponding to the two fluorophores. Appropriate filters positioned between the array and the photomultiplier tubes are used to filter the signals. The emission maxima of the fluorophores used are 565 nm for Cy3 and 650 nm for Cy5. Each array is typically scanned twice, one scan per fluorophore using the appropriate filters at the laser source, although the apparatus is capable of recording the spectra from both fluorophores simultaneously.

15 The sensitivity of the scans is typically calibrated using the signal intensity generated by a cDNA control species added to the sample mixture at a known concentration. A specific location on the array contains a complementary DNA sequence, allowing the intensity of the signal at that location to be correlated with a weight ratio of hybridizing species of 1:100,000. When two samples from different sources (e.g., representing test and control cells), each labeled with a different
20 fluorophore, are hybridized to a single array for the purpose of identifying genes that are differentially expressed, the calibration is done by labeling samples of the calibrating cDNA with the two fluorophores and adding identical amounts of each to the hybridization mixture.

The output of the photomultiplier tube is digitized using a 12-bit RTI-835H analog-to-digital (A/D) conversion board (Analog Devices, Inc., Norwood MA) installed in an IBM-compatible PC
25 computer. The digitized data are displayed as an image where the signal intensity is mapped using a linear 20-color transformation to a pseudocolor scale ranging from blue (low signal) to red (high signal). The data is also analyzed quantitatively. Where two different fluorophores are excited and measured simultaneously, the data are first corrected for optical crosstalk (due to overlapping emission spectra) between the fluorophores using each fluorophore's emission spectrum.

30 A grid is superimposed over the fluorescence signal image such that the signal from each spot is centered in each element of the grid. The fluorescence signal within each element is then integrated to obtain a numerical value corresponding to the average intensity of the signal. The software used for signal analysis is the GEMTOOLS gene expression analysis program (Incyte).

XI. Complementary Polynucleotides

Sequences complementary to the SECP-encoding sequences, or any parts thereof, are used to detect, decrease, or inhibit expression of naturally occurring SECP. For example, SEQ ID NO:135 and SEQ ID NO:136 are complementary polynucleotides to SEQ ID NO:114 and SEQ ID NO:116, respectively. Although use of oligonucleotides comprising from about 15 to 30 base pairs is described, essentially the same procedure is used with smaller or with larger sequence fragments. Appropriate oligonucleotides are designed using OLIGO 4.06 software (National Biosciences) and the coding sequence of SECP. To inhibit transcription, a complementary oligonucleotide is designed from the most unique 5' sequence and used to prevent promoter binding to the coding sequence. To inhibit translation, a complementary oligonucleotide is designed to prevent ribosomal binding to the SECP-encoding transcript.

XII. Expression of SECP

Expression and purification of SECP is achieved using bacterial or virus-based expression systems. For expression of SECP in bacteria, cDNA is subcloned into an appropriate vector containing an antibiotic resistance gene and an inducible promoter that directs high levels of cDNA transcription. Examples of such promoters include, but are not limited to, the *trp-lac (tac)* hybrid promoter and the T5 or T7 bacteriophage promoter in conjunction with the *lac* operator regulatory element. Recombinant vectors are transformed into suitable bacterial hosts, e.g., BL21(DE3). Antibiotic resistant bacteria express SECP upon induction with isopropyl beta-D-thiogalactopyranoside (IPTG). Expression of SECP in eukaryotic cells is achieved by infecting insect or mammalian cell lines with recombinant Autographica californica nuclear polyhedrosis virus (AcMNPV), commonly known as baculovirus. The nonessential polyhedrin gene of baculovirus is replaced with cDNA encoding SECP by either homologous recombination or bacterial-mediated transposition involving transfer plasmid intermediates. Viral infectivity is maintained and the strong polyhedrin promoter drives high levels of cDNA transcription. Recombinant baculovirus is used to infect Spodoptera frugiperda (Sf9) insect cells in most cases, or human hepatocytes, in some cases. Infection of the latter requires additional genetic modifications to baculovirus. (See Engelhard, E.K. et al. (1994) Proc. Natl. Acad. Sci. USA 91:3224-3227; Sandig, V. et al. (1996) Hum. Gene Ther. 7:1937-1945.)

In most expression systems, SECP is synthesized as a fusion protein with, e.g., glutathione S-transferase (GST) or a peptide epitope tag, such as FLAG or 6-His, permitting rapid, single-step, affinity-based purification of recombinant fusion protein from crude cell lysates. GST, a 26-kilodalton enzyme from Schistosoma japonicum, enables the purification of fusion proteins on immobilized glutathione under conditions that maintain protein activity and antigenicity (Amersham Pharmacia Biotech). Following purification, the GST moiety can be proteolytically cleaved from

SECP at specifically engineered sites. FLAG, an 8-amino acid peptide, enables immunoaffinity purification using commercially available monoclonal and polyclonal anti-FLAG antibodies (Eastman Kodak). 6-His, a stretch of six consecutive histidine residues, enables purification on metal-chelate resins (QIAGEN). Methods for protein expression and purification are discussed in Ausubel (1995, 5 supra, ch. 10 and 16). Purified SECP obtained by these methods can be used directly in the assays shown in Examples XVI, XVII, and XVIII, where applicable.

XIII. Functional Assays

SECP function is assessed by expressing the sequences encoding SECP at physiologically elevated levels in mammalian cell culture systems. cDNA is subcloned into a mammalian expression 10 vector containing a strong promoter that drives high levels of cDNA expression. Vectors of choice include PCMV SPORT (Life Technologies) and PCR3.1 (Invitrogen, Carlsbad CA), both of which contain the cytomegalovirus promoter. 5-10 μ g of recombinant vector are transiently transfected into a human cell line, for example, an endothelial or hematopoietic cell line, using either liposome formulations or electroporation. 1-2 μ g of an additional plasmid containing sequences encoding a 15 marker protein are co-transfected. Expression of a marker protein provides a means to distinguish transfected cells from nontransfected cells and is a reliable predictor of cDNA expression from the recombinant vector. Marker proteins of choice include, e.g., Green Fluorescent Protein (GFP; Clontech), CD64, or a CD64-GFP fusion protein. Flow cytometry (FCM), an automated, laser optics-based technique, is used to identify transfected cells expressing GFP or CD64-GFP and to evaluate 20 the apoptotic state of the cells and other cellular properties. FCM detects and quantifies the uptake of fluorescent molecules that diagnose events preceding or coincident with cell death. These events include changes in nuclear DNA content as measured by staining of DNA with propidium iodide; changes in cell size and granularity as measured by forward light scatter and 90 degree side light scatter; down-regulation of DNA synthesis as measured by decrease in bromodeoxyuridine uptake; 25 alterations in expression of cell surface and intracellular proteins as measured by reactivity with specific antibodies; and alterations in plasma membrane composition as measured by the binding of fluorescein-conjugated Annexin V protein to the cell surface. Methods in flow cytometry are discussed in Ormerod, M.G. (1994) Flow Cytometry, Oxford, New York NY.

The influence of SECP on gene expression can be assessed using highly purified populations 30 of cells transfected with sequences encoding SECP and either CD64 or CD64-GFP. CD64 and CD64-GFP are expressed on the surface of transfected cells and bind to conserved regions of human immunoglobulin G (IgG). Transfected cells are efficiently separated from nontransfected cells using magnetic beads coated with either human IgG or antibody against CD64 (DYNAL, Lake Success NY). mRNA can be purified from the cells using methods well known by those of skill in the art. 35 Expression of mRNA encoding SECP and other genes of interest can be analyzed by northern

analysis or microarray techniques.

XIV. Production of SECP Specific Antibodies

SECP substantially purified using polyacrylamide gel electrophoresis (PAGE; see, e.g., Harrington, M.G. (1990) *Methods Enzymol.* 182:488-495), or other purification techniques, is used to immunize rabbits and to produce antibodies using standard protocols.

Alternatively, the SECP amino acid sequence is analyzed using LASERGENE software (DNASTAR) to determine regions of high immunogenicity, and a corresponding oligopeptide is synthesized and used to raise antibodies by means known to those of skill in the art. Methods for selection of appropriate epitopes, such as those near the C-terminus or in hydrophilic regions are well described in the art. (See, e.g., Ausubel, 1995, *supra*, ch. 11.)

Typically, oligopeptides of about 15 residues in length are synthesized using an ABI 431A peptide synthesizer (Applied Biosystems) using Fmoc chemistry and coupled to KLH (Sigma-Aldrich, St. Louis MO) by reaction with N-maleimidobenzoyl-N-hydroxysuccinimide ester (MBS) to increase immunogenicity. (See, e.g., Ausubel, 1995, *supra*.) Rabbits are immunized with the oligopeptide-KLH complex in complete Freund's adjuvant. Resulting antisera are tested for antipeptide and anti-SECP activity by, for example, binding the peptide or SECP to a substrate, blocking with 1% BSA, reacting with rabbit antisera, washing, and reacting with radio-iodinated goat anti-rabbit IgG.

XV. Purification of Naturally Occurring SECP Using Specific Antibodies

Naturally occurring or recombinant SECP is substantially purified by immunoaffinity chromatography using antibodies specific for SECP. An immunoaffinity column is constructed by covalently coupling anti-SECP antibody to an activated chromatographic resin, such as CNBr-activated SEPHAROSE (Amersham Pharmacia Biotech). After the coupling, the resin is blocked and washed according to the manufacturer's instructions.

Media containing SECP are passed over the immunoaffinity column, and the column is washed under conditions that allow the preferential absorbance of SECP (e.g., high ionic strength buffers in the presence of detergent). The column is eluted under conditions that disrupt antibody/SECP binding (e.g., a buffer of pH 2 to pH 3, or a high concentration of a chaotrope, such as urea or thiocyanate ion), and SECP is collected.

XVI. Identification of Molecules Which Interact with SECP

SECP, or biologically active fragments thereof, are labeled with ^{125}I Bolton-Hunter reagent. (See, e.g., Bolton A.E. and W.M. Hunter (1973) *Biochem. J.* 133:529-539.) Candidate molecules previously arrayed in the wells of a multi-well plate are incubated with the labeled SECP, washed, and any wells with labeled SECP complex are assayed. Data obtained using different concentrations of SECP are used to calculate values for the number, affinity, and association of SECP with the

candidate molecules.

Alternatively, molecules interacting with SECP are analyzed using the yeast two-hybrid system as described in Fields, S. and O. Song (1989) *Nature* 340:245-246, or using commercially available kits based on the two-hybrid system, such as the MATCHMAKER system (Clontech).

- 5 SECP may also be used in the PATHCALLING process (CuraGen Corp., New Haven CT) which employs the yeast two-hybrid system in a high-throughput manner to determine all interactions between the proteins encoded by two large libraries of genes (Nandabalan, K. et al. (2000) U.S. Patent No. 6,057,101).

XVII. Demonstration of SECP Activity

- 10 An assay for growth stimulating or inhibiting activity of SECP measures the amount of DNA synthesis in Swiss mouse 3T3 cells (McKay, I. and Leigh, I., eds. (1993) Growth Factors: A Practical Approach, Oxford University Press, New York, NY). In this assay, varying amounts of SECP are added to quiescent 3T3 cultured cells in the presence of [³H]thymidine, a radioactive DNA precursor. SECP for this assay can be obtained by recombinant means or from biochemical preparations.
- 15 Incorporation of [³H]thymidine into acid-precipitable DNA is measured over an appropriate time interval, and the amount incorporated is directly proportional to the amount of newly synthesized DNA. A linear dose-response curve over at least a hundred-fold SECP concentration range is indicative of growth modulating activity. One unit of activity per milliliter is defined as the concentration of SECP producing a 50% response level, where 100% represents maximal
- 20 incorporation of [³H]thymidine into acid-precipitable DNA.

- Alternatively, an assay for SECP activity measures the stimulation or inhibition of neurotransmission in cultured cells. Cultured CHO fibroblasts are exposed to SECP. Following endocytic uptake of SECP, the cells are washed with fresh culture medium, and a whole cell voltage-clamped Xenopus myocyte is manipulated into contact with one of the fibroblasts in SECP-free
- 25 medium. Membrane currents are recorded from the myocyte. Increased or decreased current relative to control values are indicative of neuromodulatory effects of SECP (Morimoto, T. et al. (1995) *Neuron* 15:689-696).

- Alternatively, an assay for SECP activity measures the amount of SECP in secretory, membrane-bound organelles. Transfected cells as described above are harvested and lysed. The
- 30 lysate is fractionated using methods known to those of skill in the art, for example, sucrose gradient ultracentrifugation. Such methods allow the isolation of subcellular components such as the Golgi apparatus, ER, small membrane-bound vesicles, and other secretory organelles.
- Immunoprecipitations from fractionated and total cell lysates are performed using SECP-specific antibodies, and immunoprecipitated samples are analyzed using SDS-PAGE and immunoblotting
- 35 techniques. The concentration of SECP in secretory organelles relative to SECP in total cell lysate is

proportional to the amount of SECP in transit through the secretory pathway.

Alternatively, AMP binding activity is measured by combining SECP with ^{32}P -labeled AMP. The reaction is incubated at 37°C and terminated by addition of trichloroacetic acid. The acid extract is neutralized and subjected to gel electrophoresis to remove unbound label. The radioactivity retained in the gel is proportional to SECP activity.

Alternatively, SECP activity for SEQ ID NO:67, for example, can be measured as protease inhibitory activity. Trypsin (100 units) is incubated at ambient temperature in a quartz cuvette in pH 7.6 assay buffer containing 63 mM sodium phosphate, 0.23 mM N α -benzoyl-L-arginine ethyl ester, 0.06 mM hydrochloric acid, with or without SECP. Immediately after mixing by inversion, the increase in $A_{253\text{ nm}}$ is recorded for approximately 5 minutes and the enzyme activity is calculated (Bergmeyer, H.U. et al. (1974) Meth. Enzym. Anal. 1:515-516). SECP activity is proportional to its effect on the activity of trypsin.

XVIII. Demonstration of Immunoglobulin Activity

An assay for SECP activity measures the ability of SECP to recognize and precipitate antigens from serum. This activity can be measured by the quantitative precipitin reaction. (Golub, E. S. et al. (1987) Immunology: A Synthesis, Sinauer Associates, Sunderland, MA, pages 113-115.) SECP is isotopically labeled using methods known in the art. Various serum concentrations are added to constant amounts of labeled SECP. SECP-antigen complexes precipitate out of solution and are collected by centrifugation. The amount of precipitable SECP-antigen complex is proportional to the amount of radioisotope detected in the precipitate. The amount of precipitable SECP-antigen complex is plotted against the serum concentration. For various serum concentrations, a characteristic precipitin curve is obtained, in which the amount of precipitable SECP-antigen complex initially increases proportionately with increasing serum concentration, peaks at the equivalence point, and then decreases proportionately with further increases in serum concentration. Thus, the amount of precipitable SECP-antigen complex is a measure of SECP activity which is characterized by sensitivity to both limiting and excess quantities of antigen.

Alternatively, an assay for SECP activity measures the expression of SECP on the cell surface. cDNA encoding SECP is transfected into a non-leukocytic cell line. Cell surface proteins are labeled with biotin (de la Fuente, M.A. et.al. (1997) Blood 90:2398-2405). Immunoprecipitations are performed using SECP-specific antibodies, and immunoprecipitated samples are analyzed using SDS-PAGE and immunoblotting techniques. The ratio of labeled immunoprecipitant to unlabeled immunoprecipitant is proportional to the amount of SECP expressed on the cell surface.

Alternatively, an assay for SECP activity measures the amount of cell aggregation induced by overexpression of SECP. In this assay, cultured cells such as NIH3T3 are transfected with cDNA encoding SECP contained within a suitable mammalian expression vector under control of a strong

promoter. Cotransfection with cDNA encoding a fluorescent marker protein, such as Green Fluorescent Protein (CLONTECH), is useful for identifying stable transfectants. The amount of cell agglutination, or clumping, associated with transfected cells is compared with that associated with untransfected cells. The amount of cell agglutination is a direct measure of SECP activity.

5

Various modifications and variations of the described methods and systems of the invention will be apparent to those skilled in the art without departing from the scope and spirit of the invention. Although the invention has been described in connection with certain embodiments, it should be understood that the invention as claimed should not be unduly limited to such specific
10 embodiments. Indeed, various modifications of the described modes for carrying out the invention which are obvious to those skilled in molecular biology or related fields are intended to be within the scope of the following claims.

Table 1

Incyte Project ID	Polypeptide SEQ ID NO:	Incyte Polypeptide ID	Polynucleotide SEQ ID NO:	Incyte Polynucleotide ID
3211795	1	3211795CD1	68	3211795CB1
6813464	2	6813464CD1	69	6813464CB1
2156540	3	2156540CD1	70	2156540CB1
894939	4	894939CD1	71	894939CB1
4620890	5	4620890CD1	72	4620890CB1
5514146	6	5514146CD1	73	5514146CB1
7474769	7	7474769CD1	74	7474769CB1
065296	8	065296CD1	75	065296CB1
231994	9	231994CD1	76	231994CB1
538054	10	538054CD1	77	538054CB1
1259305	11	1259305CD1	78	1259305CB1
1483702	12	1483702CD1	79	1483702CB1
1519324	13	1519324CD1	80	1519324CB1
1630169	14	1630169CD1	81	1630169CB1
1664253	15	1664253CD1	82	1664253CB1
1864715	16	1864715CD1	83	1864715CB1
1929395	17	1929395CD1	84	1929395CB1
1987737	18	1987737CD1	85	1987737CB1
2122866	19	2122866CD1	86	2122866CB1
2123981	20	2123981CD1	87	2123981CB1
2200177	21	2200177CD1	88	2200177CB1
2319255	22	2319255CD1	89	2319255CB1
2792452	23	2792452CD1	90	2792452CB1
2853088	24	2853088CD1	91	2853088CB1
2949004	25	2949004CD1	92	2949004CB1
3011670	26	3011670CD1	93	3011670CB1
3242083	27	3242083CD1	94	3242083CB1
3363391	28	3363391CD1	95	3363391CB1
3703614	29	3703614CD1	96	3703614CB1
4000975	30	4000975CD1	97	4000975CB1
4598831	31	4598831CD1	98	4598831CB1

Table 1 (cont.)

Incyte Project ID	Polypeptide SEQ ID NO:	Incyte Polypeptide ID	Polynucleotide SEQ ID NO:	Incyte Polynucleotide ID
4992201	32	4992201CD1	99	4992201CB1
5441583	33	5441583CD1	100	5441583CB1
1639243	34	1639243CD1	101	1639243CB1
1335166	35	1335166CD1	102	1335166CB1
166894	36	166894CD1	103	166894CB1
217969	37	217969CD1	104	217969CB1
335237	38	335237CD1	105	335237CB1
938306	39	938306CD1	106	938306CB1
1448129	40	1448129CD1	107	1448129CB1
1761049	41	1761049CD1	108	1761049CB1
1959587	42	1959587CD1	109	1959587CB1
2303463	43	2303463CD1	110	2303463CB1
2512281	44	2512281CD1	111	2512281CB1
2755924	45	2755924CD1	112	2755924CB1
2796369	46	2796369CD1	113	2796369CB1
3010920	47	3010920CD1	114	3010920CB1
3360955	48	3360955CD1	115	3360955CB1
3409459	49	3409459CD1	116	3409459CB1
4102938	50	4102938CD1	117	4102938CB1
4124601	51	4124601CD1	118	4124601CB1
4180577	52	4180577CD1	119	4180577CB1
5265807	53	5265807CD1	120	5265807CB1
5405979	54	5405979CD1	121	5405979CB1
7481109	55	7481109CD1	122	7481109CB1
6247114	56	6247114CD1	123	6247114CB1
3243866	57	3243866CD1	124	3243866CB1
7475633	58	7475633CD1	125	7475633CB1
1431268	59	1431268CD1	126	1431268CB1
2414185	60	2414185CD1	127	2414185CB1
5266594	61	5266594CD1	128	5266594CB1
7610617	62	7610617CD1	129	7610617CB1

Table 1 (cont.)

Incyte Project ID	Polypeptide SEQ ID NO:	Incyte Polypeptide ID	Polynucleotide SEQ ID NO:	Incyte Polynucleotide ID
1902436	63	1902436CD1	130	1902436CB1
2310369	64	2310369CD1	131	2310369CB1
6180576	65	6180576CD1	132	6180576CB1
2274523	66	2274523CD1	133	2274523CB1
1801820	67	1801820CD1	134	1801820CB1

Table 2

Polypeptide SEQ ID NO:	Incyte Polypeptide ID	GenBank ID NO:	Probability score	GenBank Homolog
1	3211795CD1	g5764665	3.80E-156	[Homo sapiens] cerebral cell adhesion molecule Starzyk, R.M. et al. (2000) J. Infect. Dis. 181:181-187
2	6813464CD1	g14794726	0	[fl][Homo sapiens] CUB and sushi multiple domains 1 protein Sun, P. C. et al. (2001) Transcript map of the 8p23 putative tumor suppressor region. Genomics. 75:17-25
3	2156540CD1	g7259265	2.40E-28	[Mus musculus] contains transmembrane (TM) region Inoue, S. et al. (2000) Growth suppression of Escherichia coli by induction of expression of mammalian genes with transmembrane or ATPase domains. Biochem. Biophys. Res. Commun. 268:553-561
4	894939CD1	g7242876	4.30E-171	[Mus musculus] kidney predominant protein
5	4620890CD1	g13874437	1.00E-134	[fl][Homo sapiens] cerebral protein-11
6	5514146CD1	g1256001	5.70E-46	[Homo sapiens] LIV-1 protein
7	7474769CD1	g13603845	0	[fl][Mus musculus] ribonuclease/angiogenin inhibitor 2 Wang, P. J. et al. (2001) An abundance of X-linked genes expressed in spermatogonia. Nat. Genet. 27:422- 426
34	1639243CD1	g927209	5.00E-20	[Homo sapiens] alpha 1C adrenergic receptor isoform 2
35	1335166CD1	g3002527	1.80E-21	[Homo sapiens] neuronal thread protein AD7c-NTP
57	3243866CD1	g6273399	3.30E-27	[Homo sapiens] melanoma-associated antigen MG50 Weiler, S. R. et al. (1994) Assignment of a human melanoma associated gene MG50 (D2S448) to chromosome 2p25.3 by fluorescence in situ hybridization. Genomics 22:243-244
58	7475633CD1	g3097285	1.20E-65	[Rattus norvegicus] ZOG

Table 3

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
1	3211795CD1	622	S125 S138 S153 S262 S272 S383 S542 S569 S591 S612 T300 T406 T562 T584 Y126 Y140 Y281 Y492	N184 N381 N96	Signal_cleavage: M1-G29 Signal peptide: M1-P32 Lysyl hydrolase Lysyl hydro: V165-I240 PROCOLLAGENLYSINE 2OXOGLUTARATE 5DIOXYGENASE PRECURSOR LYSYL HYDROXYLASE OXIDOREDUCTASE DIOXYGENASE SIGNAL IRON PD009947: P52-L291 LYSYL HYDROXYLASE CHAIN DM07920 Q02809 1-726: R155-L291	SPSCAN HMMER HMMER_PFAM BLAST_PRODOR BLAST_DOMO
2	6813464CD1	529	S109 S213 S243 S251 S299 S56 T113 T192 T273 T405 T416 T99	N312 N411	Signal peptide: M1-A21 Signal peptide: M1-S22 CUB domains: C136-Y244, C309-F414 Sushi domain (SCR repeat): C74-C131, C252-C305, C422-C479 Sushi domain proteins PF00084: G93-Y104, N296-C305 GLYCOPROTEIN DOMAIN EGFLIKE PROTEIN PRECURSOR SIGNAL RECEPTOR INTRINSIC FACTORB12 REPEAT PD000165: S144-Y244, T314-F414 C1R/C1S REPEAT DM00162 I49540 748-862: C136-Y244, C309-F410 DM00162 P98063 755-862: S144-Y244, T314-F410 DM00162 I49540 592-708: C305-F414 DM00162 P13497 595-703: T314-F414	SPSCAN HMMER HMMER_PFAM HMMER_PFAM BLIMPS_PFAM BLAST_PRODOR BLAST_DOMO

Table 3 (cont.)

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
3	2156540CD1	204	S101 S124 S196 S63 S83 T46 T51 T54 T88 Y140	N122	signal_peptide: M1-A24 signal_cleavage: M1-P27 Immunoglobulin domain ig: G55-V144	HMMER SPSCAN HMMER_PFAM
4	894939CD1	406	S151 S200 S226 S249 S255 S288 S400	N134 N159 N187 N230 N333 N65 N95	signal_peptide: M1-G35 signal_cleavage: M1-G35 transmembrane_domain: L369-G387 Leucine Zipper: L371-L392	HMMER SPSCAN HMMER HMMER
5	4620890CD1	477	S151 S166 S183 S226 S235 S25 S253 S279 S360 S388 S70 T13 T325 T87	N201 N270	signal_peptide: M439-C460 transmembrane_domain: I418-F435 PROTEIN TESTISSPECIFIC TEX28KIAA0481 HH1480 PD156308: P55-Q145 PD184097: L287-W463	HMMER MOTIFS BLAST_PRODOR
6	5514146CD1	691	S130 S136 S149 S160 S250 S293 S30 S312 S329 S392 S470 S51 S513 S530 S565 S598 S645 T24 T246 T26 T362 T658 T67	N134 N162 N201 N242 N290 N59	signal_cleavage: M1-S23 transmembrane_domain: I603-L621, W632-P651, P661-Y685 PROTEIN GUFA TRANSMEMBRANE MEMBRANE INTERGENIC REGION INNER CONSERVED SIMILARITY MYXOCOCCUS PD004603: K533-L682	SPSCAN HMMER BLAST_PRODOR

Table 3 (cont.)

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
7	7474769CD1	919	S125 S146 S157 S230 S275 S37 S393 S40 S462 S523 S543 S591 S632 S638 S661 S733 T217 T574 T679 T751 T814 T885 T896 T906 T915 Y109 Y47 Y645	N228 N601 N682 N731 N852 N904	signal_cleavage: M1-T16 RIBONUCLEASE INHIBITOR REPEAT LEUCINEREPEAT 3DSTRUCTURE PLACENTAL RAI RI RECEPTOR RIBONUCLEASE/ANGIOGENIN PD017636: L737-L842, L794-E889, L681-S787 Atp_Gtp_A: G80-T87	SPSCAN BLAST_PRODOM MOTIFS
8	065296CD1	178	S24 S7 S79 S81 S87	N116 N74 N75	signal_cleavage: M1-G26	SPSCAN
9	231994CD1	310	S281 S93 T136	N101 N219	signal_cleavage: M1-P24	SPSCAN
10	538054CD1	559	S121 S150 S216 S258 S294 S385 S474 S71 T274 T28 T363 T41	N237 N26 N292 N346	signal_peptide: M1-V24 signal_cleavage: M1-N26 transmembrane domain: M1-I25 BRUSH BORDER 61.9KD PROTEINPRECURSOR SIGNAL PD144534: L8-C555	HMNER SPSCAN HMNER BLAST_PRODOM
11	1259305CD1	477	S113 S129 S211 S273 S471 T135 T139 T142 T18 T338	N111 N209	signal_cleavage: M1-S40 transmembrane domain: M23-A39, F56-I74, F250-L270, M315-L335, W406-N430	SPSCAN HMNER
12	1483702CD1	176	S108 S132 S138 S142 S3 S34 T32	N47	signal_peptide: M111-S132	HMNER
13	1519324CD1	190	S119 S164 S189 S23 S86 S93 S98		signal_cleavage: M1-S23	SPSCAN
14	1630169CD1	75		N50	signal_cleavage: M1-A27 signal_peptide: M1-S22 transmembrane domain: L3-S22 sulfatases signature sulfatase_2.prf: S22-W76	SPSCAN HMNER HMNER PROFILES SCAN

Table 3 (cont.)

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
15	1664253CD1	265	S12 S14 S187 S30 S64 T162	N197	signal_peptide: M116-A264	HMMER
16	1864715CD1	202	S33 T167 T65	N10	signal_peptide: M1-A19 signal_cleavage: M1-G15 transmembrane_domain: I173-R193	HMMER SPSCAN HMMER
17	1929395CD1	111	T4 T67		signal_cleavage: M1-G34	SPSCAN
18	1987737CD1	105	S75 T38		signal_cleavage: M1-G28	SPSCAN
19	2122866CD1	717	S172 S3 S359 S397 S454 S46 S469 S488 S5 S526 S567 S594 S7 T117 T20 T410 Y213	N130 N274 N44 N520 N666	signal_cleavage: M133-P197 PROTEIN FACTOR REPEAT CELL CHROMOSOME INTERGENIC REGION HOST C1 JASMONATE PD013240: N149-N306 RGD: R432-D434	SPSCAN BLAST_PRODOR MOTIFS
20	2123981CD1	580	S109 S120 S210 S260 S322 T121 T178 T272 T31 T327 T360 T403 T415 T450 T460 T545 T70 T73 T75 Y522	N276 N337 N543 N99	signal_peptide: M1-K21 signal_cleavage: M1-A16	HMMER SPSCAN
21	2200177CD1	172	S113 S148 S93 T138 Y121		signal_cleavage: M1-C50 transmembrane_domain: L34-I51	SPSCAN HMMER
22	2319255CD1	256	S242 S49 T116 T130 T3 T66	N200	signal_cleavage: M1-A30	SPSCAN
23	2792452CD1	93	S15 S24		signal_peptide: M1-S22 transmembrane_domain: L41-I58 signal_cleavage: M1-G52	HMMER HMMER
24	2853088CD1	112	S89		signal_peptide: M1-G36 signal_cleavage: M1-A27 J SP NEURONAL THREAD PD003801: P30-R71	HMMER SPSCAN BLAST_PRODOR

Table 3 (cont.)

SEQ ID NO.	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
25	2949004CD1	186	S26 S91 S94 T19 T65		signal_peptide: M75-S94 transmembrane_domain: I139-L163	HMMER
26	3011670CD1	487	S356 S36 S445 S89 T227 T262 T313 T44 T60 T91 Y66	N135 N203 N34 N397	signal_peptide: M128-A150 transmembrane_domain: W106-N126, I136-L155, V170-V188, M306-C323, L459-F479 Atp_Gtp_A: G53-T60	SPSCAN HMMER
27	3242083CD1	350	S311 T261 T73	N294	signal_cleavage: M1-A66 transmembrane_domain: V43-W65, M84-C102, F172-V191	MOTIFS SPSCAN HMMER
28	3363391CD1	450	S132 S140 S196 S247 S289 S380 S385 S417 S428 T208 T214 T357 T445 Y368 Y403	N131 N206 N287	signal_peptide: M144-R162 transmembrane_domain: L146-R162	HMMER HMMER
29	3703614CD1	400	S372 S71 S82 T32 T348 T358 Y362 Y76		signal_peptide: M173-T193 transmembrane_domain: M173-Y190, P198-V221, L293-V310	HMMER HMMER
30	4000975CD1	133	S54 S7		signal_peptide: M1-A24 signal_cleavage: M1-M25	HMMER SPSCAN
31	4598831CD1	359	S139 S300 S359 T355 T41 T63		signal_peptide: M327-A344 do MEMBRANE; YOL002C; CHROMOSOME; C30D11.11; DM02642 Q09749 49-323: Y55-G321 DM02642 S62569 169-441: T41-H291 DM02642 Q09910 169-441: T41-H291 DM02642 S61982 50-325: D46-Q310	HMMER BLAST_DOMO
32	4992201CD1	72	S17 S32 T7	N14	signal_cleavage: M1-G65 transmembrane_domain: M39-F58	SPSCAN HMMER
33	5441583CD1	112			signal_peptide: M53-A80 Heme oxygenase signature heme_oxygenase.prp: T9-M53	HMMER PROFILES SCAN

Table 3 (cont.)

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
34	1639243CD1	149	S93		Signal cleavage: M1-Q45 J SP NEURONAL THREAD PD003801:R89-I130	SPSCAN BLAST_PRODOM
35	1335166CD1	97	S30 S36 S52 S70		Signal peptide: M1-P21 Signal cleavage: M1-A43 RNA EDITING PROTOONCOGENE REPEAT PD005171:S3-P45	HMMER SPSCAN BLAST_PRODOM
36	166894CD1	104	S25 T60		Signal cleavage: M1-T27 Signal peptide: M1-S29 Insulin-like growth factor binding proteins IGFB: C80-M93 P-II protein signatures pi1_glnb_cter.prf: L20-G76 Ribosomal protein S19e signature ribosomal_s19e.prf: S24-P77 Plant PEC family metallothionein signature PR00877 L75-R82	SPSCAN HMMER HMMER_PFRAM PROFILES SCAN PROFILES SCAN BLIMPS_PRINTS
37	217969CD1	99	T9		Signal cleavage: M1-S35 RNA EDITING PROTOONCOGENE REPEAT PD005171: F29-L66 P_value 2.6e-07	SPSCAN BLAST_PRODOM
38	335237CD1	80	S38	N42	Signal peptide: M1-S18 Signal cleavage: M1-S18	HMMER SPSCAN
39	938306CD1	96	S41 S45 T79	N58	Signal peptide: M1-G18 Signal cleavage: M1-G18	HMMER SPSCAN
40	1448129CD1	92	S62		Signal peptide: M1-A28 Signal cleavage: M1-G31 Aminotransferases class-I pyridoxal-phosphate attachment site asp_aminotransferase.prf: K55-A92 Respiratory-chain NADH dehydrogenase 75 Kd subunit signatures complex1_75k_3.prf: F5-Q79	HMMER SPSCAN PROFILES SCAN PROFILES SCAN

Table 3 (cont.)

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
41	1761049CD1	77	S37		Signal cleavage: M1-C66	SPSCAN
42	1959587CD1	75	S48 S70		Transmembrane domain: Y40-M64	HMNER
43	2303463CD1	85			Signal cleavage: M1-G26	SPSCAN
44	2512281CD1	89	S61 S62 T29 T42 T58		Signal peptide: M1-G23	HMNER
45	2755924CD1	123			Signal cleavage: M1-A30	SPSCAN
46	2796369CD1	159	S97 T35		Transmembrane domain: L22-A48	HMNER
47	3010920CD1	77	S7		Signal peptide: M1-R31	SPSCAN
48	3360955CD1	130	S15 S32 S95 T25		Signal cleavage: M1-P22	HMNER
49	3409459CD1	97			5-hydroxytryptamine 1B receptor PR00513 R31-T42	BLIMPS_PRINTS
50	4102938CD1	74	S3		Signal peptide: M1-A19	HMNER
51	4124601CD1	74	S12		Signal cleavage: M1-S24	SPSCAN
					Transmembrane domain: I38-A53	HMNER
					Signal cleavage: M1-G27	SPSCAN
					Tyrosine specific protein phosphatases active site tyr_phosphatase.prf: L42-E100	PROFILESAN
					Signal cleavage: M1-T33	SPSCAN
					Signal cleavage: M1-T14	SPSCAN
					Orexin receptor signature PR01064 N78-T89	BLIMPS_PRINTS
					Signal peptide: M1-G20	HMNER
					Signal cleavage: M1-G20	SPSCAN
					Signal cleavage: M1-L25	SPSCAN
					Signal peptide: M1-G33	HMNER
					Hemerythrins signature hemerythrins.prf: Q21-L60	PROFILESAN
					Signal cleavage: M1-G46	SPSCAN
					Annexin Type III Signature PR00199 W17-A26	BLIMPS_PRINTS

Table 3 (cont.)

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
52	4180577CD1	151	S115 S116 S128 S63 S68 S77 T93	N40	Signal peptide: M1-A22	HMMER
53	5265807CD1	137	S123 S79		Signal cleavage: M1-S21	SPSCAN
					Signal cleavage: M1-A42	SPSCAN
					Signal peptide: M1-G31	HMMER
					Ank repeat ank: Y13-K45 L46-R78 S79-Y113	HMMER_PPFAM
					Ank repeat PD0007 L77-A89	BLIMPS_PRODOR
					Ank repeat protein motif PF00023 G80-H89	BLIMPS_PPFAM
54	5405979CD1	137	S48 T118		Signal peptide: M1-A24	HMMER
					Signal cleavage: M1-L16	SPSCAN
55	7481109CD1	205	T181 T31 Y141	N131 N143	Signal peptide: M2-Q22	HMMER
					Signal cleavage: M1-F26	SPSCAN
					HYPERGLYCEMIC HORMONE TY PR00548: L158-C168	BLIMPS_PRINTS
					Pancreatic ribonucleases RNaseA: F104-C168	HMMER_PPFAM

Table 3 (cont.)

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
56	6247114CD1	199	S108 S113 S123 S136 S146 S42 S67 S91 T162 T175 T22 T31 T77	N111 N61 N89	Signal peptide: M1-S21 Signal cleavage: M1-A18 PANCREATIC RIBONUCLEASE FAMILY DM00621 P39873 28-166: S73-V183 Pancreatic ribonuclease family signature rnase_pancreatic.prf: G92-K138 signal_peptide: M1-P21 signal_cleavage: M1-A17 transmembrane_domain: M529-V554 Polycystic kidney disease protein signature PR00500B: P241-E261 Leucine-rich repeat signature PR00019A: L149-I162 Leucine Rich Repeat LR: S172-H195, K196-A219, R52-T75, S76-R99, N100-S123, N124-F147, A148-V171. Leucine rich repeat C-terminal domain LRRCT: N240-E285 Immunoglobulin domain ig: G301-A359 RGD: R311-D313	HMMER SPSCAN BLAST_DOMO PROFILES SCAN HMMER SPSCAN HMMER BLIMPS_PRINTS BLIMPS_PRINTS HMMER_PFAM HMMER_PFAM HMMER_PFAM MOTIFS
57	3243866CD1	719	S108 S204 S255 S276 S400 S410 S618 S640 S685 S692 S76 T141 T198 T270 T305 T348 T454 T467 T480 T510 T647 T661 T668 T691 T704 T713	N330 N339 N382 N406 N452 N671 N672 N73		

Table 3 (cont.)

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
58	7475633CD1	383	S290 S31 S45 T159 T188 T258 T328	N157	signal_peptide: M1-A26 signal_cleavage: M1-G21 transmembrane_domain: V308-R330 EGF DM00003 I48324 235-285: C40-K90 DM00003 Q07645 107-174: G113-V173 DM00003 P10041 275-331: W75-K131 DM00003 P80370 72-127: Q76-K131 Laminin-type EGF-like domain BL01248 C48-C60 Type II EGF-like signature PR0010C: N194-F204 Type III EGF-like signature PR0011D: C39-C57 EGF-like domain EGF: C216-C247, C29-C57, C60-C88, C95-C128, C135-C171, C178-C209 EGF-like domain Egf: C46-C57, C77-C88, C117-C128, C160-C171, C198-C209, C236-C247 Zinc_Finger_C3hc4: C7-I16 Asp or Asn hydroxylation site found in EGF-domain proteins Asx Hydroxyl: C189-C200, C227-C238 signal_peptide: M1-A35 signal_peptide: M1-G21 signal_peptide: M1-G29 signal_peptide: M1-S21	HMMER SPSCAN HMMER BLAST_DOMO BLIMPS_BLOCKS BLIMPS_PRINTS BLIMPS_PRINTS HMMER_PFAM MOTIFS MOTIFS HMMER HMMER HMMER HMMER
59	1431268CD1	126	S2 S54			
60	2414185CD1	137	T65			
61	5266594CD1	77	T35			
62	7610617CD1	110	S46 T83			

Table 3 (cont.)

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
63	1902436CD1	103	T72		signal peptide: M1-S18 signal_cleavage: M1-S18	HMMER
					transmembrane_domain: I4-I23	SPSCAN
64	2310369CD1	192	S153 S28 S56 T186 T57		signal_cleavage: M1-A32	HMMER
					transmembrane_domain: V114-T134	SPSCAN
65	6180576CD1	310	S154 S158 S201 S307 S5 S79 S93 T225 T253 T55 T71 Y163		signal_cleavage: M1-G61	HMMER
						SPSCAN
66	2274523CD1	135	S12		signal_cleavage: M1-S44	SPSCAN
67	1801820CD1	205	S110 S126 S198 S68 T185 T34	N67	signal_cleavage: M1-X36 Bowman-Birk serine protease inhibitor family signature bowman_birk.prf: C104-C194	SPSCAN PROFILES SCAN

Table 4

Polynucleotide SEQ ID NO:	Incyte Polynucleotide ID	Sequence Length	Selected Fragment(s)	Sequence Fragments	5' Position	3' Position
68	3211795CB1	2569	1-513, 2404-2569	8052177J1 (FTUBTUE01)	1775	2401
				6756010H1 (SINTFER02)	453	1192
				6756010J1 (SINTFER02)	1047	1810
				688428H1 (BRAITDR03)	1419	1955
				6944985H1 (FTUBTUR01)	1950	2416
				2185379F6 (PROSNOT26)	2103	2569
				6900869H1 (MUSLTR02)	729	1401
				7317029H2 (BONFTXT01)	1	560
69	6813464CB1	2387	104-184, 546-2387	GNN.98051897_000002_00 2	185	519
				GNN.g9408716_002.edit	1	397
				71926854V1	1924	2387
				8085111H1 (BRACDIK08)	355	919
				71926415V1	1527	2107
				71930431V1	1025	1536
				71928622V1	1515	2047
				6813464F6 (ADRETUR01)	559	1255
				g2204647	1385	1959
				71147007V1	1221	1916
70	2156540CB1	1959	1918-1959, 723-1073	7714520H1 (SINTFEE02)	705	1255
				g5553166	1	422
				6269361H1 (BRAIFEN03)	1255	1927
				7192294H2 (BRATDIC01)	31	462
				7460559H1 (LIVRTUE01)	264	948
				825696H1 (PROSNOT06)	1674	1941
				71113024V1	1	569
				71113961V1	935	1562
71	894939CB1	1562	1-253	71111802V1	582	1258
				71178768V1	500	1205

Table 4 (cont.)

Polynucleotide SEQ ID NO:	Incyte Polynucleotide ID	Sequence Length	Selected Fragment (s)	Sequence Fragments	5' Position	3' Position
72	4620890CB1	3425	1-355, 1127-2529	3730714F6 (SMCCNON03)	3124	3425
				7126752H1 (COLNDIY01)	2199	2753
				71260484V1	964	1582
				4041777H1 (BRAITUT26)	1	299
				7089611R8 (BRAUTDR03)	439	1129
				7186380H1 (BONRPEC01)	1573	2194
				5608905T6 (MONOTXS05)	1116	1696
				6216030H1 (MUSCDIT06)	2803	3267
				7089611F8 (BRAUTDR03)	56	726
				5721967H1 (SEMVNOT05)	2457	2825
				6365636F6 (ARTANOT07)	1758	2350
				4058011F6 (SPLNNOT13)	2882	3293
				70403184D1	2245	2786
				70401863D1	2527	3130
73	5514146CB1	3130	2654-3130, 1611-2305, 992-1198	70942023V1	1409	2085
				6770687H1 (BRAUNOR01)	239	723
				7998254H1 (BRAITUC02)	1	446
				5514146F6 (BRADDIR01)	1172	1710
				70941524V1	1731	2281
				4253735F6 (BRADDIR01)	564	1091
				5511496F8 (BRADDIR01)	813	1434
				55077715J1	1	392
				5544880H1 (TESTNOC01)	1456	1649
				92100308	2911	3172
				GNN.96693125_000016_00	200	2789
				2		
				8038136J1 (SMCRUNE01)	1273	2094
				71058917V1	872	1524
74	7474769CB1	3172	646-970, 2450-2614, 1939-2217	94393469	1	357
				71247541V1	846	1456
				71059505V1	219	883
75	065296CB1	2094	1826-2094			

Table 4 (cont.)

Polynucleotide SEQ ID NO:	Incyte Polynucleotide ID	Sequence Length	Selected Fragment(s)	Sequence Fragments	5' Position	3' Position
76	231994CB1	1119	1-163	6424977H1 (BRSTUNT01)	19	703
				1735672F6 (COLNNOT22)	309	846
				3484368H1 (KIDNNOT31)	1	154
				404248R6 (TMLR3DT01)	620	1119
77	538054CB1	3321	1513-1629, 2956-3321	2304848R6 (NGANNOT01)	1671	2208
				7055893V1	546	1031
				70561551V1	403	982
				6951008H1 (BRAITDR02)	1039	1798
				530151R6 (BRAINOT03)	2887	3321
				6884875H1 (BRAHTDR03)	984	1466
78	1259305CB1	2646	1-1540	7262670H1 (UTRETMC01)	1	433
				4880167H1 (UTRMTMT01)	2499	2782
				351423T6 (LVENNOT01)	2563	3147
				6995967H1 (BRAXTDR17)	1820	2548
				661278T6 (BRAINOT03)	1959	2624
				6782356J1 (OVARDIR01)	584	1338
				8009718H1 (NOSEDIC02)	1	556
				71294610V1	2104	2646
				6782356H1 (OVARDIR01)	1332	2025
				70979178V1	1068	1745
				70978234V1	349	977
				SXAF03884V1	1	513
79	1483702CB1	1749	1712-1749, 1130-1181	1478338F1 (CORPNOT02)	404	960
				1477349F2 (CORPNOT02)	1178	1731
				1832249R6 (BRAINON01)	1421	1749
				2571527R6 (HIPOAZT01)	713	1222
80	1519324CB1	2339	1987-2339, 1-66	569251H1 (MMLR3DT01)	1203	1522
				4435164H2 (LUNGNOT38)	918	1216
				268264H1 (HNT2NOT01)	473	759
				3504827H1 (ADRENOT11)	2031	2319
				3803618H1 (BLADTUT03)	1521	1831

Table 4 (cont.)

Polynucleotide SEQ ID NO:	Incyte Polynucleotide ID	Sequence Length	Selected Fragment(s)	Sequence Fragments	5' Position	3' Position
81				318988H1 (THYMNON04)	1459	1783
				1556067T6 (BLADTUT04)	1662	2316
				1484363F6 (CORPNOT02)	1	478
				1347568H1 (PROSNOT11)	2175	2339
				2959710H2 (ADRENOT09)	379	692
				4103403F6 (BRSTTUT17)	597	1168
				3802959H1 (BLADTUT03)	1197	1499
				6771071J1 (BRAUNOR01)	1	547
				6915144J1 (PITUDIR01)	386	815
				71296019V1	654	1006
82		1006	1-1006	6975645F8 (BRAHTDR04)	571	1050
				1511079H1 (LUNGNOT14)	1	206
				7988873H1 (UTRSTUC01)	280	911
				6975645R8 (BRAHTDR04)	153	745
				2866801H1 (KIDNNOT20)	505	837
				3736166H1 (SMCCNOS01)	1503	1774
				2972289H2 (HEAONOT02)	168	461
				1864715T6 (PROSNOT19)	1175	1760
				4585982H1 (OVARNOT13)	928	1195
				1864715F6 (PROSNOT19)	396	828
83		1774	1-91, 864- 1189	95431009	1	452
				6521196H1 (CONFDT02)	584	1192
				6426607H1 (LUNGNON07)	223	830
				7159765H1 (HNT2TXC01)	1	437
				6403390H1 (UTRENOT10)	2330	2608
				5203330H2 (STOMNOT08)	1168	1429
				8184678H1 (EYERNON01)	1894	2382
				999772R1 (KIDNTUT01)	1416	1963
				5776203H1 (BRAINOT20)	806	1360
				2569101R6 (HIPAZT01)	1	375
84		2608	1-230, 2378-2608, 1364-1424, 1859-1908	1987737R6 (LUNGAST01)	855	1336
85		1336	1-362, 396- 455			

Table 4 (cont.)

Polynucleotide SEQ ID NO:	Incyte Polynucleotide ID	Sequence Length	Selected Fragment(s)	Sequence Fragments	5' Position	3' Position
86	2122866CB1	3062	2470-3062	7651140J1 (STOMTDE01)	287	883
				70822631V1	379	1013
				1344981T6 (PROSNOT11)	2525	3062
				3373914T6 (CONNTUT05)	1720	2315
				5106924H1 (PROSTUS19)	1482	1736
87	2123981CB1	2543	2458-2543	70581466V1	571	1299
				7760472J1 (THYMNOE02)	1	697
				1861920T6 (PROSNOT19)	1986	2566
				888838R1 (STOMTUT01)	1125	1703
				2502591T6 (CONUTUT01)	1788	2498
				70818388V1	760	1446
				2123981F6 (BRSTNOT07)	189	695
				7184911H1 (BONRFEC01)	1	201
				7687944J1 (PROSTME06)	658	1336
				7182169H1 (BONRFEC01)	104	694
				1746464F6 (STOMTUT02)	1479	2021
				1782982R6 (PGANNON02)	2215	2543
				70818385V1	1272	1844
				1570015F6 (UTRSNOT05)	2016	2528
				70715536V1	229	873
88	2200177CB1	873	1-873	2200177T6 (SPLNFET02)	1	653
89	2319255CB1	2134	1-1366	70972802V1	747	1350
				70972902V1	1606	2134
				2319255R6 (OVARNOT02)	1205	1770
				71291442V1	1388	1893
				70973293V1	369	927
90	2792452CB1	1886	1-675, 1174-1420, 1824-1886	2719008F6 (THYRN0T09)	1	541
				2631710F6 (COLNTUT15)	1358	1886
				2792452T6 (COLNTUT16)	1154	1816
				7056068H1 (BRALNON02)	1	611
				70747733V1	764	1290

Table 4 (cont.)

Polynucleotide SEQ ID NO:	Incyte Polynucleotide ID	Sequence Length	Selected Fragment(s)	Sequence Fragments	5' Position	3' Position
91	2853088CB1	928	1-368, 541-928	70750933V1 6909603F6 (PITUDIR01) 2853088F6 (BRSTTUT13)	541 403 1	1199 928 510
92	2949004CB1	962	1-73, 524-574	6406737H1 (UTREDIT10) 5773379H1 (BRAINOT20) 5571763H1 (TLYMNOT08) 5763826H1 (PROSBPT02) 1454503T6 (PENITUT01) 70529802V1 70527923V1 70529939V1 6784388H1 (SINITMC01) 2130537H1 (KIDNNOT05) 2493973H1 (ADRETUT05) 8035143H1 (SMCRUNE01) 70729684V1 7764378H1 (URETTUE01) 8035143J1 (SMCRUNE01) 7407517H1 (UTREDME05) 1691309F6 (PROSTUT10) 4647910F6 (PROSTUT20) 70701284V1 7600868H1 (ESOGTME01) 7600868J1 (ESOGTME01) 2055126T6 (BEPINOT01) 70977847V1 70979848V1 70977881V1 70977648V1 70255942V1 70256236V1	207 103 1 569 1958 492 1101 1266 270 2381 1 641 1272 549 1 1294 1890 1 1462 415 701 1750 1 615 579 1047 1 62	755 676 212 962 2571 1220 1684 2004 744 2644 317 1410 1875 1215 640 1790 2378 702 1803 935 1335 2348 619 1114 1096 1597 528 653
93	3011670CB1	2644	2581-2644, 2402-2433, 269-296, 1289-1370			
94	3242083CB1	1875	1605-1624, 760-1228, 140-182			
95	3363391CB1	2378	1-1315			
96	3703614CB1	1597	1-230, 1513-1597			
97	4000975CB1	653	1-653			

Table 4 (cont.)

Polynucleotide SEQ ID NO:	Incyte Polynucleotide ID	Sequence Length	Selected Fragment(s)	Sequence Fragments	5' Position	3' Position
98	4598831CB1	3090	1-265, 1019-1311, 3031-3090	6450571H1 (BRAINOC01)	2461	3039
				7604731H1 (COLRTUE01)	277	871
				7280444H1 (BMARTXE01)	1	363
				7604731J1 (COLRTUE01)	1034	1618
				6531268H1 (LUNPTMC01)	2134	2785
				1717922F6 (UCMCNOT02)	1652	2072
				9884719	2646	3090
				7069481H1 (BRAUTDR02)	498	1097
				7078716H1 (BRAUTDR04)	1807	2285
				2598065F6 (OVRTUT02)	2715	3063
99	4992201CB1	1274	1-132, 1180-1274	7361768H1 (BRAIFEE05)	1093	1715
				7408255H1 (UTREDME05)	190	754
				6169092H1 (UTRSTDT01)	1004	1274
				264080H1 (HNT2AGT01)	1	357
				4992201F6 (LIVRTUT11)	356	1023
				7619787J1 (KIDNTUE01)	1036	1514
				3216178F6 (TESTNOT07)	1	608
				71580026V1	543	1108
				71579908V1	651	1167
				2789856F6 (COLNTUT16)	1188	1375
100	5441583CB1	1514	1-341, 1360-1514, 644-737	8215917H1 (FIBRTXC01)	435	1224
				71249544V1	1	632
				91447886	892	1380
				1659932T6 (URETTUT01)	658	1319
				6929148H1 (SINITMR01)	374	942
				1335166F6 (COLNNT013)	1	512
				2914860T6 (THYMFEET03)	1046	1631
				8056981J1 (ESOGTUE01)	1	641
				6623672H1 (LIVRTMR01)	1310	1815
				5880645F8 (LIVRNON08)	516	1276
101	1639243CB1	1380	1332-1380, 1-505	71249544V1	1	632
				91447886	892	1380
				1659932T6 (URETTUT01)	658	1319
				6929148H1 (SINITMR01)	374	942
				1335166F6 (COLNNT013)	1	512
				2914860T6 (THYMFEET03)	1046	1631
				8056981J1 (ESOGTUE01)	1	641
				6623672H1 (LIVRTMR01)	1310	1815
				5880645F8 (LIVRNON08)	516	1276
				71249544V1	1	632
102	1335166CB1	942	1-356, 806- 942	71249544V1	1	632
				91447886	892	1380
				1659932T6 (URETTUT01)	658	1319
				6929148H1 (SINITMR01)	374	942
				1335166F6 (COLNNT013)	1	512
				2914860T6 (THYMFEET03)	1046	1631
				8056981J1 (ESOGTUE01)	1	641
				6623672H1 (LIVRTMR01)	1310	1815
				5880645F8 (LIVRNON08)	516	1276
				71249544V1	1	632
103	166894CB1	1815	1-488, 1719-1815	71249544V1	1	632
				91447886	892	1380
				1659932T6 (URETTUT01)	658	1319
				6929148H1 (SINITMR01)	374	942
				1335166F6 (COLNNT013)	1	512
				2914860T6 (THYMFEET03)	1046	1631
				8056981J1 (ESOGTUE01)	1	641
				6623672H1 (LIVRTMR01)	1310	1815
				5880645F8 (LIVRNON08)	516	1276
				71249544V1	1	632

Table 4 (cont.)

Polynucleotide SEQ ID NO:	Incyte Polynucleotide ID	Sequence Length	Selected Fragment(s)	Sequence Fragments	5' Position	3' Position
104	217969CB1	1120	801-1120, 1-542	1406479T6 (LATRTUT02) 1406479F6 (LATRTUT02) 7127039H1 (COLNDIY01) 6516316H1 (THYMDIT01)	238 1 546 1	897 500 1120 517
105	335237CB1	535	1-120, 389- 535	335237R6 (EOSIHET02)	188	535
106	938306CB1	1188	1-36, 806- 1010, 232- 330	6837739H1 (BRSTN02) 1843293R6 (COLNNOT08)	580 1	1188 610
107	1448129CB1	638	1-162	1448129F6 (PLACNOT02) 95152438	1 219	450 638
108	1761049CB1	648	152-576, 1- 91, 626-648	1448129T6 (PLACNOT02) 3335890F7 (BRAIFET01)	217 148	599 648
109	1959587CB1	1181	1-484	55079223J1 2418384T6 (HNT3AZT01) 6562708H1 (MCLDTXT04) 1959587H1 (BRSTN04)	1 623 227 1	647 1181 911 228
110	2303463CB1	1291	765-815, 965-1000, 1141-1291	3087044F6 (HEAONOT03) 70726533V1 92052638 70725786V1 5371544F6 (BRAINOT22) 4533095F8 (MYEPTXT01)	77 587 678 659 1 758	657 1253 1291 1270 626 1287
111	2512281CB1	594	324-594	2512281F6 (CONUTUT01) 7177623T8 (BRAXDIC01)	1 201	522 594
112	2755924CB1	852	1-173, 290- 852	70252828V1 71442317V1	384 1	852 603
113	2796369CB1	1361	1-244, 802- 1135	5289268F6 (LIVRTUS02) 5716622H1 (PANCONOT16) 2796369F6 (NPOLNOT01) 6264155H1 (MCLDTXN03)	1 772 562 483	554 1361 1128 950

Table 4 (cont.)

Polynucleotide SEQ ID NO:	Incyte Polynucleotide ID	Sequence Length	Selected Fragment(s)	Sequence Fragments	5' Position	3' Position
135	3010920CB1.comp	1650	1-235, 1416-1650	71297582V1 3010920F6 (MUSCNOT07) 70990273V1 70991132V1 70138150V1 70772246V1 3360955F6 (PROSBPT02) 70137857V1 g793031	1 805 1154 624 993 485 1272 21 1	646 1441 1650 1290 1531 1001 1845 501 360
115	3360955CB1	1845	898-1404, 162-191, 1- 29, 1714- 1845	2096304H1 (BRAITUT02) 4560677H1 (KERATXT01) 8049214H1 (LUNGTUS02) 3409459F6 (PROSTUS08) 7590996H1 (LIVRNOC07) 71665573V1 4102938F6 (BRSTTUT17)	1 942 628 377 134 330 1	230 1061 946 798 641 1085 507
136	3409459CB1.comp	1061	1-627, 435- 1061	71157275V1 4124601F6 (BRSTTUT26) 72081891D1 72079822D1 7370367H1 (ADREFEC01) 72360174D1 7371968H2 (BRAIFEE04) 7675267J2 (NOSETUE01) 72360060D1 72082904D1	1 200 2015 2570 2278 1351 413 1 1405 721	520 870 2579 3394 2821 2147 992 607 2231 1412
117	4102938CB1	1085	971-1085, 1-257, 684- 783, 464- 486	71163549V1 71397639V1 71396875V1	560 1942 1547	1218 2343 2197
118	4124601CB1	870	285-870			
119	4180577CB1	3394	1-766, 3366-3394, 1487-1519			
120	5265807CB1	2343	56-519, 671-1627			

Table 4 (cont.)

Polynucleotide SEQ ID NO:	Incyte Polynucleotide ID	Sequence Length	Selected Fragment(s)	Sequence Fragments	5' Position	3' Position
121		751	718-751	71163327V1	1285	1845
				7142204H1 (LIVRDIT07)	1183	1750
				7098309H1 (BRACDIR02)	1	574
				71166280V1	540	1183
122	5405979CB1	751	718-751	96711145	1	240
123	7481109CB1 6247114CB1	618 979	1-618 1-384	2009872R6 (TESTNOT03)	303	751
				2009872T6 (TESTNOT03)	17	749
124	3243866CB1	3012	1-43, 2836- 2885, 2307- 2520, 1275- 1779	GNN.99588388_010	1	618
				2008559R6 (TESTNOT03)	538	979
				6247114F8 (TESTNOT17)	1	741
				8184163H1 (EYERNON01)	1215	1932
				6768945J1 (BRAUNOR01)	1	569
				70960957V1	2437	3012
				70869814V1	1551	2189
				7056045H1 (BRALNON02)	2131	2749
125	7475633CB1	1600	1-72	3243866F6 (BRAINOT19)	735	1313
				70870169V1	2283	2835
				4783652F6 (BRATNOT03)	503	1074
				6874650F6 (EPIMUNN04)	1	911
				72379524V1	900	1600
				6874650T6 (EPIMUNN04)	746	1587
				5223435H1 (OVARIT07)	650	922
				6747053H1 (BRAFNOT02)	286	909
126	1431268CB1	1001	335-547	4139565H1 (BRSTTMT01)	709	1001
				70255212V1	1	559
				7034594R6 (SINTFER03)	1	607
				6897648H1 (LIVRTMR01)	899	1424
127	2414185CB1	1424	1-46, 297- 1424	2414185F6 (HNT3AZT01)	570	1052
				7764520H1 (URETTUE01)	730	1282
128	5266594CB1	1282	1-34, 990- 1282	5545864T8 (TESTNOC01)	1	829
				7373102F6 (BRAIFEE04)	29	632

Table 4 (cont.)

Polynucleotide SEQ ID NO:	Incyte Polynucleotide ID	Sequence Length	Selected Fragment (s)	Sequence Fragments	5' Position	3' Position
129	7610617CB1	642	314-642	7736885J1 (BRAITUE01)	348	963
				3881415H1 (SPINNOT11)	721	1008
				7610617F6 (KIDCTME01)	97	642
130	1902436CB1	1326	845-1326, 740-776, 40-705	7610617R6 (KIDCTME01)	1	609
				70882432V1	717	1326
				2216468F6 (SINTFET03)	1	480
				70880085V1	538	986
131	2310369CB1	1486	139-395, 67-106	70879063V1	412	953
				2310369R6 (NGANNO01)	107	557
				3375613H1 (CONNTUT05)	1	265
				72069492V1	784	1486
				6438845H1 (BRAENOT02)	252	854
132	6180576CB1	1523	1-40	6116528T8 (SINITMT04)	762	1496
				7179642H1 (BRAXDIC01)	526	1151
				6146183H1 (BRANDIT03)	1185	1523
				6577826F8 (BRANDIT04)	1	571
				2274523T6 (PROSNON01)	378	848
133	2274523CB1	848	87-567	2083066F6 (UTRSNOT08)	87	583
				2068885H1 (PROSNOT26)	1	348
				1315807F1 (BLADTUT02)	2434	2758
				2645114F6 (OVARTUT03)	1991	2502
				1798924F6 (COLNNOT27)	556	1212
134	1801820CB1	2758	2484-2758	7079130H1 (BRAUTDR04)	1	600
				7409525H1 (BRAIFEJ02)	631	1253
				6479333H1 (PROSTMC01)	910	1424
				2303084R6 (BRSTNOT05)	1294	1816
				2926367F6 (TLYMNOT04)	1693	2229

Table 5

Polynucleotide SEQ ID NO:	Incyte Project ID	Representative Library
68	3211795CB1	OVARTUE01
69	6813464CB1	ADRETUR01
70	2156540CB1	BRAINOT09
71	894939CB1	BONSTUT01
72	4620890CB1	MONOTXS05
73	5514146CB1	BRADDIR01
74	7474769CB1	TESTNOC01
75	065296CB1	PLACNOB01
76	231994CB1	BRSTTUT01
77	538054CB1	LVENNOT01
78	1259305CB1	KIDNNOT19
79	1483702CB1	CORPNOT02
80	1519324CB1	KIDNNOT09
81	1630169CB1	BRAGNON02
82	1664253CB1	BRAHTR04
83	1864715CB1	PROSNOT19
84	1929395CB1	COLNTUT03
85	1987737CB1	BRAINON01
86	2122866CB1	LIVRNON08
87	2123981CB1	PGANNON02
88	2200177CB1	KERANOT01
89	2319255CB1	TESTNOT03
90	2792452CB1	COLNTUT16
91	2853088CB1	PITUDIR01
92	2949004CB1	BRAITUT07
93	3011670CB1	PENITUT01
94	3242083CB1	FIBRPXS07
95	3363391CB1	PROSTUT10
96	3703614CB1	SINJNOT03
97	4000975CB1	HNT2AZS07
98	4598831CB1	UCMCNOT02

Table 5 (cont.)

Polynucleotide SEQ ID NO:	Incyte Project ID	Representative Library
99	4992201CB1	HNT2AGT01
100	5441583CB1	BSTMNON02
101	1639243CB1	UTRSNOT06
102	1335166CB1	COLANOT13
103	166894CB1	LIVRNOT01
104	217969CB1	LATRTUT02
105	335237CB1	EOSIHT02
106	938306CB1	CERVNOT01
107	1448129CB1	PLACNOT02
108	1761049CB1	PITUNOT03
109	1959587CB1	HEAONOT03
110	2303463CB1	BRSTNOT05
111	2512281CB1	CONUTUT01
112	2755924CB1	THPLAZS08
113	2796369CB1	LIVRTUS02
135	3010920CB1.comp	MUSCNOT07
115	3360955CB1	PROSBPT02
136	3409459CB1.comp	STOMTUT02
117	4102938CB1	BRSTTUT17
118	4124601CB1	BRSTTUT26
119	4180577CB1	BRAIFEE04
120	5265807CB1	UTRENON03
121	5405979CB1	TESTNOT03
123	6247114CB1	TESTNOT17
124	3243866CB1	BRAIFEN03
125	7475633CB1	EPIMUNN04
126	1431268CB1	SINTBST01
127	2414185CB1	BRSTNOT04
128	5266594CB1	STOMFET01
129	7610617CB1	KIDCTME01
130	1902436CB1	OVARNOT07
131	2310369CB1	FIBRTXS07

Table 5 (cont.)

Polynucleotide SEQ ID NO:	Incyte Project ID	Representative Library
132	6180576CB1	HIPONOT01
133	2274523CB1	PROSNON01
134	1801820CB1	COLNNOT27

Table 6

Library	Vector	Library Description
ADRETUR01	PCDNA2.1	This random primed library was constructed using RNA isolated from left upper pole, adrenal gland tumor tissue removed from a 52-year-old Caucasian male during nephroureterectomy and local destruction of renal lesion. Pathology indicated grade 3 adrenal cortical carcinoma forming a mass that infiltrated almost the whole adrenal parenchyma and extended to adjacent adipose tissue. A metastatic tumor nodule was identified in the hilar region. The renal vein was infiltrated by tumor and the neoplastic process was present at the resection margin of the renal vein. Fragments of adrenal cortical carcinoma and thrombus were found in the inferior vena cava. Patient history included abnormal weight loss. Family history included skin cancer, type I diabetes, and neurotic depression.
BONSTUT01	pINCY	Library was constructed using RNA isolated from sacral bone tumor tissue removed from an 18-year-old Caucasian female during an exploratory laparotomy with soft tissue excision. Pathology indicated giant cell tumor of the sacrum. Patient history included a soft tissue malignant neoplasm. Family history included prostate cancer.
BRADDIR01	pINCY	Library was constructed using RNA isolated from diseased choroid plexus tissue of the lateral ventricle, removed from the brain of a 57-year-old Caucasian male, who died from a cerebrovascular accident.
BRAGNON02	pINCY	This normalized substantia nigra tissue library was constructed from 4.2x10e7 independent clones from a substantia nigra tissue library. Starting RNA was made from RNA isolated from substantia nigra tissue removed from an 81-year-old Caucasian female who died from a hemorrhage and ruptured thoracic aorta due to atherosclerosis. Pathology indicated moderate atherosclerosis involving the internal carotids, bilaterally; microscopic infarcts of the frontal cortex and hippocampus; and scattered diffuse amyloid plaques and neurofibrillary tangles, consistent with age. Grossly, the leptomeninges showed only mild thickening and hyalinization along the superior sagittal sinus. The remainder of the leptomeninges was thin and contained some congested blood vessels. Mild atrophy was found mostly in the frontal poles and lobes, and temporal lobes, bilaterally. Microscopically, there were pairs of Alzheimer type II astrocytes within the deep layers of the neocortex. There was increased satellitosis around neurons in the deep gray matter in the middle frontal cortex. The amygdala contained rare diffuse plaques and neurofibrillary tangles. The posterior hippocampus contained a

Table 6 (cont.)

Library	Vector	Library Description
BRAHTDR04	PCDNA2.1	microscopic area of cystic cavitation with hemosiderin-laden macrophages surrounded by reactive gliosis. Patient history included sepsis, cholangitis, post-operative atelectasis, pneumonia CAD, cardiomegaly due to left ventricular hypertrophy, splenomegaly, arteriolonephrosclerosis, nodular colloidal goiter, emphysema, CHF, hypothyroidism, and peripheral vascular disease. The library was normalized in two rounds using conditions adapted from Soares et al., PNAS (1994) 91:9228-9232 and Bonaldo et al., Genome Research 6 (1996):791, except that a significantly longer (48 hours/round) reannealing hybridization was used.
		This random primed library was constructed using RNA isolated from archaocortex, anterior hippocampus tissue removed from a 55-year-old Caucasian female who died from cholangiocarcinoma. Pathology indicated mild meningeal fibrosis predominately over the convexities, scattered axonal spheroids in the white matter of the cingulate cortex and the thalamus, and a few scattered neurofibrillary tangles in the entorhinal cortex and the periaqueductal gray region. Pathology for the associated tumor tissue indicated well-differentiated cholangiocarcinoma of the liver with residual or relapsed tumor. Patient history included cholangiocarcinoma, post-operative Budd-Chiari syndrome, biliary ascites, hydrothorax, dehydration, malnutrition, oliguria and acute renal failure. Previous surgeries included cholecystectomy and resection of 85% of the liver.
BRAIFEE04	PINCY	This 5' biased random primed library was constructed using RNA isolated from brain tissue removed from a Caucasian male fetus who was stillborn with a hypoplastic left heart at 23 weeks' gestation.
BRAIFEN03	PINCY	This normalized fetal brain tissue library was constructed from 3.26 million independent clones from a fetal brain library. Starting RNA was made from brain tissue removed from a Caucasian male fetus, who was stillborn with a hypoplastic left heart at 23 weeks' gestation. The library was normalized in 2 rounds using conditions adapted from Soares et al., PNAS (1994) 91:9228 and Bonaldo et al., Genome Research (1996) 6:791, except that a significantly longer (48 hours/round) reannealing hybridization was used.
BRAINON01	PSPORT1	This normalized library was made from 4.88 million independent clones from a brain tissue library. Starting RNA was isolated from brain tissue removed from a 26-year-old Caucasian male during cranioplasty and excision of a cerebral meningeal lesion. Pathology for the associated tumor tissue indicated a grade 4

Table 6 (cont.)

Library	Vector	Library Description
BRAINOT09	pINCY	oligoastrocytoma in the right fronto-parietal part of the brain. Library was constructed using RNA isolated from brain tissue removed from a Caucasian male fetus, who died at 23 weeks' gestation.
BRAITUT07	pINCY	Library was constructed using RNA isolated from left frontal lobe tumor tissue removed from the brain of a 32-year-old Caucasian male during excision of a cerebral meningeal lesion. Pathology indicated low grade desmoplastic neuronal neoplasm, type not otherwise specified. The lesion formed a firm, circumscribed cyst-associated mass involving white matter and cortex. No definite glial component was evident to suggest a diagnosis of ganglioglioma. Family history included atherosclerotic coronary artery disease.
BRSTNOT04	PSPORT1	Library was constructed using RNA isolated from breast tissue removed from a 62-year-old East Indian female during a unilateral extended simple mastectomy. Pathology for the associated tumor tissue indicated an invasive grade 3 ductal carcinoma. Patient history included benign hypertension, hyperlipidemia, and hematuria. Family history included cerebrovascular and cardiovascular disease, hyperlipidemia, and liver cancer.
BRSTNOT05	PSPORT1	Library was constructed using RNA isolated from breast tissue removed from a 58-year-old Caucasian female during a unilateral extended simple mastectomy. Pathology for the associated tumor tissue indicated multicentric invasive grade 4 lobular carcinoma. Patient history included skin cancer, rheumatic heart disease, osteoarthritis, and tuberculosis. Family history included cerebrovascular and cardiovascular disease, breast and prostate cancer, and type I diabetes.
BRSTTUT01	PSPORT1	Library was constructed using RNA isolated from breast tumor tissue removed from a 55-year-old Caucasian female during a unilateral extended simple mastectomy. Pathology indicated invasive grade 4 mammary adenocarcinoma of mixed lobular and ductal type, extensively involving the left breast. The tumor was identified in the deep dermis near the lactiferous ducts with extracapsular extension. Seven mid and low and five high axillary lymph nodes were positive for tumor. Proliferative fibrocytic changes were characterized by apocrine metaplasia, sclerosing adenosis, cyst formation, and ductal hyperplasia without atypia. Patient history included atrial tachycardia, blood in the stool, and a benign breast neoplasm. Family history included benign hypertension, atherosclerotic coronary artery disease, cerebrovascular disease, and depressive disorder.

Table 6 (cont.)

Library	Vector	Library Description
BRSTTUT17	pINCY	Library was constructed using RNA isolated from left breast tumor tissue removed from a 65-year-old Caucasian female during a unilateral radical mastectomy. Pathology indicated invasive and in situ grade 3, nuclear grade 2 ductal carcinoma. Patient history included hyperlipidemia and uterine leiomyoma. Family history included stomach cancer, myocardial infarction, atherosclerotic coronary artery disease, prostate cancer, benign hypertension, breast cancer, and hyperlipidemia.
BRSTTUT26	pINCY	Library was constructed using RNA isolated from breast tumor tissue removed from an adult female. The breast carcinoma tumor tissue was found to have low vascular density and was considered resting.
BSTMNON02	PSPORT1	This normalized brain stem library was constructed from 2.84 million independent clones from a brain stem library. Starting RNA was made from the brain stem tissue of a 72-year-old Caucasian male who died from myocardial infarction. Patient history included coronary artery disease, insulin-dependent diabetes mellitus, and arthritis. Normalization and hybridization conditions were adapted from Soares et al. (PNAS (1994) 91:9228).
CERVNOT01	PSPORT1	Library was constructed using RNA isolated from the uterine cervical tissue of a 35-year-old Caucasian female during a vaginal hysterectomy with dilation and curettage. Pathology indicated mild chronic cervicitis. Family history included atherosclerotic coronary artery disease and type II diabetes.
COLNNOT13	pINCY	Library was constructed using RNA isolated from ascending colon tissue of a 28-year-old Caucasian male with moderate chronic ulcerative colitis.
COLNNOT27	pINCY	Library was constructed using RNA isolated from diseased cecal tissue removed from 31-year-old Caucasian male during a total intra-abdominal colectomy, appendectomy, and permanent ileostomy. Pathology indicated severe active Crohn's disease involving the colon from the cecum to the rectum. There were deep rake-like ulcerations which spared the intervening mucosa. The ulcers extended into the muscularis, and there was transmural inflammation. Patient history included an irritable colon. Previous surgeries included a colonoscopy.
COLNTUT03	pINCY	Library was constructed using RNA isolated from colon tumor tissue obtained from the sigmoid colon of a 62-year-old Caucasian male during a sigmoidectomy and permanent colostomy. Pathology indicated invasive grade 2 adenocarcinoma. One lymph node contained metastasis with extranodal extension. Patient history

Table 6 (cont.)

Library	Vector	Library Description
		included hyperlipidemia, cataract disorder, and dermatitis. Family history included benign hypertension, atherosclerotic coronary artery disease, hyperlipidemia, breast cancer, and prostate cancer.
COLNTUT16	pINCY	Library was constructed using RNA isolated from colon tumor tissue obtained from a 60-year-old Caucasian male during a left hemicolectomy. Pathology indicated an invasive grade 2 adenocarcinoma, forming a sessile mass. Patient history included thrombophlebitis, inflammatory polyarthropathy, prostatic inflammatory disease, and depressive disorder. Previous surgeries included resection of the rectum. Family history included atherosclerotic coronary artery disease and colon cancer.
CONUTUT01	pINCY	Library was constructed using RNA isolated from sigmoid mesentery tumor tissue obtained from a 61-year-old female during a total abdominal hysterectomy and bilateral salpingo-oophorectomy with regional lymph node excision. Pathology indicated a metastatic grade 4 malignant mixed mullerian tumor present in the sigmoid mesentery at two sites.
CORPNOT02	pINCY	Library was constructed using RNA isolated from diseased corpus callosum tissue removed from the brain of a 74-year-old Caucasian male who died from Alzheimer's disease.
EOSIHET02	PBLUESCRIPT	Library was constructed using RNA isolated from peripheral blood cells apheresed from a 48-year-old Caucasian male. Patient history included hypereosinophilia. The cell population was determined to be greater than 77% eosinophils by Wright's staining.
EPIMUNN04	pINCY	This normalized mammary epithelial cell tissue library was constructed from 3.28 million independent clones from an epithelial cell tissue library. Starting RNA was made from untreated mammary epithelial cell tissue removed from a 21-year-old female. The library was normalized in two rounds using conditions adapted from Soares et al., PNAS (1994) 91:9228-9232 and Bonaldo et al., Genome Research 6 (1996):791, except that a significantly longer (48 hours/round) reannealing hybridization was used.
FIBRTXS07	pINCY	This subtracted library was constructed using 1.3 million clones from a dermal fibroblast library and was subjected to two rounds of subtraction hybridization with 2.8 million clones from the an untreated dermal fibroblast tissue library. The starting library for subtraction was constructed using RNA isolated from treated dermal fibroblast tissue removed from the breast of a 31-year-old

Table 6 (cont.)

Library	Vector	Library Description
		Caucasian female. The cells were treated with 9CIS retinoic acid. The hybridization probe for subtraction was derived from a similarly constructed library from RNA isolated from untreated dermal fibroblast tissue from the same donor. Subtractive hybridization conditions were based on the methodologies of Swaroop et al., NAR (1991) 19:1954 and Bonaldo, et al., Genome Research (1996) 6:791.
HEAONOT03	pINCY	Library was constructed using RNA isolated from aortic tissue removed from a 27-year-old Caucasian female, who died from an intracranial bleed.
HIPONOT01	PELUESCRIPT	Library was constructed using RNA isolated from the hippocampus tissue of a 72-year-old Caucasian female who died from an intracranial bleed. Patient history included nose cancer, hypertension, and arthritis.
HNT2AGT01	PELUESCRIPT	Library was constructed at Stratagene (STR937233), using RNA isolated from the hNT2 cell line derived from a human teratocarcinoma that exhibited properties characteristic of a committed neuronal precursor. Cells were treated with retinoic acid for 5 weeks and with mitotic inhibitors for two weeks and allowed to mature for an additional 4 weeks in conditioned medium.
HNT2AZS07	PSPORT1	This subtracted library was constructed from RNA isolated from an hNT2 cell line (derived from a human teratocarcinoma that exhibited properties characteristic of a committed neuronal precursor) treated for three days with 0.35 micromolar AZ. The hybridization probe for subtraction was derived from a similarly constructed library from untreated hNT2 cells. 3.08M clones from the AZ-treated library were subjected to three rounds of subtractive hybridization with 3.04M clones from the untreated library. Subtractive hybridization conditions were based on the methodologies of Swaroop et al. (NAR (1991) 19:1954) and Bonaldo et al. (Genome Research (1996) 6:791).
KERANOT01	PELUESCRIPT	Library was constructed using RNA isolated from neonatal keratinocytes obtained from the leg skin of a spontaneously aborted black male.
KIDCTME01	PCDNA2.1	This 5' biased random primed library was constructed using RNA isolated from kidney cortex tissue removed from a 65-year-old male during nephroureterectomy. Pathology indicated the margins of resection were free of involvement. Pathology for the matched tumor tissue indicated grade 3 renal cell carcinoma, clear cell type, forming a variegated multicystic mass situated within the mid-portion of the kidney. The tumor invaded deeply into but not through the renal capsule.

Table 6 (cont.)

Library	Vector	Library Description
KIDNNOT09	pINCY	Library was constructed using RNA isolated from the kidney tissue of a Caucasian male fetus, who died at 23 weeks' gestation.
KIDNNOT19	pINCY	Library was constructed using RNA isolated from kidney tissue removed a 65-year-old Caucasian male during an exploratory laparotomy and nephroureterectomy. Pathology for the associated tumor tissue indicated a grade 1 renal cell carcinoma within the upper pole of the left kidney. Patient history included malignant melanoma of the abdominal skin, benign neoplasm of colon, cerebrovascular disease, and umbilical hernia. Family history included myocardial infarction, atherosclerotic coronary artery disease, cerebrovascular disease, prostate cancer, myocardial infarction, and atherosclerotic coronary artery disease.
LATRUT02	pINCY	Library was constructed using RNA isolated from a myxoma removed from the left atrium of a 43-year-old Caucasian male during annuloplasty. Pathology indicated atrial myxoma. Patient history included pulmonary insufficiency, acute myocardial infarction, atherosclerotic coronary artery disease, hyperlipidemia, and tobacco use. Family history included benign hypertension, acute myocardial infarction, atherosclerotic coronary artery disease, and type II diabetes.
LIVRN08	pINCY	This normalized library was constructed from 5.7 million independent clones from a pooled liver tissue library. Starting RNA was made from pooled liver tissue removed from a 4-year-old Hispanic male who died from anoxia and a 16 week female fetus who died after 16-weeks gestation from anencephaly. Serologies were positive for cytomegalovirus in the 4-year-old. Patient history included asthma in the 4-year-old. Family history included taking daily prenatal vitamins and mitral valve prolapse in the mother of the fetus. The library was normalized in 2 rounds using conditions adapted from Soares et al., PNAS (1994) 91:9228 and Bonaldo et al., Genome Research 6 (1996):791, except that a significantly longer (48 hours/round) reannealing hybridization was used.
LIVRN01	PBLUESCRIPT	Library was constructed at Stratagene, using RNA isolated from the liver tissue of a 49-year-old male.
LIVRTUS02	pINCY	This subtracted C3A liver tumor cell line tissue library was constructed using 6.4 million clones from a 3-methylcholthrene-treated hepatocyte library and was subjected to two rounds of subtraction hybridization with 1.72 million clones from an untreated C3A hepatocyte library. The starting library for subtraction was constructed using RNA isolated from a treated C3A hepatocyte cell line which is a

Table 6 (cont.)

Library	Vector	Library Description
LIVENOT01	PBLUESCRIPT	Library was constructed using RNA isolated from the left ventricle of a 51-year-old Caucasian female, who died from an intracranial bleed.
MONOTXS05	pINCY	Library was constructed using 7.5 million clones from a treated monocyte library and subjected to two rounds of subtraction hybridization with 1.03×10^7 clones from a second treated monocyte library. The starting library for subtraction was constructed using RNA from treated monocytes from peripheral blood obtained from a 42-year-old female. The cells were pre-treated for 1 hour with 10 ng/ml anti-interleukin-10 (anti-IL-10). Lipopolysaccharide (LPS) at 5 ng/ml was added and monocytes harvested after 24 hours. Monocytes were isolated from buffy coat by adherence to plastic. The hybridization probe for subtraction was derived from a similarly constructed library using RNA isolated from monocyte tissue, treated with interleukin-10 (IL10) and lipopolysaccharide (LPS) from the same donor. Subtractive hybridization conditions were based on the methodologies of Swaroop et al. NAR 19 (1991):1954 and Bonaldo, et al. Genome Research 6 (1996):791.
MUSCNOT07	pINCY	Library was constructed using RNA isolated from muscle tissue removed from the forearm of a 38-year-old Caucasian female during a soft tissue excision. Pathology for the associated tumor tissue indicated intramuscular hemangioma. Family history included breast cancer, benign hypertension, cerebrovascular disease, colon cancer, and type II diabetes.
OVARNOT07	pINCY	Library was constructed using RNA isolated from left ovarian tissue removed from a 28-year-old Caucasian female during a vaginal hysterectomy and removal of the fallopian tubes and ovaries. The tissue was associated with multiple follicular cysts, endometriosis in a weakly proliferative phase, and chronic cervicitis of the cervix with squamous metaplasia. Family history included benign hypertension, hyperlipidemia, and atherosclerotic coronary artery disease.

Table 6 (cont.)

Library	Vector	Library Description
OVARTUE01	PCDNA2.1	This 5' biased random primed library was constructed using RNA isolated from left ovary tumor tissue removed from a 44-year-old female. Pathology indicated grade 4 (of 4) serous carcinoma replacing both the right and left ovaries forming solid mass cystic masses. Neoplastic deposits are identified in para-ovarian soft tissue, on the surface of the uterus and scattered throughout the myometrium and cervix. Transverse colon was positive for metastatic disease. Multiple staging biopsies including diaphragm, bladder, liver, sigmoid rectal serosa, inguinal lymph nodes and left colic gutter are metastatically involved.
PENITUT01	PINCY	Library was constructed using RNA isolated from tumor tissue removed from the penis of a 64-year-old Caucasian male during penile amputation. Pathology indicated a fungating invasive grade 4 squamous cell carcinoma involving the inner wall of the foreskin and extending onto the glans penis. Patient history included benign neoplasm of the large bowel, atherosclerotic coronary artery disease, angina pectoris, gout, and obesity. Family history included malignant pharyngeal neoplasm, chronic lymphocytic leukemia, and chronic liver disease.
PGANNON02	PSPORT1	This normalized paraganglion library was constructed with 5.48e6 independent clones from a paraganglionic tissue library. Starting RNA was made from paraganglionic tissue removed from a 46-year-old Caucasian male during exploratory laparotomy. Pathology indicated a benign paraganglioma and was associated with a grade 2 renal cell carcinoma. The normalization and hybridization conditions were adapted from Soares et al. (PNAS (1994) 91:9228-9232) using a significantly longer (48-hour) reannealing hybridization period.
PITUDIR01	PCDNA2.1	This random primed library was constructed using RNA isolated from pituitary gland tissue removed from a 70-year-old female who died from metastatic adenocarcinoma. Pathology for the brain indicated moderate Alzheimer disease and mild carotid and cerebral atherosclerosis. The cerebral hemispheres, frontal and temporal lobes, white matter, and hippocampus showed mild atrophy, bilaterally. There were numerous neurofibrillary tangles, neuritic and diffuse amyloid plaques deposited throughout most neocortical areas. Most of the diffuse plaques were in the superficial layers, with more core and neuritic amyloid plaques in the deep cortical layers. Most of the tangles were found in small interneurons, rather than in the large pyramidal neurons. The areas that were most involved with plaques and tangles were the entorhinal cortex, temporal cortex, and superior parietal lobes.

Table 6 (cont.)

Library	Vector	Library Description
		There was marked vacuolization of the superficial layers throughout all neocortical areas examined. The hippocampus contained numerous neurofibrillary tangles (predominantly in the CA-1 field), diffuse and neuritic plaques, as well as granulovacuolar degeneration within the pyramidal cell neurons. There were neuritic plaques with scattered neurofibrillary tangles within the amygdala. The thalamus had scattered diffuse plaques. There was mild pigment incontinence in the substantia nigra compacta. The periaqueductal gray matter showed mild gliosis. Diffuse plaques were found within the superior colliculus. Neurofibrillary tangles were found within the pons. The neurons of the locus ceruleus were ballooned and contain eosinophilic foamy material with very little neuromelanin pigment.
PITUNOT03	PSPORT1	Library was constructed using RNA isolated from pituitary tissue of a 46-year-old Caucasian male, who died from colon cancer. Serologies were negative. Patient history included arthritis, peptic ulcer disease, and tobacco use. Patient medications included Tagamet and muscle relaxants.
PLACNOB01	PELUESCRIPT	Library was constructed using RNA isolated from placenta.
PLACNOT02	pINCY	Library was constructed using RNA isolated from the placental tissue of a Hispanic female fetus, who was prematurely delivered at 21 weeks' gestation. Serologies of the mother's blood were positive for CMV (cytomegalovirus).
PROSBPT02	pINCY	Library was constructed using RNA isolated from diseased prostate tissue removed from a 65-year-old Caucasian male during a radical prostatectomy. Pathology indicated benign prostatic hyperplasia (BPH). Pathology for the associated tumor tissue indicated adenocarcinoma, Gleason grade 3+4. One (of 7) right pelvic lymph nodes was positive for metastatic adenocarcinoma. The patient presented with induration and elevated prostate specific antigen (PSA). Patient history included a benign neoplasm of the large bowel and benign hypertension.
PROSNON01	PSPORT1	This normalized prostate library was constructed from 4.4 M independent clones from a prostate library. Starting RNA was made from prostate tissue removed from a 28-year-old Caucasian male who died from a self-inflicted gunshot wound. The normalization and hybridization conditions were adapted from Soares, M.B. et al. (1994) Proc. Natl. Acad. Sci. USA 91:9228-9232, using a longer (19 hour) reannealing hybridization period.

Table 6 (cont.)

Library	Vector	Library Description
PROSNOT19	pINCY	Library was constructed using RNA isolated from diseased prostate tissue removed from a 59-year-old Caucasian male during a radical prostatectomy with regional lymph node excision. Pathology indicated adenofibromatous hyperplasia. Pathology for the associated tumor tissue indicated an adenocarcinoma (Gleason grade 3+3). The patient presented with elevated prostate-specific antigen (PSA). Patient history included colon diverticuli, asbestosis, and thrombophlebitis. Previous surgeries included a partial colectomy. Family history included benign hypertension, multiple myeloma, hyperlipidemia and rheumatoid arthritis.
PROSTUT10	pINCY	Library was constructed using RNA isolated from prostatic tumor tissue removed from a 66-year-old Caucasian male during radical prostatectomy and regional lymph node excision. Pathology indicated an adenocarcinoma (Gleason grade 2+3). Adenofibromatous hyperplasia was also present. The patient presented with elevated prostate specific antigen (PSA). Family history included prostate cancer and secondary bone cancer.
SINJNOT03	pINCY	Library was constructed using RNA isolated from duodenum tissue removed from the small intestine of a 16-year-old Caucasian male who died from head trauma. Patient history included a kidney infection.
SINTBST01	pINCY	Library was constructed using RNA isolated from ileum tissue obtained from an 18-year-old Caucasian female during bowel anastomosis. Pathology indicated Crohn's disease of the ileum, involving 15 cm of the small bowel. Family history included cerebrovascular disease and atherosclerotic coronary artery disease.
STOMFET01	pINCY	Library was constructed using RNA isolated from the stomach tissue of a Caucasian female fetus, who died at 20 weeks' gestation.
STOMTUT02	pINCY	Library was constructed using RNA isolated from stomach tumor tissue obtained from a 68-year-old Caucasian female during a partial gastrectomy. Pathology indicated a malignant lymphoma of diffuse large-cell type. Previous surgeries included cholecystectomy. Patient history included thalassemia. Family history included acute leukemia, malignant neoplasm of the esophagus, malignant stomach neoplasm, and atherosclerotic coronary artery disease.
TESTNOC01	PELUESCRIPT	This large size fractionated library was constructed using RNA isolated from testicular tissue removed from a pool of eleven, 10 to 61-year-old Caucasian males.

Table 6 (cont.)

Library	Vector	Library Description
TESTNOT03	PBLUESCRIPT	Library was constructed using RNA isolated from testicular tissue removed from a 37-year-old Caucasian male, who died from liver disease. Patient history included cirrhosis, jaundice, and liver failure.
TESTNOT17	PINCY	Library was constructed from testis tissue removed from a 26-year-old Caucasian male who died from head trauma due to a motor vehicle accident. Serologies were negative. Patient history included a hernia at birth, tobacco use (1 1/2 ppd), marijuana use, and daily alcohol use (beer and hard liquor).
THP1AZS08	PSPORT1	This subtracted THP-1 promonocyte cell line library was constructed using 5.76 x 1e6 clones from a 5-aza-2'-deoxycytidine (AZ) treated THP-1 cell library. Starting RNA was made from THP-1 promonocyte cells treated for three days with 0.8 micromolar AZ. The donor had acute monocytic leukemia. The hybridization probe for subtraction was derived from a similarly constructed library, made from 1 microgram of polyA RNA isolated from untreated THP-1 cells. 5.76 million clones from the AZ-treated THP-1 cell library were then subjected to two rounds of subtractive hybridization with 5 million clones from the untreated THP-1 cell library. Subtractive hybridization conditions were based on the methodologies of Swaroop et al., NAR (1991) 19:1954, and Bonaldo et al., Genome Research (1996) 6:791.
UCMCNOT02	PINCY	Library was constructed using RNA isolated from mononuclear cells obtained from the umbilical cord blood of nine individuals.

Table 6 (cont.)

Library	Vector	Library Description
UTRENON03	pINCY	<p>Library was constructed from 1.2 x 10⁶ independent clones from a uterine endometrial tissue library. Starting RNA was made from uterine endometrium tissue obtained from a 29-year-old Caucasian female during a vaginal hysterectomy and cystocele repair. Pathology indicated the endometrium was secretory and the cervix showed mild chronic cervicitis with focal squamous metaplasia. Pathology for the associated tumor tissue indicated an intramural uterine leiomyoma. Patient history included hypothyroidism, pelvic floor relaxation, incomplete T-12 injury (due to a motor vehicle accident) causing paraplegia and self catheterization. Previous surgeries included a normal delivery, a laminectomy, and a rhinoplasty. Family history included benign hypertension, type II diabetes, and hyperlipidemia. The libraries were normalized in two rounds using conditions adapted from Soares et al 1., PNAS (1994) 91:9928 and Bonaldo et al., Genome Research (1996) 6:791, except that a significantly longer (48 hours/round) reannealing hybridization was used.</p>
UTRSNOT06	pINCY	<p>Library was constructed using RNA isolated from myometrial tissue removed from a 50-year-old Caucasian female during a vaginal hysterectomy. Pathology indicated residual atypical complex endometrial hyperplasia. Pathology for the associated tissue removed during dilation and curettage indicated fragments of atypical complex hyperplasia and a single microscopic focus suspicious for grade 1 adenocarcinoma. Patient history included benign breast neoplasm, hypothyroid disease, polypectomy, and arthralgia. Family history included cerebrovascular disease, atherosclerotic coronary artery disease, hyperlipidemia, and chronic hepatitis.</p>

Table 7

Program	Description	Reference	Parameter Threshold
ABI FACTURA	A program that removes vector sequences and masks ambiguous bases in nucleic acid sequences.	Applied Biosystems, Foster City, CA.	
ABI/PARACEL FDF	A Fast Data Finder useful in comparing and annotating amino acid or nucleic acid sequences.	Applied Biosystems, Foster City, CA; Paracel Inc., Pasadena, CA.	Mismatch <50%
ABI AutoAssembler	A program that assembles nucleic acid sequences.	Applied Biosystems, Foster City, CA.	
BLAST	A Basic Local Alignment Search Tool useful in sequence similarity search for amino acid and nucleic acid sequences. BLAST includes five functions: blastp, blastn, blastx, tblastn, and tblastx.	Altschul, S.F. et al. (1990) J. Mol. Biol. 215:403-410; Altschul, S.F. et al. (1997) Nucleic Acids Res. 25:3389-3402.	ESTs: Probability value= 1.0E-8 or less Full Length sequences: Probability value= 1.0E-10 or less
FASTA	A Pearson and Lipman algorithm that searches for similarity between a query sequence and a group of sequences of the same type. FASTA comprises at least five functions: fasta, tfasta, fastx, tfastx, and ssearch.	Pearson, W.R. and D.J. Lipman (1988) Proc. Natl. Acad. Sci. USA 85:2444-2448; Pearson, W.R. (1990) Methods Enzymol. 183:63-98; and Smith, T.F. and M.S. Waterman (1981) Adv. Appl. Math. 2:482-489.	ESTs: fasta E value=1.06E-6 Assembled ESTs: fasta Identity= 95% or greater and Match length=200 bases or greater; fastx E value=1.0E-8 or less Full Length sequences: fastx score=100 or greater
BLIMPS	A BLOCKS IMProved Searcher that matches a sequence against those in BLOCKS, PRINTS, DOMO, PRODOM, and PFAM databases to search for gene families, sequence homology, and structural fingerprint regions.	Henikoff, S. and J.G. Henikoff (1991) Nucleic Acids Res. 19:6565-6572; Henikoff, J.G. and S. Henikoff (1996) Methods Enzymol. 266:88-105; and Attwood, T.K. et al. (1997) J. Chem. Inf. Comput. Sci. 37:417-424.	Probability value= 1.0E-3 or less
HMMER	An algorithm for searching a query sequence against hidden Markov model (HMM)-based databases of protein family consensus sequences, such as PFAM.	Krogh, A. et al. (1994) J. Mol. Biol. 235:1501-1531; Sonnhammer, E.L.L. et al. (1988) Nucleic Acids Res. 26:320-322; Durbin, R. et al. (1998) Our World View, in a Nutshell, Cambridge Univ. Press, pp. 1-350.	PFAM hits: Probability value= 1.0E-3 or less Signal peptide hits: Score= 0 or greater

Table 7 (cont.)

Program	Description	Reference	Parameter Threshold
ProfileScan	An algorithm that searches for structural and sequence motifs in protein sequences that match sequence patterns defined in Prosite.	Gribskov, M. et al. (1988) CABIOS 4:61-66; Gribskov, M. et al. (1989) Methods Enzymol. 183:146-159; Bairoch, A. et al. (1997) Nucleic Acids Res. 25:217-221.	Normalized quality score \geq GCG-specified "HIGH" value for that particular Prosite motif. Generally, score=1.4-2.1.
Phred	A base-calling algorithm that examines automated sequencer traces with high sensitivity and probability.	Ewing, B. et al. (1998) Genome Res. 8:175-185; Ewing, B. and P. Green (1998) Genome Res. 8:186-194.	
Phrap	A Phils Revised Assembly Program including SWAT and CrossMatch, programs based on efficient implementation of the Smith-Waterman algorithm, useful in searching sequence homology and assembling DNA sequences.	Smith, T.F. and M.S. Waterman (1981) Adv. Appl. Math. 2:482-489; Smith, T.F. and M.S. Waterman (1981) J. Mol. Biol. 147:195-197; and Green, P., University of Washington, Seattle, WA.	Score= 120 or greater; Match length= 56 or greater
Consed	A graphical tool for viewing and editing Phrap assemblies.	Gordon, D. et al. (1998) Genome Res. 8:195-202.	
SPScan	A weight matrix analysis program that scans protein sequences for the presence of secretory signal peptides.	Nielson, H. et al. (1997) Protein Engineering 10:1-6; Claverie, J.M. and S. Audic (1997) CABIOS 12:431-439.	Score=3.5 or greater
TMAP	A program that uses weight matrices to delineate transmembrane segments on protein sequences and determine orientation.	Persson, B. and P. Argos (1994) J. Mol. Biol. 237:182-192; Persson, B. and P. Argos (1996) Protein Sci. 5:363-371.	
TMHMMER	A program that uses a hidden Markov model (HMM) to delineate transmembrane segments on protein sequences and determine orientation.	Sonnhammer, E.L. et al. (1998) Proc. Sixth Intl. Conf. on Intelligent Systems for Mol. Biol., Glasgow et al., eds., The Am. Assoc. for Artificial Intelligence Press, Menlo Park, CA, pp. 175-182.	
Motifs	A program that searches amino acid sequences for patterns that matched those defined in Prosite.	Bairoch, A. et al. (1997) Nucleic Acids Res. 25:217-221; Wisconsin Package Program Manual, version 9, page M51-59, Genetics Computer Group, Madison, WI.	

What is claimed is:

1. An isolated polypeptide selected from the group consisting of:
 - a) a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-67,
 - b) a polypeptide comprising a naturally occurring amino acid sequence at least 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-67,
 - c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-67, and
 - d) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-67.
2. An isolated polypeptide of claim 1 selected from the group consisting of SEQ ID NO:1-67.
3. An isolated polynucleotide encoding a polypeptide of claim 1.
4. An isolated polynucleotide encoding a polypeptide of claim 2.
5. An isolated polynucleotide of claim 4 selected from the group consisting of SEQ ID NO:68-134.
6. A recombinant polynucleotide comprising a promoter sequence operably linked to a polynucleotide of claim 3.
7. A cell transformed with a recombinant polynucleotide of claim 6.
8. A transgenic organism comprising a recombinant polynucleotide of claim 6.
9. A method of producing a polypeptide of claim 1, the method comprising:
 - a) culturing a cell under conditions suitable for expression of the polypeptide, wherein said cell is transformed with a recombinant polynucleotide, and said recombinant polynucleotide comprises a promoter sequence operably linked to a polynucleotide encoding the polypeptide of claim 1, and
 - b) recovering the polypeptide so expressed.

10. A method of claim 9, wherein the polypeptide has an amino acid sequence selected from the group consisting of SEQ ID NO:1-67.

11. An isolated antibody which specifically binds to a polypeptide of claim 1.

12. An isolated polynucleotide selected from the group consisting of:

- a) a polynucleotide comprising a polynucleotide sequence selected from the group consisting of SEQ ID NO:68-134,
- b) a polynucleotide comprising a naturally occurring polynucleotide sequence at least 90% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NO:68-134,
- c) a polynucleotide complementary to a polynucleotide of a),
- d) a polynucleotide complementary to a polynucleotide of b), and
- e) an RNA equivalent of a)-d).

13. An isolated polynucleotide comprising at least 60 contiguous nucleotides of a polynucleotide of claim 12.

14. A method of detecting a target polynucleotide in a sample, said target polynucleotide having a sequence of a polynucleotide of claim 12, the method comprising:

- a) hybridizing the sample with a probe comprising at least 20 contiguous nucleotides comprising a sequence complementary to said target polynucleotide in the sample, and which probe specifically hybridizes to said target polynucleotide, under conditions whereby a hybridization complex is formed between said probe and said target polynucleotide or fragments thereof, and
- b) detecting the presence or absence of said hybridization complex, and, optionally, if present, the amount thereof.

15. A method of claim 14, wherein the probe comprises at least 60 contiguous nucleotides.

16. A method of detecting a target polynucleotide in a sample, said target polynucleotide having a sequence of a polynucleotide of claim 12, the method comprising:

- a) amplifying said target polynucleotide or fragment thereof using polymerase chain reaction amplification, and
- b) detecting the presence or absence of said amplified target polynucleotide or fragment

thereof, and, optionally, if present, the amount thereof.

17. A composition comprising a polypeptide of claim 1 and a pharmaceutically acceptable excipient.

5

18. A composition of claim 17, wherein the polypeptide has an amino acid sequence selected from the group consisting of SEQ ID NO:1-67.

19. A method for treating a disease or condition associated with decreased expression of functional SECP, comprising administering to a patient in need of such treatment the composition of claim 17.

20. A method of screening a compound for effectiveness as an agonist of a polypeptide of claim 1, the method comprising:

- 15
- a) exposing a sample comprising a polypeptide of claim 1 to a compound, and
 - b) detecting agonist activity in the sample.

21. A composition comprising an agonist compound identified by a method of claim 20 and a pharmaceutically acceptable excipient.

20

22. A method for treating a disease or condition associated with decreased expression of functional SECP, comprising administering to a patient in need of such treatment a composition of claim 21.

23. A method of screening a compound for effectiveness as an antagonist of a polypeptide of claim 1, the method comprising:

- 25
- a) exposing a sample comprising a polypeptide of claim 1 to a compound, and
 - b) detecting antagonist activity in the sample.

24. A composition comprising an antagonist compound identified by a method of claim 23 and a pharmaceutically acceptable excipient.

30

25. A method for treating a disease or condition associated with overexpression of functional SECP, comprising administering to a patient in need of such treatment a composition of claim 24.

35

26. A method of screening for a compound that specifically binds to the polypeptide of claim 1, the method comprising:

- a) combining the polypeptide of claim 1 with at least one test compound under suitable conditions, and
- 5 b) detecting binding of the polypeptide of claim 1 to the test compound, thereby identifying a compound that specifically binds to the polypeptide of claim 1.

27. A method of screening for a compound that modulates the activity of the polypeptide of claim 1, the method comprising:

- 10 a) combining the polypeptide of claim 1 with at least one test compound under conditions permissive for the activity of the polypeptide of claim 1,
- b) assessing the activity of the polypeptide of claim 1 in the presence of the test compound, and
- 15 c) comparing the activity of the polypeptide of claim 1 in the presence of the test compound with the activity of the polypeptide of claim 1 in the absence of the test compound, wherein a change in the activity of the polypeptide of claim 1 in the presence of the test compound is indicative of a compound that modulates the activity of the polypeptide of claim 1.

28. A method of screening a compound for effectiveness in altering expression of a target polynucleotide, wherein said target polynucleotide comprises a sequence of claim 5, the method comprising:

- a) exposing a sample comprising the target polynucleotide to a compound, under conditions suitable for the expression of the target polynucleotide,
- 25 b) detecting altered expression of the target polynucleotide, and
- c) comparing the expression of the target polynucleotide in the presence of varying amounts of the compound and in the absence of the compound.

29. A method of assessing toxicity of a test compound, the method comprising:

- 30 a) treating a biological sample containing nucleic acids with the test compound,
- b) hybridizing the nucleic acids of the treated biological sample with a probe comprising at least 20 contiguous nucleotides of a polynucleotide of claim 12 under conditions whereby a specific hybridization complex is formed between said probe and a target polynucleotide in the biological sample, said target polynucleotide comprising a
- 35 polynucleotide sequence of a polynucleotide of claim 12 or fragment thereof,

- c) quantifying the amount of hybridization complex, and
- d) comparing the amount of hybridization complex in the treated biological sample with the amount of hybridization complex in an untreated biological sample, wherein a difference in the amount of hybridization complex in the treated biological sample is indicative of toxicity of the test compound.

30. A diagnostic test for a condition or disease associated with the expression of SECP in a biological sample, the method comprising:

- a) combining the biological sample with an antibody of claim 11, under conditions suitable for the antibody to bind the polypeptide and form an antibody:polypeptide complex, and
- b) detecting the complex, wherein the presence of the complex correlates with the presence of the polypeptide in the biological sample.

31. The antibody of claim 11, wherein the antibody is:

- a) a chimeric antibody,
- b) a single chain antibody,
- c) a Fab fragment,
- d) a $F(ab')_2$ fragment, or
- e) a humanized antibody.

32. A composition comprising an antibody of claim 11 and an acceptable excipient.

33. A method of diagnosing a condition or disease associated with the expression of SECP in a subject, comprising administering to said subject an effective amount of the composition of claim 32.

34. A composition of claim 32, wherein the antibody is labeled.

35. A method of diagnosing a condition or disease associated with the expression of SECP in a subject, comprising administering to said subject an effective amount of the composition of claim 34.

36. A method of preparing a polyclonal antibody with the specificity of the antibody of claim 11, the method comprising:

- a) immunizing an animal with a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-67, or an immunogenic fragment thereof, under conditions to elicit an antibody response,
- b) isolating antibodies from said animal, and
- 5 c) screening the isolated antibodies with the polypeptide, thereby identifying a polyclonal antibody which binds specifically to a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-67.

37. A polyclonal antibody produced by a method of claim 36.

38. A composition comprising the polyclonal antibody of claim 37 and a suitable carrier.

39. A method of making a monoclonal antibody with the specificity of the antibody of claim 11, the method comprising:

- 15 a) immunizing an animal with a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-67, or an immunogenic fragment thereof, under conditions to elicit an antibody response,
- b) isolating antibody producing cells from the animal,
- c) fusing the antibody producing cells with immortalized cells to form monoclonal antibody-producing hybridoma cells,
- 20 d) culturing the hybridoma cells, and
- e) isolating from the culture monoclonal antibody which binds specifically to a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-67.

40. A monoclonal antibody produced by a method of claim 39.

41. A composition comprising the monoclonal antibody of claim 40 and a suitable carrier.

42. The antibody of claim 11, wherein the antibody is produced by screening a Fab expression library.

43. The antibody of claim 11, wherein the antibody is produced by screening a recombinant immunoglobulin library.

44. A method of detecting a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-67 in a sample, the method comprising:

- a) incubating the antibody of claim 11 with a sample under conditions to allow specific binding of the antibody and the polypeptide, and
- 5 b) detecting specific binding, wherein specific binding indicates the presence of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-67 in the sample.

45. A method of purifying a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-67 from a sample, the method comprising:

- a) incubating the antibody of claim 11 with a sample under conditions to allow specific binding of the antibody and the polypeptide, and
- b) separating the antibody from the sample and obtaining the purified polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-15 67.

46. A microarray wherein at least one element of the microarray is a polynucleotide of claim 13.

20 47. A method of generating a transcript image of a sample which contains polynucleotides, the method comprising:

- a) labeling the polynucleotides of the sample,
- b) contacting the elements of the microarray of claim 46 with the labeled polynucleotides of the sample under conditions suitable for the formation of a hybridization complex, and
- 25 c) quantifying the expression of the polynucleotides in the sample.

48. An array comprising different nucleotide molecules affixed in distinct physical locations on a solid substrate, wherein at least one of said nucleotide molecules comprises a first oligonucleotide or polynucleotide sequence specifically hybridizable with at least 30 contiguous nucleotides of a target polynucleotide, and wherein said target polynucleotide is a polynucleotide of claim 12.

49. An array of claim 48, wherein said first oligonucleotide or polynucleotide sequence is completely complementary to at least 30 contiguous nucleotides of said target polynucleotide.

50. An array of claim 48, wherein said first oligonucleotide or polynucleotide sequence is completely complementary to at least 60 contiguous nucleotides of said target polynucleotide.

51. An array of claim 48, wherein said first oligonucleotide or polynucleotide sequence is
5 completely complementary to said target polynucleotide.

52. An array of claim 48, which is a microarray.

53. An array of claim 48, further comprising said target polynucleotide hybridized to a
10 nucleotide molecule comprising said first oligonucleotide or polynucleotide sequence.

54. An array of claim 48, wherein a linker joins at least one of said nucleotide molecules to said solid substrate.

15 55. An array of claim 48, wherein each distinct physical location on the substrate contains multiple nucleotide molecules, and the multiple nucleotide molecules at any single distinct physical location have the same sequence, and each distinct physical location on the substrate contains nucleotide molecules having a sequence which differs from the sequence of nucleotide molecules at another distinct physical location on the substrate.

20

56. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:1.

57. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:2.

25 58. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:3.

59. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:4.

60. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:5.

30

61. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:6.

62. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:7.

35 63. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:8.

64. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:9.
65. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:10.
- 5 66. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:11.
67. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:12.
68. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:13.
- 10 69. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:14.
70. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:15.
- 15 71. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:16.
72. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:17.
73. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:18.
- 20 74. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:19.
75. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:20.
- 25 76. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:21.
77. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:22.
78. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:23.
- 30 79. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:24.
80. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:25.
- 35 81. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:26.

82. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:27.
83. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:28.
- 5 84. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:29.
85. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:30.
86. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:31.
- 10 87. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:32.
88. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:33.
- 15 89. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:34.
90. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:35.
91. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:36.
- 20 92. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:37.
93. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:38.
- 25 94. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:39.
95. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:40.
96. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:41.
- 30 97. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:42.
98. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:43.
- 35 99. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:44.

100. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:45.
101. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:46.
- 5 102. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:47.
103. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:48.
104. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:49.
- 10 105. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:50.
106. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:51.
- 15 107. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:52.
108. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:53.
109. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:54.
- 20 110. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:55.
111. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:56.
- 25 112. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:57.
113. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:58.
114. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:59.
- 30 115. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:60.
116. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:61.
- 35 117. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:62.

118. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:63.
119. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:64.
- 5 120. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:65.
121. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:66.
122. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:67.
- 10 123. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID NO:68.
124. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID NO:69.
- 15 125. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID NO:70.
126. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID NO:71.
- 20 127. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID NO:72.
128. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID NO:73.
- 25 129. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID NO:74.
- 30 130. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID NO:75.
131. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID
- 35

NO:76.

132. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID
NO:77.

5

133. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID
NO:78.

134. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID
10 NO:79.

135. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID
NO:80.

15 136. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID
NO:81.

137. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID
NO:82.

20

138. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID
NO:83.

139. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID
25 NO:84.

140. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID
NO:85.

30 141. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID
NO:86.

142. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID
NO:87.

35

143. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID
NO:88.

144. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID
5 NO:89.

145. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID
NO:90.

10 146. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID
NO:91.

147. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID
NO:92.

15 148. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID
NO:93.

149. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID
20 NO:94.

150. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID
NO:95.

25 151. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID
NO:96.

152. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID
NO:97.

30 153. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID
NO:98.

154. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID
35 NO:99.

155. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID
NO:100.

156. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID
5 NO:101.

157. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID
NO:102.

158. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID
10 NO:103.

159. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID
NO:104.
15

160. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID
NO:105.

161. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID
20 NO:106.

162. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID
NO:107.

163. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID
25 NO:108.

164. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID
NO:109.
30

165. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID
NO:110.

166. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID
35 NO:111.

167. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID
NO:112.

168. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID
5 NO:113.

169. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID
NO:114.

10 170. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID
NO:115.

171. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID
NO:116.

15 172. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID
NO:117.

173. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID
20 NO:118.

174. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID
NO:119.

25 175. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID
NO:120.

176. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID
NO:121.

30 177. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID
NO:122.

178. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID
35 NO:123.

179. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID
NO:124.

180. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID
5 NO:125.

181. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID
NO:126.

10 182. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID
NO:127.

183. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID
NO:128.

15 184. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID
NO:129.

185. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID
20 NO:130.

186. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID
NO:131.

25 187. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID
NO:132.

188. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID
NO:133.

30 189. A polynucleotide of claim 12, comprising the polynucleotide sequence of SEQ ID
NO:134.

<110> INCYTE GENOMICS, INC.

YUE, Henry
 TANG, Y. Tom
 NGUYEN, Danniel B.
 YAO, Monique G.
 XU, Yuming
 TRIBOULEY, Catherine M.
 SANJANWALA, Madhu S.
 WALIA, Narinder K.
 BAUGHN, Mariah R.
 SAPPERSTEIN, Stephanie
 LAL, Preeti
 THORNTON, Michael
 GANDHI, Ameena R.
 RAMKUMAR, Jayalaxmi
 ELLIOTT, Vicki S.
 ARVIZU, Chandra
 THANGAVELU, Kavitha
 GIETZEN, Kimberly
 DING, Li
 AU-YOUNG, Janice
 TRAN, Bao
 POLICKY, Jennifer L.
 LEE, Sally
 LU, Dyung Aina M.
 BURFORD, Neil
 WARREN, Bridget A.
 GURURAJAN, Rajagopal
 DUGGAN, Brendan M.
 HONCHELL, Cynthia D.
 HAFALIA, April J.A.

<120> SECRETED PROTEINS

<130> PI-0240 PCT

<140> To Be Assigned

<141> Herewith

<150> 60/236,869; 60/239,812; 60/240,108; 60/241,282; 60/242,218

<151> 2000-09-29; 2000-10-11; 2000-10-12; 2000-10-17; 2000-10-20

<160> 136

<170> PERL Program

<210> 1

<211> 622

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 3211795CD1

<400> 1

Met	Ala	Ala	Ala	Pro	Arg	Ala	Gly	Arg	Arg	Arg	Gly	Gln	Pro	Leu
1				5					10					15
Leu	Ala	Leu	Leu	Leu	Leu	Leu	Ala	Pro	Leu	Pro	Pro	Gly	Ala	
				20					25					30
Pro	Pro	Gly	Ala	Asp	Ala	Tyr	Phe	Pro	Glu	Glu	Arg	Trp	Ser	Pro
				35					40					45
Glu	Ser	Pro	Leu	Gln	Ala	Pro	Arg	Val	Leu	Ile	Ala	Leu	Leu	Ala

Arg Asn Ala Ala His	50	55	60
Ala Leu Pro Thr Thr	65	70	75
Leu Gly Ala Leu Glu	80	85	90
Arg Leu Arg His Pro Arg Glu Arg Thr Ala Leu Trp Val Ala Thr	95	100	105
Asp His Asn Met Asp Asn Thr Ser Thr Val Leu Arg Glu Trp Leu	110	115	120
Val Ala Val Lys Ser Leu Tyr His Ser Val Glu Trp Arg Pro Ala	125	130	135
Glu Glu Pro Arg Ser Tyr Pro Asp Glu Glu Gly Pro Lys His Trp	140	145	150
Ser Asp Ser Arg Tyr Glu His Val Met Lys Leu Arg Gln Ala Ala	155	160	165
Leu Lys Ser Ala Arg Asp Met Trp Ala Asp Tyr Ile Leu Phe Val	170	175	180
Asp Ala Asp Asn Leu Ile Leu Asn Pro Asp Thr Leu Ser Leu Leu	185	190	195
Ile Ala Glu Asn Lys Thr Val Val Ala Pro Met Leu Asp Ser Arg	200	205	210
Ala Ala Tyr Ser Asn Phe Trp Cys Gly Met Thr Ser Gln Gly Tyr	215	220	225
Tyr Lys Arg Thr Pro Ala Tyr Ile Pro Ile Arg Lys Arg Asp Arg	230	235	240
Arg Gly Cys Phe Ala Val Pro Met Val His Ser Thr Phe Leu Ile	245	250	255
Asp Leu Arg Lys Ala Ala Ser Arg Asn Leu Ala Phe Tyr Pro Pro	260	265	270
His Pro Asp Tyr Thr Trp Ser Phe Asp Asp Ile Ile Val Phe Ala	275	280	285
Phe Ser Cys Lys Gln Ala Glu Val Gln Met Tyr Val Cys Asn Lys	290	295	300
Glu Glu Tyr Gly Phe Leu Pro Val Pro Leu Arg Ala His Ser Thr	305	310	315
Leu Gln Asp Glu Ala Glu Ser Phe Met His Val Gln Leu Glu Val	320	325	330
Met Val Lys His Pro Pro Ala Glu Pro Ser Arg Phe Ile Ser Ala	335	340	345
Pro Thr Lys Thr Pro Asp Lys Met Gly Phe Asp Glu Val Phe Met	350	355	360
Ile Asn Leu Arg Arg Arg Gln Asp Arg Arg Glu Arg Met Leu Arg	365	370	375
Ala Leu Gln Ala Gln Glu Ile Glu Cys Arg Leu Val Glu Ala Val	380	385	390
Asp Gly Lys Ala Met Asn Thr Ser Gln Val Glu Ala Leu Gly Ile	395	400	405
Gln Met Leu Pro Gly Tyr Arg Asp Pro Tyr His Gly Arg Pro Leu	410	415	420
Thr Lys Gly Glu Leu Gly Cys Phe Leu Ser His Tyr Asn Ile Trp	425	430	435
Lys Glu Val Val Asp Arg Gly Leu Gln Lys Ser Leu Val Phe Glu	440	445	450
Asp Asp Leu Arg Phe Glu Ile Phe Phe Lys Arg Arg Leu Met Asn	455	460	465
Leu Met Arg Asp Val Glu Arg Glu Gly Leu Asp Trp Asp Leu Ile	470	475	480
Tyr Val Gly Arg Lys Arg Met Gln Val Glu His Pro Glu Lys Ala	485	490	495
Val Pro Arg Val Arg Asn Leu Val Glu Ala Asp Tyr Ser Tyr Trp	500	505	510
Thr Leu Ala Tyr Val Ile Ser Leu Gln Gly Ala Arg Lys Leu Leu	515	520	525
Ala Ala Glu Pro Leu Ser Lys Met Leu Pro Val Asp Glu Phe Leu			

Pro	Val	Met	Phe	Asp	Lys	His	Pro	Val	Ser	Glu	Tyr	Lys	Ala	His
				530					535					540
Phe	Ser	Leu	Arg	Asn	Leu	His	Ala	Phe	Ser	Val	Glu	Pro	Leu	Leu
				545					550					555
Ile	Tyr	Pro	Thr	His	Tyr	Thr	Gly	Asp	Asp	Gly	Tyr	Val	Ser	Asp
				560					565					570
Thr	Glu	Thr	Ser	Val	Val	Trp	Asn	Asn	Glu	His	Val	Lys	Thr	Asp
				575					580					585
Trp	Asp	Arg	Ala	Lys	Ser	Gln	Lys	Met	Arg	Glu	Gln	Gln	Ala	Leu
				590					595					600
Ser	Arg	Glu	Ala	Lys	Asn	Ser	Asp	Val	Leu	Gln	Ser	Pro	Leu	Asp
				605					610					615
Ser	Ala	Ala	Arg	Asp	Glu	Leu								
				620										

<210> 2

<211> 529

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 6813464CD1

<400> 2

Met	Ala	Gly	Ala	Pro	Pro	Pro	Ala	Leu	Leu	Leu	Pro	Cys	Ser	Leu
1				5					10					15
Ile	Ser	Asp	Cys	Cys	Ala	Ser	Asn	Gln	Arg	His	Ser	Val	Gly	Val
				20					25					30
Gly	Pro	Ser	Glu	Leu	Val	Lys	Lys	Gln	Ile	Glu	Leu	Lys	Ser	Arg
				35					40					45
Gly	Val	Lys	Leu	Met	Pro	Ser	Lys	Asp	Asn	Ser	Gln	Lys	Thr	Ser
				50					55					60
Val	Leu	Thr	Gln	Val	Gly	Val	Ser	Gln	Gly	His	Asn	Met	Cys	Pro
				65					70					75
Asp	Pro	Gly	Ile	Pro	Glu	Arg	Gly	Lys	Arg	Leu	Gly	Ser	Asp	Phe
				80					85					90
Arg	Leu	Gly	Ser	Ser	Val	Gln	Phe	Thr	Cys	Asn	Glu	Gly	Tyr	Asp
				95					100					105
Leu	Gln	Gly	Ser	Lys	Arg	Ile	Thr	Cys	Met	Lys	Val	Ser	Asp	Met
				110					115					120
Phe	Ala	Ala	Trp	Ser	Asp	His	Arg	Pro	Val	Cys	Arg	Ala	Arg	Met
				125					130					135
Cys	Asp	Ala	His	Leu	Arg	Gly	Pro	Ser	Gly	Ile	Ile	Thr	Ser	Pro
				140					145					150
Asn	Phe	Pro	Ile	Gln	Tyr	Asp	Asn	Asn	Ala	His	Cys	Val	Trp	Ile
				155					160					165
Ile	Thr	Ala	Leu	Asn	Pro	Ser	Lys	Val	Ile	Lys	Leu	Ala	Phe	Glu
				170					175					180
Glu	Phe	Asp	Leu	Glu	Arg	Gly	Tyr	Asp	Thr	Leu	Thr	Val	Gly	Asp
				185					190					195
Gly	Gly	Gln	Asp	Gly	Asp	Gln	Lys	Thr	Val	Leu	Tyr	Ile	Leu	Thr
				200					205					210
Gly	Thr	Ser	Val	Pro	Asp	Leu	Ile	Val	Ser	Thr	Asn	His	Gln	Met
				215					220					225
Trp	Leu	Leu	Phe	Glu	Thr	Asp	Gly	Ser	Gly	Ser	Ser	Leu	Gly	Phe
				230					235					240
Lys	Ala	Ser	Tyr	Glu	Glu	Ile	Glu	Gln	Gly	Ser	Cys	Gly	Asp	Pro
				245					250					255
Gly	Ile	Pro	Ala	Tyr	Gly	Arg	Arg	Glu	Gly	Ser	Arg	Phe	His	His
				260					265					270
Gly	Asp	Thr	Leu	Lys	Phe	Glu	Cys	Gln	Pro	Ala	Phe	Glu	Leu	Val
				275					280					285

Gly	Gln	Lys	Ala	Ile	Thr	Cys	Gln	Lys	Asn	Asn	Gln	Trp	Ser	Ala	
				290					295					300	
Lys	Lys	Pro	Gly	Cys	Val	Phe	Ser	Cys	Phe	Phe	Asn	Phe	Thr	Ser	
				305					310					315	
Pro	Ser	Gly	Val	Val	Leu	Ser	Pro	Asn	Tyr	Pro	Glu	Asp	Tyr	Gly	
				320					325					330	
Asn	His	Leu	His	Cys	Val	Trp	Leu	Ile	Leu	Ala	Arg	Pro	Glu	Ser	
				335					340					345	
Arg	Ile	His	Leu	Ala	Phe	Asn	Asp	Ile	Asp	Val	Glu	Pro	Gln	Phe	
				350					355					360	
Asp	Phe	Leu	Val	Ile	Lys	Asp	Gly	Ala	Thr	Ala	Glu	Ala	Pro	Val	
				365					370					375	
Leu	Gly	Thr	Phe	Ser	Gly	Asn	Gln	Leu	Pro	Ser	Ser	Ile	Thr	Ser	
				380					385					390	
Ser	Gly	His	Val	Ala	Arg	Leu	Glu	Phe	Gln	Thr	Asp	His	Ser	Thr	
				395					400					405	
Gly	Lys	Arg	Gly	Phe	Asn	Ile	Thr	Phe	Thr	Thr	Phe	Arg	His	Asn	
				410					415					420	
Glu	Cys	Pro	Asp	Pro	Gly	Val	Pro	Val	Asn	Gly	Lys	Arg	Phe	Gly	
				425					430					435	
Asp	Ser	Leu	Gln	Leu	Gly	Ser	Ser	Ile	Ser	Phe	Leu	Cys	Asp	Glu	
				440					445					450	
Gly	Phe	Leu	Gly	Thr	Gln	Gly	Ser	Glu	Thr	Ile	Thr	Cys	Val	Leu	
				455					460					465	
Lys	Glu	Gly	Ser	Val	Val	Trp	Asn	Ser	Ala	Val	Leu	Arg	Cys	Glu	
				470					475					480	
Ala	Pro	Cys	Gly	Gly	His	Leu	Thr	Ser	Pro	Ser	Gly	Thr	Ile	Leu	
				485					490					495	
Ser	Pro	Gly	Trp	Pro	Gly	Phe	Tyr	Lys	Asp	Ala	Leu	Ser	Cys	Ala	
				500					505					510	
Trp	Val	Ile	Glu	Ala	Gln	Pro	Gly	Tyr	Pro	Ile	Lys	Ile	Thr	Phe	
				515					520					525	
Asp	Arg	Cys	Leu												

<210> 3

<211> 204

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 2156540CD1

<400> 3

Met	Gly	Ala	Pro	Leu	Ala	Val	Ala	Leu	Gly	Ala	Leu	His	Tyr	Leu	
1				5					10					15	
Ala	Leu	Phe	Leu	Gln	Leu	Gly	Gly	Ala	Thr	Arg	Pro	Ala	Gly	His	
				20					25					30	
Ala	Pro	Trp	Asp	Asn	His	Val	Ser	Gly	His	Ala	Leu	Phe	Thr	Glu	
				35					40					45	
Thr	Pro	His	Asp	Met	Thr	Ala	Arg	Thr	Gly	Glu	Asp	Val	Glu	Met	
				50					55					60	
Ala	Cys	Ser	Phe	Arg	Gly	Ser	Gly	Ser	Pro	Ser	Tyr	Ser	Leu	Glu	
				65					70					75	
Ile	Gln	Trp	Trp	Tyr	Val	Arg	Ser	His	Arg	Asp	Trp	Thr	Asp	Lys	
				80					85					90	
Gln	Ala	Trp	Ala	Ser	Asn	Gln	Leu	Lys	Ala	Ser	Gln	Gln	Glu	Asp	
				95					100					105	
Ala	Gly	Lys	Glu	Ala	Thr	Lys	Ile	Ser	Val	Val	Lys	Val	Val	Gly	
				110					115					120	
Ser	Asn	Ile	Ser	His	Lys	Leu	Arg	Leu	Ser	Arg	Val	Lys	Pro	Thr	
				125					130					135	

Asp	Glu	Gly	Thr	Tyr	Glu	Cys	Arg	Val	Ile	Asp	Phe	Ser	Asp	Gly
				140					145					150
Lys	Ala	Arg	His	His	Lys	Val	Lys	Ala	Tyr	Leu	Arg	Val	Gln	Pro
				155					160					165
Gly	Glu	Asn	Ser	Val	Leu	His	Leu	Pro	Glu	Ala	Pro	Pro	Ala	Ala
				170					175					180
Pro	Ala	Pro	Pro	Pro	Pro	Lys	Pro	Gly	Lys	Glu	Leu	Arg	Lys	Arg
				185					190					195
Ser	Val	Asp	Gln	Glu	Ala	Cys	Ser	Leu						
				200										

<210> 4

<211> 406

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 894939CD1

<400> 4

Met	Arg	Gly	Ser	Val	Glu	Cys	Thr	Trp	Gly	Trp	Gly	His	Cys	Ala
1				5					10					15
Pro	Ser	Pro	Leu	Leu	Leu	Trp	Thr	Leu	Leu	Leu	Phe	Ala	Ala	Pro
				20					25					30
Phe	Gly	Leu	Leu	Gly	Glu	Lys	Thr	Arg	Gln	Val	Ser	Leu	Glu	Val
				35					40					45
Ile	Pro	Asn	Trp	Leu	Gly	Pro	Leu	Gln	Asn	Leu	Leu	His	Ile	Arg
				50					55					60
Ala	Val	Gly	Thr	Asn	Ser	Thr	Leu	His	Tyr	Val	Trp	Ser	Ser	Leu
				65					70					75
Gly	Pro	Leu	Ala	Val	Val	Met	Val	Ala	Thr	Asn	Thr	Pro	His	Ser
				80					85					90
Thr	Leu	Ser	Val	Asn	Trp	Ser	Leu	Leu	Leu	Ser	Pro	Glu	Pro	Asp
				95					100					105
Gly	Gly	Leu	Met	Val	Leu	Pro	Lys	Asp	Ser	Ile	Gln	Phe	Ser	Ser
				110					115					120
Ala	Leu	Val	Phe	Thr	Arg	Leu	Leu	Glu	Phe	Asp	Ser	Thr	Asn	Val
				125					130					135
Ser	Asp	Thr	Ala	Ala	Lys	Pro	Leu	Gly	Arg	Pro	Tyr	Pro	Pro	Tyr
				140					145					150
Ser	Leu	Ala	Asp	Phe	Ser	Trp	Asn	Asn	Ile	Thr	Asp	Ser	Leu	Asp
				155					160					165
Pro	Ala	Thr	Leu	Ser	Ala	Thr	Phe	Gln	Gly	His	Pro	Met	Asn	Asp
				170					175					180
Pro	Thr	Arg	Thr	Phe	Ala	Asn	Gly	Ser	Leu	Ala	Phe	Arg	Val	Gln
				185					190					195
Ala	Phe	Ser	Arg	Ser	Ser	Arg	Pro	Ala	Gln	Pro	Pro	Arg	Leu	Leu
				200					205					210
His	Thr	Ala	Asp	Thr	Cys	Gln	Leu	Glu	Val	Ala	Leu	Ile	Gly	Ala
				215					220					225
Ser	Pro	Arg	Gly	Asn	Arg	Ser	Leu	Phe	Gly	Leu	Glu	Val	Ala	Thr
				230					235					240
Leu	Gly	Gln	Gly	Pro	Asp	Cys	Pro	Ser	Met	Gln	Glu	Gln	His	Ser
				245					250					255
Ile	Asp	Asp	Glu	Tyr	Ala	Pro	Ala	Val	Phe	Gln	Leu	Asp	Gln	Leu
				260					265					270
Leu	Trp	Gly	Ser	Leu	Pro	Ser	Gly	Phe	Ala	Gln	Trp	Arg	Pro	Val
				275					280					285
Ala	Tyr	Ser	Gln	Lys	Pro	Gly	Gly	Arg	Glu	Ser	Ala	Leu	Pro	Cys
				290					295					300
Gln	Ala	Ser	Pro	Leu	His	Pro	Ala	Leu	Ala	Tyr	Ser	Leu	Pro	Gln
				305					310					315

```

Ser Pro Ile Val Arg Ala Phe Phe Gly Ser Gln Asn Asn Phe Cys
      320      325      330
Ala Phe Asn Leu Thr Phe Gly Ala Ser Thr Gly Pro Gly Tyr Trp
      335      340      345
Asp Gln His Tyr Leu Ser Trp Ser Met Leu Leu Gly Val Gly Phe
      350      355      360
Pro Pro Val Asp Gly Leu Ser Pro Leu Val Leu Gly Ile Met Ala
      365      370      375
Val Ala Leu Gly Ala Pro Gly Leu Met Leu Leu Gly Gly Gly Leu
      380      385      390
Val Leu Leu Leu His His Lys Lys Tyr Ser Glu Tyr Gln Ser Ile
      395      400      405
Asn

```

```

<210> 5
<211> 477
<212> PRT
<213> Homo sapiens

```

```

<220>
<221> misc_feature
<223> Incyte ID No: 4620890CD1

```

```

<400> 5
Met Pro Gly Ser Asp Thr Ala Leu Thr Val Asp Arg Thr Tyr Ser
  1      5      10      15
Asp Pro Gly Arg His His Arg Cys Lys Ser Arg Val Glu Arg His
      20      25      30
Asp Met Asn Thr Leu Ser Leu Pro Leu Asn Ile Arg Arg Gly Gly
      35      40      45
Ser Asp Thr Asn Gly Leu Asn Phe Asp Val Pro Asp Gly Ile Leu Asp
      50      55      60
Phe His Lys Val Lys Leu Thr Ala Asp Ser Leu Lys Gln Lys Ile
      65      70      75
Leu Lys Val Thr Glu Gln Ile Lys Ile Glu Gln Thr Ser Arg Asp
      80      85      90
Gly Asn Val Ala Glu Tyr Leu Lys Leu Val Asn Asn Ala Asp Lys
      95      100     105
Gln Gln Ala Gly Arg Ile Lys Gln Val Phe Glu Lys Lys Asn Gln
      110     115     120
Lys Ser Ala His Ser Ile Ala Gln Leu Gln Lys Lys Leu Glu Gln
      125     130     135
Tyr His Arg Lys Leu Arg Glu Ile Glu Gln Asn Gly Ala Ser Arg
      140     145     150
Ser Ser Lys Asp Ile Ser Lys Asp His Leu Lys Asp Ile His Arg
      155     160     165
Ser Leu Lys Asp Ala His Val Lys Ser Arg Thr Ala Pro His Cys
      170     175     180
Met Glu Ser Ser Lys Ser Gly Met Pro Gly Val Ser Leu Thr Pro
      185     190     195
Pro Val Phe Val Phe Asn Lys Ser Arg Glu Phe Ala Asn Leu Ile
      200     205     210
Arg Asn Lys Phe Gly Ser Ala Asp Asn Ile Ala His Leu Lys Asn
      215     220     225
Ser Leu Glu Glu Phe Arg Pro Glu Ala Ser Ala Arg Ala Tyr Gly
      230     235     240
Gly Ser Ala Thr Ile Val Asn Lys Pro Lys Tyr Gly Ser Asp Asp
      245     250     255
Glu Cys Ser Ser Gly Thr Ser Gly Ser Ala Asp Ser Asn Gly Asn
      260     265     270
Gln Ser Phe Gly Ala Gly Gly Ala Ser Thr Leu Asp Ser Gln Gly
      275     280     285

```

```

Lys Leu Ala Val Ile Leu Glu Glu Leu Arg Glu Ile Lys Asp Thr
290 295 300
Gln Ala Gln Leu Ala Glu Asp Ile Glu Ala Leu Lys Val Gln Phe
305 310 315
Lys Arg Glu Tyr Gly Phe Ile Ser Gln Thr Leu Gln Glu Glu Arg
320 325 330
Tyr Arg Tyr Glu Arg Leu Glu Asp Gln Leu His Asp Leu Thr Asp
335 340 345
Leu His Gln His Glu Thr Ala Asn Leu Lys Gln Glu Leu Ala Ser
350 355 360
Ile Glu Glu Lys Val Ala Tyr Gln Ala Tyr Glu Arg Ser Arg Asp
365 370 375
Ile Gln Glu Ala Leu Glu Ser Cys Gln Thr Arg Ile Ser Lys Leu
380 385 390
Glu Leu His Gln Gln Glu Gln Gln Ala Leu Gln Thr Asp Thr Val
395 400 405
Asn Ala Lys Val Leu Leu Gly Arg Cys Ile Asn Val Ile Leu Ala
410 415 420
Phe Met Thr Val Ile Leu Val Cys Val Ser Thr Ile Ala Lys Phe
425 430 435
Val Ser Pro Met Met Lys Ser Arg Cys His Ile Leu Gly Thr Phe
440 445 450
Phe Ala Val Thr Leu Leu Ala Ile Phe Cys Lys Asn Trp Asp His
455 460 465
Ile Leu Cys Ala Ile Glu Arg Met Ile Ile Pro Arg
470 475

```

<210> 6

<211> 691

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 5514146CD1

<400> 6

```

Met Cys Phe Arg Thr Lys Leu Ser Val Ser Trp Val Pro Leu Phe
1 5 10 15
Leu Leu Leu Ser Arg Val Phe Ser Thr Glu Thr Asp Lys Pro Ser
20 25 30
Ala Gln Asp Ser Arg Ser Arg Gly Ser Ser Gly Gln Pro Ala Asp
35 40 45
Leu Leu Gln Val Leu Ser Ala Gly Asp His Pro Pro His Asn His
50 55 60
Ser Arg Ser Leu Ile Lys Thr Leu Leu Glu Lys Thr Gly Cys Pro
65 70 75
Arg Arg Arg Asn Gly Met Gln Gly Asp Cys Asn Leu Cys Phe Glu
80 85 90
Pro Asp Ala Leu Leu Leu Ile Ala Gly Gly Asn Phe Glu Asp Gln
95 100 105
Leu Arg Glu Glu Val Val Gln Arg Val Ser Leu Leu Leu Leu Tyr
110 115 120
Tyr Ile Ile His Gln Glu Glu Ile Cys Ser Ser Lys Leu Asn Met
125 130 135
Ser Asn Lys Glu Tyr Lys Phe Tyr Leu His Ser Leu Leu Ser Leu
140 145 150
Arg Gln Asp Glu Asp Ser Ser Phe Leu Ser Gln Asn Glu Thr Glu
155 160 165
Asp Ile Leu Ala Phe Thr Arg Gln Tyr Phe Asp Thr Ser Gln Ser
170 175 180
Gln Cys Met Glu Thr Lys Thr Leu Gln Lys Lys Ser Gly Ile Val
185 190 195

```

Ser	Ser	Glu	Gly	Ala	Asn	Glu	Ser	Thr	Leu	Pro	Gln	Leu	Ala	Ala
				200					205					210
Met	Ile	Ile	Thr	Leu	Ser	Leu	Gln	Gly	Val	Cys	Leu	Gly	Gln	Gly
				215					220					225
Asn	Leu	Pro	Ser	Pro	Asp	Tyr	Phe	Thr	Glu	Tyr	Ile	Phe	Ser	Ser
				230					235					240
Leu	Asn	Arg	Thr	Asn	Thr	Leu	Arg	Leu	Ser	Glu	Leu	Asp	Gln	Leu
				245					250					255
Leu	Asn	Thr	Leu	Trp	Thr	Arg	Ser	Thr	Cys	Ile	Lys	Asn	Glu	Lys
				260					265					270
Ile	His	Gln	Phe	Gln	Arg	Lys	Gln	Asn	Asn	Ile	Ile	Thr	His	Asp
				275					280					285
Gln	Asp	Tyr	Ser	Asn	Phe	Ser	Ser	Ser	Met	Glu	Lys	Glu	Ser	Glu
				290					295					300
Asp	Gly	Pro	Ile	Ser	Trp	Asp	Gln	Thr	Cys	Phe	Ser	Ala	Arg	Gln
				305					310					315
Leu	Val	Glu	Ile	Phe	Leu	Gln	Lys	Gly	Leu	Ser	Leu	Ile	Ser	Lys
				320					325					330
Glu	Asp	Phe	Lys	Gln	Met	Ser	Pro	Gly	Ile	Ile	Gln	Gln	Leu	Leu
				335					340					345
Ser	Cys	Ser	Cys	His	Leu	Pro	Lys	Asp	Gln	Gln	Ala	Lys	Leu	Pro
				350					355					360
Pro	Thr	Thr	Leu	Glu	Lys	Tyr	Gly	Tyr	Ser	Thr	Val	Ala	Val	Thr
				365					370					375
Leu	Leu	Thr	Leu	Gly	Ser	Met	Leu	Gly	Thr	Ala	Leu	Val	Leu	Phe
				380					385					390
His	Ser	Cys	Glu	Glu	Asn	Tyr	Arg	Leu	Ile	Leu	Gln	Leu	Phe	Val
				395					400					405
Gly	Leu	Ala	Val	Gly	Thr	Leu	Ser	Gly	Asp	Ala	Leu	Leu	His	Leu
				410					415					420
Ile	Pro	Gln	Val	Leu	Gly	Leu	His	Lys	Gln	Glu	Ala	Pro	Glu	Phe
				425					430					435
Gly	His	Phe	His	Glu	Ser	Lys	Gly	His	Ile	Trp	Lys	Leu	Met	Gly
				440					445					450
Leu	Ile	Gly	Gly	Ile	His	Gly	Phe	Phe	Leu	Ile	Glu	Lys	Cys	Phe
				455					460					465
Ile	Leu	Leu	Val	Ser	Pro	Asn	Asp	Lys	Gln	Gly	Leu	Ser	Leu	Val
				470					475					480
Asn	Gly	His	Val	Gly	His	Ser	His	His	Leu	Ala	Leu	Asn	Ser	Glu
				485					490					495
Leu	Ser	Asp	Gln	Ala	Gly	Arg	Gly	Lys	Ser	Ala	Ser	Thr	Ile	Gln
				500					505					510
Leu	Lys	Ser	Pro	Glu	Asp	Ser	Gln	Ala	Ala	Glu	Met	Pro	Ile	Gly
				515					520					525
Ser	Met	Thr	Ala	Ser	Asn	Arg	Lys	Cys	Lys	Ala	Ile	Ser	Leu	Leu
				530					535					540
Ala	Ile	Met	Ile	Leu	Val	Gly	Asp	Ser	Leu	His	Asn	Phe	Ala	Asp
				545					550					555
Gly	Leu	Ala	Ile	Gly	Ala	Ala	Phe	Ser	Ser	Ser	Ser	Glu	Ser	Gly
				560					565					570
Val	Thr	Thr	Thr	Ile	Ala	Ile	Leu	Cys	His	Glu	Ile	Pro	His	Glu
				575					580					585
Met	Gly	Asp	Phe	Ala	Val	Leu	Leu	Ser	Ser	Gly	Leu	Ser	Met	Lys
				590					595					600
Thr	Ala	Ile	Leu	Met	Asn	Phe	Ile	Ser	Ser	Leu	Thr	Ala	Phe	Met
				605					610					615
Gly	Leu	Tyr	Ile	Gly	Leu	Ser	Val	Ser	Ala	Asp	Pro	Cys	Val	Gln
				620					625					630
Asp	Trp	Ile	Phe	Thr	Val	Thr	Ala	Gly	Met	Phe	Leu	Tyr	Leu	Ser
				635					640					645
Leu	Val	Glu	Met	Leu	Pro	Glu	Met	Thr	His	Val	Gln	Thr	Gln	Arg
				650					655					660
Pro	Trp	Met	Met	Phe	Leu	Leu	Gln	Asn	Phe	Gly	Leu	Ile	Leu	Gly

	665		670		675
Trp	Leu	Ser	Leu	Leu	Leu
	680		685		690
Ile					

<210> 7

<211> 919

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 7474769CD1

<400> 7

Met	Gln	Glu	Cys	Leu	Thr	Leu	Trp	Val	Phe	Ser	Pro	Leu	Ala	Leu
1				5					10					15
Thr	Asp	Ser	Gly	Tyr	Thr	Lys	Thr	Tyr	Gln	Ala	His	Ala	Lys	Gln
				20					25					30
Lys	Phe	Ser	Arg	Leu	Trp	Ser	Ser	Lys	Ser	Val	Thr	Glu	Ile	His
				35					40					45
Leu	Tyr	Phe	Glu	Glu	Glu	Val	Lys	Gln	Glu	Glu	Cys	Asp	His	Leu
				50					55					60
Asp	Arg	Leu	Phe	Ala	Pro	Lys	Glu	Ala	Gly	Lys	Gln	Pro	Arg	Thr
				65					70					75
Val	Ile	Ile	Gln	Gly	Pro	Gln	Gly	Ile	Gly	Lys	Thr	Thr	Leu	Leu
				80					85					90
Met	Lys	Leu	Met	Met	Ala	Trp	Ser	Asp	Asn	Lys	Ile	Phe	Arg	Asp
				95					100					105
Arg	Phe	Leu	Tyr	Thr	Phe	Tyr	Phe	Cys	Cys	Arg	Glu	Leu	Arg	Glu
				110					115					120
Leu	Pro	Pro	Thr	Ser	Leu	Ala	Asp	Leu	Ile	Ser	Arg	Glu	Trp	Pro
				125					130					135
Asp	Pro	Ala	Ala	Pro	Ile	Thr	Glu	Ile	Val	Ser	Gln	Pro	Glu	Arg
				140					145					150
Leu	Leu	Phe	Val	Ile	Asp	Ser	Phe	Glu	Glu	Leu	Gln	Gly	Gly	Leu
				155					160					165
Asn	Glu	Pro	Asp	Ser	Asp	Leu	Cys	Gly	Asp	Leu	Met	Glu	Lys	Arg
				170					175					180
Pro	Val	Gln	Val	Leu	Leu	Ser	Ser	Leu	Leu	Arg	Lys	Lys	Met	Leu
				185					190					195
Pro	Glu	Ala	Ser	Leu	Leu	Ile	Ala	Ile	Lys	Pro	Val	Cys	Pro	Lys
				200					205					210
Glu	Leu	Arg	Asp	Gln	Val	Thr	Ile	Ser	Glu	Ile	Tyr	Gln	Pro	Arg
				215					220					225
Gly	Phe	Asn	Glu	Ser	Asp	Arg	Leu	Val	Tyr	Phe	Cys	Cys	Phe	Phe
				230					235					240
Lys	Asp	Pro	Lys	Arg	Ala	Met	Glu	Ala	Phe	Asn	Leu	Val	Arg	Glu
				245					250					255
Ser	Glu	Gln	Leu	Phe	Ser	Ile	Cys	Gln	Ile	Pro	Leu	Leu	Cys	Trp
				260					265					270
Ile	Leu	Cys	Thr	Ser	Leu	Lys	Gln	Glu	Met	Gln	Lys	Gly	Lys	Asp
				275					280					285
Leu	Ala	Leu	Thr	Cys	Gln	Ser	Thr	Thr	Ser	Val	Tyr	Ser	Ser	Phe
				290					295					300
Val	Phe	Asn	Leu	Phe	Thr	Pro	Glu	Gly	Ala	Glu	Gly	Pro	Thr	Pro
				305					310					315
Gln	Thr	Gln	His	Gln	Leu	Lys	Ala	Leu	Cys	Ser	Leu	Ala	Ala	Glu
				320					325					330
Gly	Met	Trp	Thr	Asp	Thr	Phe	Glu	Phe	Cys	Glu	Asp	Asp	Leu	Arg
				335					340					345
Arg	Asn	Gly	Val	Val	Asp	Ala	Asp	Ile	Pro	Ala	Leu	Leu	Gly	Thr

	350		355		360
Lys Ile Leu Leu	Lys Tyr Gly Glu Arg	Glu Ser Ser Tyr Val	Phe		
	365		370		375
Leu His Val Cys	Ile Gln Glu Phe Cys	Ala Ala Leu Phe Tyr	Leu		
	380		385		390
Leu Lys Ser His	Leu Asp His Pro His	Pro Ala Val Arg Cys	Val		
	395		400		405
Gln Glu Leu Leu	Val Ala Asn Phe Glu	Lys Ala Arg Arg Ala	His		
	410		415		420
Trp Ile Phe Leu	Gly Cys Phe Leu Thr	Gly Leu Leu Asn Lys	Lys		
	425		430		435
Glu Gln Glu Lys	Leu Asp Ala Phe Phe	Gly Phe Gln Leu Ser	Gln		
	440		445		450
Glu Ile Lys Gln	Gln Ile His Gln Cys	Leu Lys Ser Leu Gly	Glu		
	455		460		465
Arg Gly Asn Pro	Gln Gly Gln Val Asp	Ser Leu Ala Ile Phe	Tyr		
	470		475		480
Cys Leu Phe Glu	Met Gln Asp Pro Ala	Phe Val Lys Gln Ala	Val		
	485		490		495
Asn Leu Leu Gln	Glu Ala Asn Phe His	Ile Ile Asp Asn Val	Asp		
	500		505		510
Leu Val Val Ser	Ala Tyr Cys Leu Lys	Tyr Cys Ser Ser Leu	Arg		
	515		520		525
Lys Leu Cys Phe	Ser Val Gln Asn Val	Phe Lys Lys Glu Asp	Glu		
	530		535		540
His Ser Ser Thr	Ser Asp Tyr Ser Leu	Ile Cys Trp His His	Ile		
	545		550		555
Cys Ser Val Leu	Thr Thr Ser Gly His	Leu Arg Glu Leu Gln	Val		
	560		565		570
Gln Asp Ser Thr	Leu Ser Glu Ser Thr	Phe Val Thr Trp Cys	Asn		
	575		580		585
Gln Leu Arg His	Pro Ser Cys Arg Leu	Gln Lys Leu Gly Ile	Asn		
	590		595		600
Asn Val Ser Phe	Ser Gly Gln Ser Val	Leu Leu Phe Glu Val	Leu		
	605		610		615
Phe Tyr Gln Pro	Asp Leu Lys Tyr Leu	Ser Phe Thr Leu Thr	Lys		
	620		625		630
Leu Ser Arg Asp	Asp Ile Arg Ser Leu	Cys Asp Ala Leu Asn	Tyr		
	635		640		645
Pro Ala Gly Asn	Val Lys Glu Leu Ala	Leu Val Asn Cys His	Leu		
	650		655		660
Ser Pro Ile Asp	Cys Glu Val Leu Ala	Gly Leu Leu Thr Asn	Asn		
	665		670		675
Lys Lys Leu Thr	Tyr Leu Asn Val Ser	Cys Asn Gln Leu Asp	Thr		
	680		685		690
Gly Val Pro Leu	Leu Cys Glu Ala Leu	Cys Ser Pro Asp Thr	Val		
	695		700		705
Leu Val Tyr Leu	Met Leu Ala Phe Cys	His Leu Ser Glu Gln	Cys		
	710		715		720
Cys Glu Tyr Ile	Ser Glu Met Leu Leu	Arg Asn Lys Ser Val	Arg		
	725		730		735
Tyr Leu Asp Leu	Ser Ala Asn Val Leu	Lys Asp Glu Gly Leu	Lys		
	740		745		750
Thr Leu Cys Glu	Ala Leu Lys His Pro	Asp Cys Cys Leu Asp	Ser		
	755		760		765
Leu Cys Leu Val	Lys Cys Phe Ile Thr	Ala Ala Gly Cys Glu	Asp		
	770		775		780
Leu Ala Ser Ala	Leu Ile Ser Asn Gln	Asn Leu Lys Ile Leu	Gln		
	785		790		795
Ile Gly Cys Asn	Glu Ile Gly Asp Val	Gly Val Gln Leu Leu	Cys		
	800		805		810
Arg Ala Leu Thr	His Thr Asp Cys Arg	Leu Glu Ile Leu Gly	Leu		
	815		820		825

Glu	Glu	Cys	Gly	Leu	Thr	Ser	Thr	Cys	Cys	Lys	Asp	Leu	Ala	Ser	
				830					835					840	
Val	Leu	Thr	Cys	Ser	Lys	Thr	Leu	Gln	Gln	Leu	Asn	Leu	Thr	Leu	
				845					850					855	
Asn	Thr	Leu	Asp	His	Thr	Gly	Val	Val	Val	Leu	Cys	Glu	Ala	Leu	
				860					865					870	
Arg	His	Pro	Glu	Cys	Ala	Leu	Gln	Val	Leu	Gly	Leu	Arg	Lys	Thr	
				875					880					885	
Asp	Phe	Asp	Glu	Glu	Thr	Gln	Ala	Leu	Leu	Thr	Ala	Glu	Glu	Glu	
				890					895					900	
Arg	Asn	Pro	Asn	Leu	Thr	Ile	Thr	Asp	Asp	Cys	Asp	Thr	Ile	Thr	
				905					910					915	
Arg	Val	Glu	Ile												

<210> 8
 <211> 178
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 065296CD1

<400>	8														
Met	Ala	Gly	Lys	Pro	Tyr	Ser	Leu	Arg	Gly	Ser	Ser	His	Thr	Thr	
1				5					10					15	
Gly	Thr	Phe	Leu	Leu	Leu	Ser	Gln	Ser	Ser	Gly	Glu	Leu	Gln	Ile	
				20					25					30	
Ile	Lys	Tyr	Phe	Lys	Met	Lys	Phe	Lys	Thr	Glu	Met	Phe	Leu	Leu	
				35					40					45	
Leu	Leu	Leu	Trp	Arg	Asp	Cys	Met	Lys	Thr	His	Thr	Gly	Met	Asn	
				50					55					60	
His	Arg	Leu	His	Val	Pro	Glu	Leu	Ser	Asn	Ala	Gln	Asp	Asn	Asn	
				65					70					75	
Ser	Ser	Ala	Ser	Ile	Ser	Asp	Lys	Val	Gly	Phe	Ser	Lys	Ala	Glu	
				80					85					90	
Leu	Arg	Met	Cys	Leu	Ala	Ile	Trp	Thr	Phe	Ser	Pro	Ile	Lys	Gln	
				95					100					105	
Val	Tyr	Lys	Ile	Leu	Lys	Ile	Glu	Cys	Leu	Asn	Phe	Ser	Ile	Val	
				110					115					120	
Leu	Ser	Val	Leu	Lys	Pro	Ile	Arg	Ile	Pro	Arg	Ile	Asn	Met	Phe	
				125					130					135	
Val	Phe	Leu	Gly	Ala	Leu	Ser	Met	Thr	Gln	Asp	Asn	Glu	Trp	Tyr	
				140					145					150	
Leu	Asn	Tyr	Ile	Phe	Phe	Thr	Leu	Glu	Ile	Ser	Arg	Gln	Lys	Val	
				155					160					165	
Phe	Phe	Glu	Trp	Val	Asn	Ser	Ala	Leu	Ser	Phe	Ser	Gln			
				170					175						

<210> 9
 <211> 310
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 231994CD1

<400>	9														
Met	Ser	Cys	Pro	Val	Gln	Thr	Met	Asp	Pro	Glu	Val	Thr	Leu	Leu	
1				5					10					15	
Leu	Gln	Cys	Pro	Gly	Gly	Gly	Leu	Pro	Gln	Glu	Gln	Ile	Gln	Ala	

	20		25		30
Glu Leu Ser Pro Ala His Asp Arg Arg Pro Leu Pro Gly Gly Asp					
	35		40		45
Glu Ala Ile Thr Ala Ile Trp Glu Thr Arg Leu Lys Ala Gln Pro					
	50		55		60
Trp Leu Phe Asp Ala Pro Lys Phe Arg Leu His Ser Ala Thr Leu					
	65		70		75
Ala Pro Ile Gly Ser Arg Gly Pro Gln Leu Leu Leu Arg Leu Gly					
	80		85		90
Leu Thr Ser Tyr Arg Asp Phe Leu Gly Thr Asn Trp Ser Ser Ser					
	95		100		105
Ala Ala Trp Leu Arg Gln Gln Gly Ala Thr Asp Trp Gly Asp Thr					
	110		115		120
Gln Ala Tyr Leu Ala Asp Pro Leu Gly Val Gly Ala Ala Leu Ala					
	125		130		135
Thr Ala Asp Asp Phe Leu Val Phe Leu Arg Arg Ser Arg Gln Val					
	140		145		150
Ala Glu Ala Pro Gly Leu Val Asp Val Pro Gly Gly His Pro Glu					
	155		160		165
Pro Gln Ala Leu Cys Pro Gly Gly Ser Pro Gln His Gln Asp Leu					
	170		175		180
Ala Gly Gln Leu Val Val His Glu Leu Phe Ser Ser Val Leu Gln					
	185		190		195
Glu Ile Cys Asp Glu Val Asn Leu Pro Leu Leu Thr Leu Ser Gln					
	200		205		210
Pro Leu Leu Leu Gly Ile Ala Arg Asn Glu Thr Ser Ala Gly Arg					
	215		220		225
Ala Ser Ala Glu Phe Tyr Val Gln Cys Ser Leu Thr Ser Glu Gln					
	230		235		240
Val Arg Lys His Tyr Leu Ser Gly Gly Pro Glu Ala His Glu Ser					
	245		250		255
Thr Gly Ile Phe Phe Val Glu Thr Gln Asn Val Arg Arg Leu Pro					
	260		265		270
Glu Thr Glu Met Trp Ala Glu Leu Cys Pro Ser Ala Lys Gly Ala					
	275		280		285
Ile Ile Leu Tyr Asn Arg Val Gln Gly Ser Pro Thr Gly Ala Ala					
	290		295		300
Leu Gly Ser Pro Ala Leu Leu Pro Pro Leu					
	305		310		

<210> 10

<211> 559

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 538054CD1

<400> 10

Met Trp Thr Asn Phe Phe Lys Leu Arg Leu Phe Cys Cys Leu Leu					
1	5		10		15
Ala Val Leu Met Val Val Val Leu Val Ile Asn Val Thr Gln Val					
	20		25		30
Glu Tyr Leu Asp His Glu Thr Val Ser Ala Thr Phe Ile Asp Ser					
	35		40		45
Ser Gly Gln Phe Val Ser Ser Gln Val Thr Gly Ile Ser Arg Asn					
	50		55		60
Pro Tyr Cys Gly Tyr Asp Gln Gln Thr Leu Ser Ser Gln Glu Arg					
	65		70		75
Met Glu Glu Asp Ser Leu Leu Ala Ala Leu His Arg Gln Val Pro					
	80		85		90
Asp Val Gly Pro Val Pro Phe Val Lys Ser Thr Asp Pro Ser Ser					

	95	100	105
Ser Tyr Phe Val	Ile Leu Asn Ser Ala	Ala Phe Phe Lys Val Gly	
	110	115	120
Ser Gln Leu Glu	Val Leu Val His Val	Gln Asp Phe Gln Arg Lys	
	125	130	135
Pro Lys Lys Tyr	Gly Gly Asp Tyr Leu	Gln Ala Arg Ile His Ser	
	140	145	150
Leu Lys Leu Gln	Ala Gly Ala Val Gly	Arg Val Val Asp Tyr Gln	
	155	160	165
Asn Gly Phe Tyr	Lys Val Phe Phe Thr	Leu Leu Trp Pro Gly Lys	
	170	175	180
Val Lys Val Ser	Val Ser Leu Val His	Pro Ser Glu Gly Ile Arg	
	185	190	195
Val Leu Gln Arg	Leu Gln Glu Asp Lys	Pro Asp Arg Val Tyr Phe	
	200	205	210
Lys Ser Leu Phe	Arg Ser Gly Arg Ile	Ser Glu Thr Thr Glu Cys	
	215	220	225
Asn Val Cys Leu	Pro Gly Asn Leu Pro	Leu Cys Asn Phe Thr Asp	
	230	235	240
Leu Tyr Thr Gly	Glu Pro Trp Phe Cys	Phe Lys Pro Lys Lys Leu	
	245	250	255
Pro Cys Ser Ser	Arg Ile Thr His Phe	Lys Gly Gly Tyr Leu Lys	
	260	265	270
Gly Leu Leu Thr	Ala Ala Glu Ser Ala	Phe Phe Gln Ser Gly Val	
	275	280	285
Asn Ile Lys Met	Pro Val Asn Ser Ser	Gly Pro Asp Trp Val Thr	
	290	295	300
Val Ile Pro Arg	Arg Ile Lys Glu Thr	Asn Ser Leu Glu Leu Ser	
	305	310	315
Gln Gly Ser Gly	Thr Phe Pro Ser Gly	Tyr Tyr Tyr Lys Asp Gln	
	320	325	330
Trp Arg Pro Arg	Lys Phe Lys Met Arg	Gln Phe Asn Asp Pro Asp	
	335	340	345
Asn Ile Thr Glu	Cys Leu Gln Arg Lys	Val Val His Leu Phe Gly	
	350	355	360
Asp Ser Thr Ile	Arg Gln Trp Phe Glu	Tyr Leu Thr Thr Phe Val	
	365	370	375
Pro Asp Leu Val	Glu Phe Asn Leu Gly	Ser Pro Lys Asn Val Gly	
	380	385	390
Pro Phe Leu Ala	Val Asp Gln Lys His	Asn Ile Leu Leu Lys Tyr	
	395	400	405
Arg Cys His Gly	Pro Pro Ile Arg Phe	Thr Thr Val Phe Ser Asn	
	410	415	420
Glu Leu His Tyr	Val Ala Asn Glu Leu	Asn Gly Ile Val Gly Gly	
	425	430	435
Lys Asn Thr Val	Val Ala Ile Ala Val	Trp Ser His Phe Ser Thr	
	440	445	450
Phe Pro Leu Glu	Val Tyr Ile Arg Arg	Leu Arg Asn Ile Arg Arg	
	455	460	465
Ala Val Val Arg	Leu Leu Asp Arg Ser	Pro Lys Thr Val Val Val	
	470	475	480
Ile Arg Thr Ala	Asn Ala Gln Glu Leu	Gly Pro Glu Val Ser Leu	
	485	490	495
Phe Asn Ser Asp	Trp Tyr Asn Phe Gln	Leu Asp Thr Ile Leu Arg	
	500	505	510
Arg Met Phe Ser	Gly Val Gly Val Tyr	Leu Val Asp Ala Trp Glu	
	515	520	525
Met Thr Leu Ala	His Tyr Leu Pro His	Lys Leu His Pro Asp Glu	
	530	535	540
Val Ile Val Lys	Asn Gln Leu Asp Met	Phe Leu Ser Phe Val Cys	
	545	550	555
Pro Leu Glu Thr			

```
<220>  
<221> misc_feature  
<223> Incyte ID No: 1259305CD1
```

Met	Val	Cys	Val	Phe	Val	Met	Asn	Arg	Met	Asn	Ser	Gln	Asn	Ser
1				5					10					15
Gly	Phe	Thr	Gln	Arg	Arg	Arg	Met	Ala	Leu	Gly	Ile	Val	Ile	Leu
				20					25					30
Leu	Leu	Val	Asp	Val	Ile	Trp	Val	Ala	Ser	Ser	Glu	Leu	Thr	Ser
				35					40					45
Tyr	Val	Phe	Thr	Gln	Tyr	Asn	Lys	Pro	Phe	Phe	Ser	Thr	Phe	Ala
				50					55					60
Lys	Thr	Ser	Met	Phe	Val	Leu	Tyr	Leu	Leu	Gly	Phe	Ile	Ile	Trp
				65					70					75
Lys	Pro	Trp	Arg	Gln	Gln	Cys	Thr	Arg	Gly	Leu	Arg	Gly	Lys	His
				80					85					90
Ala	Ala	Phe	Phe	Ala	Asp	Ala	Glu	Gly	Tyr	Phe	Ala	Ala	Cys	Thr
				95					100					105
Thr	Asp	Thr	Thr	Met	Asn	Ser	Ser	Leu	Ser	Glu	Pro	Leu	Tyr	Val
				110					115					120
Pro	Val	Lys	Phe	His	Asp	Leu	Pro	Ser	Glu	Lys	Pro	Glu	Ser	Thr
				125					130					135
Asn	Ile	Asp	Thr	Glu	Lys	Thr	Pro	Lys	Lys	Ser	Arg	Val	Arg	Phe
				140					145					150
Ser	Asn	Ile	Met	Glu	Ile	Arg	Gln	Leu	Pro	Ser	Ser	His	Ala	Leu
				155					160					165
Glu	Ala	Lys	Leu	Ser	Arg	Met	Ser	Tyr	Pro	Val	Lys	Glu	Gln	Glu
				170					175					180
Ser	Ile	Leu	Lys	Thr	Val	Gly	Lys	Leu	Thr	Ala	Thr	Gln	Val	Ala
				185					190					195
Lys	Ile	Ser	Phe	Phe	Phe	Cys	Phe	Val	Trp	Phe	Leu	Ala	Asn	Leu
				200					205					210
Ser	Tyr	Gln	Glu	Ala	Leu	Ser	Asp	Thr	Gln	Val	Ala	Ile	Val	Asn
				215					220					225
Ile	Leu	Ser	Ser	Thr	Ser	Gly	Leu	Phe	Thr	Leu	Ile	Leu	Ala	Ala
				230					235					240
Val	Phe	Pro	Ser	Asn	Ser	Gly	Asp	Arg	Phe	Thr	Leu	Ser	Lys	Leu
				245					250					255
Leu	Ala	Val	Ile	Leu	Ser	Ile	Gly	Gly	Val	Val	Leu	Val	Asn	Leu
				260					265					270
Ala	Gly	Ser	Glu	Lys	Pro	Ala	Gly	Arg	Asp	Thr	Val	Gly	Ser	Ile
				275					280					285
Trp	Ser	Leu	Ala	Gly	Ala	Met	Leu	Tyr	Ala	Val	Tyr	Ile	Val	Met
				290					295					300
Ile	Lys	Arg	Lys	Val	Asp	Arg	Glu	Asp	Lys	Leu	Asp	Ile	Pro	Met
				305					310					315
Phe	Phe	Gly	Phe	Val	Gly	Leu	Phe	Asn	Leu	Leu	Leu	Leu	Trp	Pro
				320					325					330
Gly	Phe	Phe	Leu	Leu	His	Tyr	Thr	Gly	Phe	Glu	Asp	Phe	Glu	Phe
				335					340					345
Pro	Asn	Lys	Val	Val	Leu	Met	Cys	Ile	Ile	Ile	Asn	Gly	Leu	Ile

395	400	405
Trp Leu Phe Phe	Ala Gly Ala Ile Pro Val Phe Phe Ser Phe Phe	
410	415	420
Ile Val Thr Leu	Leu Cys His Tyr Asn Asn Trp Asp Pro Val Met	
425	430	435
Val Gly Ile Arg	Arg Ile Phe Ala Phe Ile Cys Arg Lys His Arg	
440	445	450
Ile Gln Arg Val	Pro Glu Asp Ser Glu Gln Cys Glu Ser Leu Ile	
455	460	465
Ser Met His Ser	Val Ser Gln Glu Asp Gly Ala Ser	
470	475	

<210> 12
 <211> 176
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 1483702CD1

<400> 12
Met Leu Ser Leu Lys Leu Pro Gln Leu Leu Gln Val His Gln Val
1 5 10 15
Pro Arg Val Phe Trp Glu Asp Gly Ile Met Ser Gly Tyr Arg Arg
20 25 30
Pro Thr Ser Ser Ala Leu Asp Cys Val Leu Ser Ser Phe Gln Met
35 40 45
Thr Asn Glu Thr Val Asn Ile Trp Thr His Phe Leu Pro Thr Trp
50 55 60
Tyr Phe Leu Trp Arg Leu Leu Ala Leu Ala Gly Gly Pro Gly Phe
65 70 75
Arg Ala Glu Pro Tyr His Trp Pro Leu Leu Val Phe Leu Leu Pro
80 85 90
Ala Cys Leu Tyr Pro Phe Ala Ser Cys Cys Ala His Thr Phe Ser
95 100 105
Ser Met Ser Pro Arg Met Arg His Ile Cys Tyr Phe Leu Asp Tyr
110 115 120
Gly Ala Leu Ser Leu Tyr Ser Leu Val Ser Trp Ser Trp Lys Ala
125 130 135
Leu Gly Ser Val Arg Ser Ser Ala Gln Glu Pro Ser Pro Ile His
140 145 150
Ser Cys Ser Thr Thr Ser His Ser Phe Ile Gly Ser Gly Cys Ala
155 160 165
Gly Ala Gly Ala Thr Ala Val Gly Arg Arg Pro
170 175

<210> 13
 <211> 190
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 1519324CD1

<400> 13
Met Ala Gly Met Met Lys Gly Ile Arg Trp Ser Cys Pro Ala Ile
1 5 10 15
Ala Ser Ile Ser Gln Thr Arg Ser Ser Gln Glu Lys Asp Ser Ser
20 25 30
Ser Pro Pro Trp Asp Leu Arg Arg Ala Ala Thr Glu Trp Gly Gly
35 40 45

```

Pro Arg Cys Ala Val Pro Lys Pro Gly Pro Arg Pro Lys Phe Ser
      50      55      60
Leu Pro Ser Leu Val Pro Ser Cys Pro Phe Leu Leu His Ala Trp
      65      70      75
Ala Cys Arg Pro Thr Pro Ala Thr Thr Glu Ser Thr Arg Ser Ala
      80      85      90
Leu Cys Ser Trp Arg Arg His Ser Arg Val Glu Ser Cys Pro Ser
      95     100     105
Leu Ser Leu Gly His Leu Gly Gly Glu Ser Gly Leu Arg Ser Glu
     110     115     120
Leu Asp Pro Gly Asp Leu Gly Ser Phe Phe Leu Ala His Gln Pro
     125     130     135
Cys Arg Pro His Leu Ser Gln Asn Pro Leu Cys Leu Gly Gly Ser
     140     145     150
Gly Ser Ala Leu Leu Cys Ser Arg Gly Trp Gly Val Asp Ser Ile
     155     160     165
Arg Trp Glu Ser Gly Val His Pro His Val Ser Val Gly Phe Ser
     170     175     180
Pro Trp Gly Trp Lys Lys Arg Ala Ser Thr
     185     190

```

<210> 14
 <211> 75
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 1630169CD1

```

<400> 14
Met Ala Leu Gly Lys Val Leu Ala Met Ala Leu Val Leu Ala Leu
      1      5      10      15
Ala Val Leu Gly Ser Leu Ser Pro Gly Ala Arg Ala Gly Asp Cys
      20      25      30
Lys Gly Gln Arg Gln Val Leu Arg Glu Ala Pro Gly Phe Val Thr
      35      40      45
Asp Gly Ala Gly Asn Tyr Ser Val Asn Gly Asn Cys Glu Trp Leu
      50      55      60
Ile Glu Gly Glu Trp Gly Arg Val Gly His Ser Leu Ile Arg Trp
      65      70      75

```

<210> 15
 <211> 265
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 1664253CD1

```

<400> 15
Met Ser Ala Ala Pro Leu Val Gly Tyr Ser Ser Ser Gly Ser Glu
      1      5      10      15
Asp Glu Ser Glu Asp Gly Met Arg Thr Arg Pro Gly Asp Gly Ser
      20      25      30
His Arg Arg Gly Gln Ser Pro Leu Pro Arg Gln Arg Phe Pro Val
      35      40      45
Pro Asp Ser Val Leu Asn Met Phe Pro Gly Thr Glu Glu Gly Pro
      50      55      60
Glu Asp Asp Ser Thr Lys His Gly Gly Arg Val Arg Thr Phe Pro
      65      70      75

```

```

His Glu Arg Gly Asn Trp Ala Thr His Val Tyr Val Pro Tyr Glu
      80                      85                      90
Ala Lys Glu Glu Phe Leu Asp Leu Leu Asp Val Leu Leu Pro His
      95                      100                     105
Ala Gln Thr Tyr Val Pro Arg Leu Val Arg Met Lys Val Phe His
      110                     115                     120
Leu Ser Leu Ser Gln Ser Val Val Leu Arg His His Trp Ile Leu
      125                     130                     135
Pro Phe Val Gln Ala Leu Lys Ala Arg Met Thr Ser Phe His Arg
      140                     145                     150
Phe Phe Phe Thr Ala Asn Gln Val Lys Ile Tyr Thr Asn Gln Glu
      155                     160                     165
Lys Thr Arg Thr Phe Ile Gly Leu Glu Val Thr Ser Gly His Ala
      170                     175                     180
Gln Phe Leu Asp Leu Val Ser Glu Val Asp Arg Val Met Glu Glu
      185                     190                     195
Phe Asn Leu Thr Thr Phe Tyr Gln Asp Pro Ser Phe His Leu Ser
      200                     205                     210
Leu Ala Trp Cys Val Gly Asp Ala Arg Leu Gln Leu Glu Gly Gln
      215                     220                     225
Cys Leu Gln Glu Leu Gln Ala Ile Val Asp Gly Phe Glu Asp Ala
      230                     235                     240
Glu Val Leu Leu Arg Val His Thr Glu Gln Val Arg Cys Lys Ser
      245                     250                     255
Gly Asn Lys Phe Phe Ser Met Pro Leu Lys
      260                     265

```

<210> 16
 <211> 202
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 1864715CD1

```

<400> 16
Met Glu Ala Ala Leu Leu Gly Leu Cys Asn Trp Ser Thr Leu Gly
  1          5          10          15
Val Cys Ala Ala Leu Lys Leu Pro Gln Ile Ser Ala Val Leu Ala
  20          25          30
Ala Arg Ser Ala Arg Gly Leu Ser Leu Pro Ser Leu Leu Leu Glu
  35          40          45
Leu Ala Gly Phe Leu Val Phe Leu Arg Tyr Gln Cys Tyr Tyr Gly
  50          55          60
Tyr Pro Pro Leu Thr Tyr Leu Glu Tyr Pro Ile Leu Ile Ala Gln
  65          70          75
Asp Val Ile Leu Leu Leu Cys Ile Phe His Phe Asn Gly Asn Val
  80          85          90
Lys Gln Ala Thr Pro Tyr Ile Ala Val Leu Val Ser Ser Trp Phe
  95          100         105
Ile Leu Ala Leu Gln Lys Trp Ile Ile Asp Leu Ala Met Asn Leu
  110         115         120
Cys Thr Phe Ile Ser Ala Ala Ser Lys Phe Ala Gln Leu Gln Cys
  125         130         135
Leu Trp Lys Thr Arg Asp Ser Gly Thr Val Ser Ala Leu Thr Trp
  140         145         150
Ser Leu Ser Ser Tyr Thr Cys Ala Thr Arg Ile Ile Thr Thr Leu
  155         160         165
Met Thr Thr Asn Asp Phe Thr Ile Leu Leu Arg Phe Val Ile Met
  170         175         180
Leu Ala Leu Asn Ile Trp Val Thr Val Thr Val Leu Arg Tyr Arg
  185         190         195

```


Lys Thr Ala Ile Lys Ala Glu
200

<210> 17
<211> 111
<212> PRT
<213> Homo sapiens

<220>
<221> misc_feature
<223> Incyte ID No: 1929395CD1

<400> 17
Met Tyr Pro Thr Ala Pro Glu Leu Leu Val Pro Gln Pro Arg Pro
1 5 10 15
Gln Gly Ser Pro Ala Ser Leu Leu Leu Gly Thr Pro Val Leu Ala
20 25 30
Ala Val Tyr Gly Ala Ser Cys Leu Pro Leu Gly Arg His Pro Cys
35 40 45
Thr Pro Ala Ser Phe Pro Trp Pro Phe Leu Ala Pro Val Leu Leu
50 55 60
Leu Tyr Ile Asp Leu Phe Thr Gln Lys Arg Ala Arg Pro Leu Phe
65 70 75
Ser Ala Thr Ser Pro Val Ser Glu Ile Gln Pro Pro Arg Leu His
80 85 90
Arg Lys Ile Asp Ile Leu Glu Ile Met Lys Ser Asp Ile Phe Ala
95 100 105
Tyr Glu Arg Lys Lys Gly
110

<210> 18
<211> 105
<212> PRT
<213> Homo sapiens

<220>
<221> misc_feature
<223> Incyte ID No: 1987737CD1

<400> 18
Met Cys Val His Arg Cys Glu Cys Val Cys Met Arg Ala Cys Leu
1 5 10 15
Cys Ala Gly Val Cys Met Cys Val Ala Ser Cys Leu Gly Leu Pro
20 25 30
Met Asn Val Val Glu Cys Tyr Thr Trp Arg Val Leu Val Phe His
35 40 45
Gln Phe Gln Asp Glu Glu Leu His Asp Thr Val Asp Leu Glu Thr
50 55 60
Ile Pro Leu Glu Arg Gln Pro Arg Asp Val Gln His Pro Val Ser
65 70 75
Thr Arg Ile Leu Tyr Leu His Val Tyr Phe Val Ala Val Thr Leu
80 85 90
Thr Leu Ile Arg Ile Leu Gln Leu Trp Thr Glu Ala Phe Ser Pro
95 100 105

<210> 19
<211> 717
<212> PRT
<213> Homo sapiens

<220>
<221> misc_feature

<223> Incyte ID No: 2122866CD1

<400> 19

Met	Ala	Ser	Ser	Ser	Asp	Ser	Glu	Asp	Asp	Ser	Phe	Met	Ala	Val
1				5					10					15
Asp	Gln	Glu	Glu	Thr	Val	Leu	Glu	Gly	Thr	Met	Asp	Gln	Asp	Glu
				20					25					30
Glu	Pro	His	Pro	Val	Leu	Glu	Ala	Glu	Thr	Arg	His	Asn	Arg	
				35					40					45
Ser	Met	Ser	Glu	Leu	Pro	Glu	Glu	Val	Leu	Glu	Tyr	Ile	Leu	Ser
				50					55					60
Phe	Leu	Ser	Pro	Tyr	Gln	Glu	His	Lys	Thr	Ala	Ala	Leu	Val	Cys
				65					70					75
Lys	Gln	Trp	Tyr	Arg	Leu	Ile	Lys	Gly	Val	Ala	His	Gln	Cys	Tyr
				80					85					90
His	Gly	Phe	Met	Lys	Ala	Val	Gln	Glu	Gly	Asn	Ile	Gln	Trp	Glu
				95					100					105
Ser	Arg	Thr	Tyr	Pro	Tyr	Pro	Gly	Thr	Pro	Ile	Thr	Gln	Arg	Phe
				110					115					120
Ser	His	Ser	Ala	Cys	Tyr	Tyr	Asp	Ala	Asn	Gln	Ser	Met	Tyr	Val
				125					130					135
Phe	Gly	Gly	Cys	Thr	Gln	Ser	Ser	Cys	Asn	Ala	Ala	Phe	Asn	Asp
				140					145					150
Leu	Trp	Arg	Leu	Asp	Leu	Asn	Ser	Lys	Glu	Trp	Ile	Arg	Pro	Leu
				155					160					165
Ala	Ser	Gly	Ser	Tyr	Pro	Ser	Pro	Lys	Ala	Gly	Ala	Thr	Leu	Val
				170					175					180
Val	Tyr	Lys	Asp	Leu	Leu	Val	Leu	Phe	Gly	Gly	Trp	Thr	Arg	Pro
				185					190					195
Ser	Pro	Tyr	Pro	Leu	His	Gln	Pro	Glu	Arg	Phe	Phe	Asp	Glu	Ile
				200					205					210
His	Thr	Tyr	Ser	Pro	Ser	Lys	Asn	Trp	Trp	Asn	Cys	Ile	Val	Thr
				215					220					225
Thr	His	Gly	Pro	Pro	Pro	Met	Ala	Gly	His	Ser	Ser	Cys	Val	Ile
				230					235					240
Asp	Asp	Lys	Met	Ile	Val	Phe	Gly	Gly	Ser	Leu	Gly	Ser	Arg	Gln
				245					250					255
Met	Ser	Asn	Asp	Val	Trp	Val	Leu	Asp	Leu	Glu	Gln	Trp	Ala	Trp
				260					265					270
Ser	Lys	Pro	Asn	Ile	Ser	Gly	Pro	Ser	Pro	His	Pro	Arg	Gly	Gly
				275					280					285
Gln	Ser	Gln	Ile	Val	Ile	Asp	Asp	Ala	Thr	Ile	Leu	Ile	Leu	Gly
				290					295					300
Gly	Cys	Gly	Gly	Pro	Asn	Ala	Leu	Phe	Lys	Asp	Ala	Trp	Leu	Leu
				305					310					315
His	Met	His	Ser	Gly	Pro	Trp	Ala	Trp	Gln	Pro	Leu	Lys	Val	Glu
				320					325					330
Asn	Glu	Glu	His	Gly	Ala	Pro	Glu	Leu	Trp	Cys	His	Pro	Ala	Cys
				335					340					345
Arg	Val	Gly	Gln	Cys	Val	Val	Val	Phe	Ser	Gln	Ala	Pro	Ser	Gly
				350					355					360
Arg	Ala	Pro	Leu	Ser	Pro	Ser	Leu	Asn	Ser	Arg	Pro	Ser	Pro	Ile
				365					370					375
Ser	Ala	Thr	Pro	Pro	Ala	Leu	Val	Pro	Glu	Thr	Arg	Glu	Tyr	Arg
				380					385					390
Ser	Gln	Ser	Pro	Val	Arg	Ser	Met	Asp	Glu	Ala	Pro	Cys	Val	Asn
				395					400					405
Gly	Arg	Trp	Gly	Thr	Leu	Arg	Pro	Arg	Ala	Gln	Arg	Gln	Thr	Pro
				410					415					420
Ser	Gly	Ser	Arg	Glu	Gly	Ser	Leu	Ser	Pro	Ala	Arg	Gly	Asp	Gly
				425					430					435
Ser	Pro	Ile	Leu	Asn	Gly	Gly	Ser	Leu	Ser	Pro	Gly	Thr	Ala	Ala
				440					445					450

```

Val Gly Gly Ser Ser Leu Asp Ser Pro Val Gln Ala Ile Ser Pro
455 460 465
Ser Thr Pro Ser Ala Pro Glu Gly Tyr Asp Leu Lys Ile Gly Leu
470 475 480
Ser Leu Ala Pro Arg Arg Gly Ser Leu Pro Asp Gln Lys Asp Leu
485 490 495
Arg Leu Gly Ser Ile Asp Leu Asn Trp Asp Leu Lys Pro Ala Ser
500 505 510
Ser Ser Asn Pro Met Asp Gly Met Asp Asn Arg Thr Val Gly Gly
515 520 525
Ser Met Arg His Pro Pro Glu Gln Thr Asn Gly Val His Thr Pro
530 535 540
Pro His Val Ala Ser Ala Leu Ala Gly Ala Val Ser Pro Gly Ala
545 550 555
Leu Arg Arg Ser Leu Glu Ala Ile Lys Ala Met Ser Ser Lys Gly
560 565 570
Pro Ser Ala Ser Ala Ala Leu Ser Pro Pro Leu Gly Ser Ser Pro
575 580 585
Gly Ser Pro Gly Ser Gln Ser Leu Ser Ser Gly Glu Thr Val Pro
590 595 600
Ile Pro Arg Pro Gly Pro Ala Gln Gly Asp Gly His Ser Leu Pro
605 610 615
Pro Ile Ala Arg Arg Leu Gly His His Pro Pro Gln Ser Leu Asn
620 625 630
Val Gly Lys Pro Leu Tyr Gln Ser Met Asn Cys Lys Pro Met Gln
635 640 645
Met Tyr Val Leu Asp Ile Lys Asp Thr Lys Glu Lys Gly Arg Val
650 655 660
Lys Trp Lys Val Phe Asn Ser Ser Ser Val Val Gly Pro Pro Glu
665 670 675
Thr Ser Leu His Thr Val Val Gln Gly Arg Gly Glu Leu Ile Ile
680 685 690
Phe Gly Gly Leu Met Asp Lys Lys Gln Asn Val Lys Tyr Tyr Pro
695 700 705
Lys Thr Asn Ala Leu Tyr Phe Val Arg Ala Lys Arg
710 715

```

<210> 20

<211> 580

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 2123981CD1

<400> 20

```

Met His Leu Ser Ala Val Phe Asn Ala Leu Leu Val Ser Val Leu
1 5 10 15
Ala Ala Val Leu Trp Lys His Val Arg Leu Arg Glu His Ala Ala
20 25 30
Thr Leu Glu Glu Glu Leu Ala Leu Ser Arg Gln Ala Thr Glu Pro
35 40 45
Ala Pro Ala Leu Arg Ile Asp Tyr Pro Lys Ala Leu Gln Ile Leu
50 55 60
Met Glu Gly Gly Thr His Met Val Cys Thr Gly Arg Thr His Thr
65 70 75
Asp Arg Ile Cys Arg Phe Lys Trp Leu Cys Tyr Ser Asn Glu Ala
80 85 90
Glu Glu Phe Ile Phe Phe His Gly Asn Thr Ser Val Met Leu Pro
95 100 105
Asn Leu Gly Ser Arg Arg Phe Gln Pro Ala Leu Leu Asp Leu Ser
110 115 120

```

Thr Val Glu Asp	His Asn Thr Gln Tyr Phe Asn Phe Val Glu Leu	125	130	135
Pro Ala Ala Ala	Leu Arg Phe Met Pro Lys Pro Val Phe Val Pro	140	145	150
Asp Val Ala Leu	Ile Ala Asn Arg Phe Asn Pro Asp Asn Leu Met	155	160	165
His Val Phe His	Asp Asp Leu Leu Pro Leu Phe Tyr Thr Leu Arg	170	175	180
Gln Phe Pro Gly	Leu Ala His Glu Ala Arg Leu Phe Phe Met Glu	185	190	195
Gly Trp Gly Glu	Gly Ala His Phe Asp Leu Tyr Lys Leu Leu Ser	200	205	210
Pro Lys Gln Pro	Leu Leu Arg Ala Gln Leu Lys Thr Leu Gly Arg	215	220	225
Leu Leu Cys Phe	Ser His Ala Phe Val Gly Leu Ser Lys Ile Thr	230	235	240
Thr Trp Tyr Gln	Tyr Gly Phe Val Gln Pro Gln Gly Pro Lys Ala	245	250	255
Asn Ile Leu Val	Ser Gly Asn Glu Ile Arg Gln Phe Ala Arg Phe	260	265	270
Met Thr Glu Lys	Leu Asn Val Ser His Thr Gly Val Pro Leu Gly	275	280	285
Glu Glu Tyr Ile	Leu Val Phe Ser Arg Thr Gln Asn Arg Leu Ile	290	295	300
Leu Asn Glu Ala	Glu Leu Leu Leu Ala Leu Ala Gln Glu Phe Gln	305	310	315
Met Lys Thr Val	Thr Val Ser Leu Glu Asp His Thr Phe Ala Asp	320	325	330
Val Val Arg Leu	Val Ser Asn Ala Ser Met Leu Val Ser Met His	335	340	345
Gly Ala Gln Leu	Val Thr Thr Leu Phe Leu Pro Arg Gly Ala Thr	350	355	360
Val Val Glu Leu	Phe Pro Tyr Ala Val Asn Pro Asp His Tyr Thr	365	370	375
Pro Tyr Lys Thr	Leu Ala Met Leu Pro Gly Met Asp Leu Gln Tyr	380	385	390
Val Ala Trp Arg	Asn Met Met Pro Glu Asn Thr Val Thr His Pro	395	400	405
Glu Arg Pro Trp	Asp Gln Gly Gly Ile Thr His Leu Asp Arg Ala	410	415	420
Glu Gln Ala Arg	Ile Leu Gln Ser Arg Glu Val Pro Arg His Leu	425	430	435
Cys Cys Arg Asn	Pro Glu Trp Leu Phe Arg Ile Tyr Gln Asp Thr	440	445	450
Lys Val Asp Ile	Pro Ser Leu Ile Gln Thr Ile Arg Arg Val Val	455	460	465
Lys Gly Arg Pro	Gly Pro Arg Lys Gln Lys Trp Thr Val Gly Leu	470	475	480
Tyr Pro Gly Lys	Val Arg Glu Ala Arg Cys Gln Ala Ser Val His	485	490	495
Gly Ala Ser Glu	Ala Arg Leu Thr Val Ser Trp Gln Ile Pro Trp	500	505	510
Asn Leu Lys Tyr	Leu Lys Val Arg Glu Val Lys Tyr Glu Val Trp	515	520	525
Leu Gln Glu Gln	Gly Glu Asn Thr Tyr Val Pro Tyr Ile Leu Ala	530	535	540
Leu Gln Asn His	Thr Phe Thr Glu Asn Ile Lys Pro Phe Thr Thr	545	550	555
Tyr Leu Val Trp	Val Arg Cys Ile Phe Asn Lys Ile Leu Leu Gly	560	565	570
Pro Phe Ala Asp	Val Leu Val Cys Asn Thr	575	580	

<210> 21
 <211> 172
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 2200177CD1

<400> 21
 Met Leu Ser Glu Asp Lys Leu Phe Gln Ile Ile His Ser Leu Leu
 1 5 10 15
 Ile Thr Gln Leu Ala Ser Glu Thr Lys Ile Ser Ala Thr Ile Cys
 20 25 30
 Leu Pro Leu Leu Phe His Cys Leu Phe Leu Leu Val Leu Ser Phe
 35 40 45
 Pro Ile Thr Leu Cys Ile Arg His Ser Gly Pro Tyr His Ile Tyr
 50 55 60
 Pro Leu Leu Gln Val Ser Asn Leu Ile Phe Leu Gln Thr His Phe
 65 70 75
 Leu Ser Tyr Ile Ala Gly Ile Met Gln Lys Leu Leu Ser Asn Val
 80 85 90
 Val His Ser Gln Lys Ile His Pro Glu Ile Leu Arg Phe Gly Lys
 95 100 105
 Val Cys Ala Gln Ser Thr Ile Ser Lys Lys Phe Lys Glu Glu Lys
 110 115 120
 Tyr Lys Thr Pro His Thr Ile Ser Leu Ile Ser Gln Ile His Glu
 125 130 135
 Thr Ala Thr Ile Lys Ser Lys Val Phe Arg Lys Leu Ser Thr Tyr
 140 145 150
 Phe Ser Ile Val Leu Lys Leu Lys Glu Ile Lys Ile Ala Gly Phe
 155 160 165
 Lys Tyr Leu Trp Ser Ser Asn
 170

<210> 22
 <211> 256
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 2319255CD1

<400> 22
 Met Asn Thr Phe Lys Ala Val Gly Lys Ile Arg Gly Lys Pro Leu
 1 5 10 15
 Pro Leu Leu Leu Phe Phe Glu Ala Leu Phe Ile Thr Ser His Ala
 20 25 30
 Phe Pro Cys Pro Val Asp Ala Ala Leu Thr Leu Glu Gly Ile Lys
 35 40 45
 Cys Gly Leu Ser Glu Lys Arg Leu Asp Leu Val Thr Asn Trp Val
 50 55 60
 Thr Gln Glu Arg Leu Thr Phe Ser Glu Glu Ala Gly Asp Val Ile
 65 70 75
 Cys Asp Tyr Gly Glu Gln Asp Thr Tyr Asn Lys Ala Lys Cys Leu
 80 85 90
 Ala Leu Ala Gln Ile Ile Tyr Ser Glu Cys Gly Leu His Lys Lys
 95 100 105
 Ala Ile Leu Cys Leu Cys Lys Gln Gly Gln Thr His Arg Val Met
 110 115 120
 Glu Tyr Ile Gln Gln Leu Lys Asp Phe Thr Thr Asp Asp Leu Leu
 125 130 135

Gln Leu Leu Met Ser Cys Pro Gln Val Glu Leu Ile Gln Cys Leu
 140 145 150
 Thr Lys Glu Leu Asn Glu Lys Gln Pro Ser Leu Ser Phe Gly Leu
 155 160 165
 Ala Ile Leu His Leu Phe Ser Ala Asp Met Lys Lys Val Gly Ile
 170 175 180
 Lys Leu Leu Gln Glu Ile Asn Lys Gly Gly Ile Asp Ala Val Glu
 185 190 195
 Ser Leu Met Ile Asn Asp Ser Phe Cys Ser Ile Glu Lys Trp Gln
 200 205 210
 Glu Val Ala Asn Ile Cys Ser Gln Asn Gly Phe Asp Lys Leu Ser
 215 220 225
 Asn Asp Ile Thr Ser Ile Leu Arg Ser Gln Ala Ala Val Thr Glu
 230 235 240
 Ile Ser Glu Glu Asp Asp Ala Val Asn Leu Met Glu His Val Phe
 245 250 255
 Trp

<210> 23
 <211> 93
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 2792452CD1

<400> 23
 Met Ser Pro Met Trp Ala Pro Pro Trp Leu Pro Leu Leu Leu Ser
 1 5 10 15
 Lys Ser Glu Pro Thr Gln Ser Pro Ser Pro Arg Arg Pro Leu Pro
 20 25 30
 Pro Gly Lys Met Thr Leu Gly Gln Gly Ser Leu Leu Met Ser Val
 35 40 45
 Phe Cys Leu Val Gly Leu Gly Val Pro Leu Pro Leu Ile Arg Arg
 50 55 60
 Gly Phe Arg Ala Glu Ile Lys Pro Gln Thr Gly Glu Pro Leu Trp
 65 70 75
 His Met Ala Pro Arg Ala Ser His Ala Ser Gly Phe Ser Pro Cys
 80 85 90
 Gln Asp Thr

<210> 24
 <211> 112
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 2853088CD1

<400> 24
 Met Pro Thr Asp Val Pro Lys Ala Arg Leu Glu Leu Thr Ser Leu
 1 5 10 15
 Leu Leu Leu Leu Phe Leu Arg Trp Ser Leu Ala Leu Leu Pro
 20 25 30
 Arg Leu Asp Cys Ser Gly Ala Val Leu Ala His Cys Asn Phe Arg
 35 40 45
 Leu Trp Gly Ser Ser Asp Ser Ser Ala Ser Ala Ser Ser Gln Val
 50 55 60
 Ala Gly Ser Thr Gly Ala Cys His Gln Ala Arg Ala Lys Glu Arg

	65		70		75
Asp Ser Ile Ser Lys Ile Ile Thr Ile Ile Ile Met Arg Ser Ile					
	80		85		90
Pro Asp Val Leu Leu Gly Arg Leu Trp Ala Tyr Ser Leu Glu Leu					
	95		100		105
Arg Arg Asp Ile Lys Ala Ser					
	110				

<210> 25
 <211> 186
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 2949004CD1

<400> 25	
Met Lys Leu Pro Arg Glu Glu Ala Ser Ala Ser Phe Val Arg Arg	
1 5 10 15	
Ala Asp Leu Thr Arg Glu Asp Leu Ala Pro Ser Ser Val Asp Ser	
20 25 30	
Gly Gln Ala Gly Phe Gly Gly Cys Cys Glu Ser Gly Leu Pro Asn	
35 40 45	
Thr Met Pro Ser Ala Phe Ser Val Ser Ser Phe Pro Val Ser Ile	
50 55 60	
Pro Ala Val Leu Thr Gln Thr Asp Trp Thr Glu Pro Trp Leu Met	
65 70 75	
Gly Leu Ala Thr Phe His Ala Leu Cys Val Leu Leu Thr Cys Leu	
80 85 90	
Ser Ser Arg Ser Tyr Arg Leu Gln Ile Gly His Phe Leu Cys Leu	
95 100 105	
Val Ile Leu Val Tyr Cys Ala Glu Tyr Ile Asn Glu Ala Ala Ala	
110 115 120	
Met Asn Trp Arg Leu Phe Ser Lys Tyr Gln Tyr Phe Asp Ser Arg	
125 130 135	
Gly Met Phe Ile Ser Ile Val Phe Ser Ala Pro Leu Leu Val Asn	
140 145 150	
Ala Met Ile Ile Val Val Met Trp Val Trp Lys Thr Leu Asn Val	
155 160 165	
Met Thr Asp Leu Lys Asn Ala Gln Glu Arg Arg Lys Glu Lys Lys	
170 175 180	
Arg Arg Arg Lys Glu Asp	
185	

<210> 26
 <211> 487
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 3011670CD1

<400> 26	
Met Gly Val Ile Gly Ile Gln Leu Val Val Thr Met Val Met Ala	
1 5 10 15	
Ser Val Met Gln Lys Ile Ile Pro His Tyr Ser Leu Ala Arg Trp	
20 25 30	
Leu Leu Cys Asn Gly Ser Leu Arg Trp Tyr Gln His Pro Thr Glu	
35 40 45	
Glu Glu Leu Arg Ile Leu Ala Gly Lys Gln Gln Lys Gly Lys Thr	
50 55 60	

Lys	Lys	Asp	Arg	Lys	Tyr	Asn	Gly	His	Ile	Glu	Ser	Lys	Pro	Leu	
				65					70					75	
Thr	Ile	Pro	Lys	Asp	Ile	Asp	Leu	His	Leu	Glu	Thr	Lys	Ser	Val	
				80					85					90	
Thr	Glu	Val	Asp	Thr	Leu	Ala	Leu	His	Tyr	Phe	Pro	Glu	Tyr	Gln	
				95					100					105	
Trp	Leu	Val	Asp	Phe	Thr	Val	Ala	Ala	Thr	Val	Val	Tyr	Leu	Val	
				110					115					120	
Thr	Glu	Val	Tyr	Tyr	Asn	Phe	Met	Lys	Pro	Thr	Gln	Glu	Met	Asn	
				125					130					135	
Ile	Ser	Leu	Val	Trp	Cys	Leu	Leu	Val	Leu	Ser	Phe	Ala	Ile	Lys	
				140					145					150	
Val	Leu	Phe	Ser	Leu	Thr	Thr	His	Tyr	Phe	Lys	Val	Glu	Asp	Gly	
				155					160					165	
Gly	Glu	Arg	Ser	Val	Cys	Val	Thr	Phe	Gly	Phe	Phe	Phe	Phe	Val	
				170					175					180	
Lys	Ala	Met	Ala	Val	Leu	Ile	Val	Thr	Glu	Asn	Tyr	Leu	Glu	Phe	
				185					190					195	
Gly	Leu	Glu	Thr	Gly	Phe	Thr	Asn	Phe	Ser	Asp	Ser	Ala	Met	Gln	
				200					205					210	
Phe	Leu	Glu	Lys	Gln	Gly	Leu	Glu	Ser	Gln	Ser	Pro	Val	Ser	Lys	
				215					220					225	
Leu	Thr	Phe	Lys	Phe	Phe	Leu	Ala	Ile	Phe	Cys	Ser	Phe	Ile	Gly	
				230					235					240	
Ala	Phe	Leu	Thr	Phe	Pro	Gly	Leu	Arg	Leu	Ala	Gln	Met	His	Leu	
				245					250					255	
Asp	Ala	Leu	Asn	Leu	Ala	Thr	Glu	Lys	Ile	Thr	Gln	Thr	Leu	Leu	
				260					265					270	
His	Ile	Asn	Phe	Leu	Ala	Pro	Leu	Phe	Met	Val	Leu	Leu	Trp	Val	
				275					280					285	
Lys	Pro	Ile	Thr	Lys	Asp	Tyr	Ile	Met	Asn	Pro	Pro	Leu	Gly	Lys	
				290					295					300	
Glu	Ser	Ile	Pro	Leu	Met	Thr	Glu	Ala	Thr	Phe	Asp	Thr	Leu	Arg	
				305					310					315	
Leu	Trp	Leu	Ile	Ile	Leu	Leu	Cys	Ala	Leu	Arg	Leu	Ala	Met	Met	
				320					325					330	
Arg	Ser	His	Leu	Gln	Ala	Tyr	Leu	Asn	Leu	Ala	Gln	Lys	Cys	Val	
				335					340					345	
Asp	Gln	Met	Lys	Lys	Glu	Ala	Gly	Arg	Ile	Ser	Thr	Val	Glu	Leu	
				350					355					360	
Gln	Lys	Met	Val	Ala	Arg	Val	Phe	Tyr	Tyr	Leu	Cys	Val	Ile	Ala	
				365					370					375	
Leu	Gln	Tyr	Val	Ala	Pro	Leu	Val	Met	Leu	Leu	His	Thr	Thr	Leu	
				380					385					390	
Leu	Leu	Lys	Thr	Leu	Gly	Asn	His	Ser	Trp	Gly	Ile	Tyr	Pro	Glu	
				395					400					405	
Ser	Ile	Ser	Thr	Leu	Pro	Val	Asp	Asn	Ser	Leu	Leu	Ser	Asn	Ser	
				410					415					420	
Val	Tyr	Ser	Glu	Leu	Pro	Ser	Ala	Glu	Gly	Lys	Met	Lys	Val	Thr	
				425					430					435	
Val	Thr	Gln	Ile	Thr	Val	Ala	Leu	Ser	Ser	Leu	Lys	Asn	Ile	Phe	
				440					445					450	
Thr	Pro	Leu	Leu	Phe	Arg	Gly	Leu	Leu	Ser	Phe	Leu	Thr	Trp	Trp	
				455					460					465	
Ile	Ala	Ala	Cys	Leu	Phe	Ser	Thr	Ser	Leu	Phe	Gly	Leu	Phe	Tyr	
				470					475					480	
His	Gln	Tyr	Leu	Thr	Val	Ala									
				485											

<210> 27

<211> 350

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 3242083CD1

<400> 27

```

Met Asn Pro Arg Gly Leu Phe Gln Asp Phe Asn Pro Ser Lys Phe
 1          5          10          15
Leu Ile Tyr Thr Cys Leu Leu Leu Phe Ser Val Leu Leu Pro Leu
          20          25          30
Arg Leu Asp Gly Ile Ile Gln Trp Ser Tyr Trp Ala Val Phe Ala
          35          40          45
Pro Ile Trp Leu Trp Lys Leu Leu Val Val Ala Gly Ala Ser Val
          50          55          60
Gly Ala Gly Val Trp Ala Arg Asn Pro Arg Tyr Arg Thr Glu Gly
          65          70          75
Glu Ala Cys Val Glu Phe Lys Ala Met Leu Ile Ala Val Gly Ile
          80          85          90
His Leu Leu Leu Leu Met Phe Glu Val Leu Val Cys Asp Arg Val
          95          100          105
Glu Arg Gly Thr His Phe Trp Leu Leu Val Phe Met Pro Leu Phe
          110          115          120
Phe Val Ser Pro Val Ser Val Ala Ala Cys Val Trp Gly Phe Arg
          125          130          135
His Asp Arg Ser Leu Glu Leu Glu Ile Leu Cys Ser Val Asn Ile
          140          145          150
Leu Gln Phe Ile Phe Ile Ala Leu Lys Leu Asp Arg Ile Ile His
          155          160          165
Trp Pro Trp Leu Val Val Phe Val Pro Leu Trp Ile Leu Met Ser
          170          175          180
Phe Leu Cys Leu Val Val Leu Tyr Tyr Ile Val Trp Ser Leu Leu
          185          190          195
Phe Leu Arg Ser Leu Asp Val Val Ala Glu Gln Arg Arg Thr His
          200          205          210
Val Thr Met Ala Ile Ser Trp Ile Thr Ile Val Val Pro Leu Leu
          215          220          225
Thr Phe Glu Val Leu Leu Val His Arg Leu Asp Gly His Asn Thr
          230          235          240
Phe Ser Tyr Val Ser Ile Phe Val Pro Leu Trp Leu Ser Leu Leu
          245          250          255
Thr Leu Met Ala Thr Thr Phe Arg Arg Lys Gly Gly Asn His Trp
          260          265          270
Trp Phe Gly Ile Arg Arg Asp Phe Cys Gln Phe Leu Leu Glu Ile
          275          280          285
Phe Pro Phe Leu Arg Glu Tyr Gly Asn Ile Ser Tyr Asp Leu His
          290          295          300
His Glu Asp Ser Glu Asp Ala Glu Glu Thr Ser Val Pro Glu Ala
          305          310          315
Pro Lys Ile Ala Pro Ile Phe Gly Lys Lys Ala Arg Val Val Ile
          320          325          330
Thr Gln Ser Pro Gly Lys Tyr Val Pro Pro Pro Pro Lys Leu Asn
          335          340          345
Ile Asp Met Pro Asp
          350

```

<210> 28

<211> 450

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 3363391CD1

<400> 28

Met	Leu	Pro	Ser	Cys	Leu	Trp	Phe	Arg	Gly	Thr	Gly	Leu	Ile	Trp
1				5					10					15
Trp	Val	Thr	Gly	Thr	Ala	Ser	Ala	Ala	Gly	Leu	Leu	Tyr	Leu	His
				20					25					30
Thr	Trp	Ala	Ala	Ala	Val	Ser	Gly	Cys	Val	Phe	Ala	Ile	Phe	Thr
				35					40					45
Ala	Ser	Met	Trp	Pro	Gln	Thr	Leu	Gly	His	Leu	Ile	Asn	Ser	Gly
				50					55					60
Thr	Asn	Pro	Gly	Lys	Thr	Met	Thr	Ile	Ala	Met	Ile	Phe	Tyr	Leu
				65					70					75
Leu	Glu	Ile	Phe	Phe	Cys	Ala	Trp	Cys	Thr	Ala	Phe	Lys	Phe	Val
				80					85					90
Pro	Gly	Gly	Val	Tyr	Ala	Arg	Glu	Arg	Ser	Asp	Val	Leu	Leu	Gly
				95					100					105
Thr	Met	Met	Leu	Ile	Ile	Gly	Leu	Asn	Met	Leu	Phe	Gly	Pro	Lys
				110					115					120
Lys	Asn	Leu	Asp	Leu	Leu	Leu	Gln	Thr	Lys	Asn	Ser	Ser	Lys	Val
				125					130					135
Leu	Phe	Arg	Lys	Ser	Glu	Lys	Tyr	Met	Lys	Leu	Phe	Leu	Trp	Leu
				140					145					150
Leu	Val	Gly	Val	Gly	Leu	Leu	Gly	Leu	Gly	Leu	Arg	His	Lys	Ala
				155					160					165
Tyr	Glu	Arg	Lys	Leu	Gly	Lys	Val	Ala	Pro	Thr	Lys	Glu	Val	Ser
				170					175					180
Ala	Ala	Ile	Trp	Pro	Phe	Arg	Phe	Gly	Tyr	Asp	Asn	Glu	Gly	Trp
				185					190					195
Ser	Ser	Leu	Glu	Arg	Ser	Ala	His	Leu	Leu	Asn	Glu	Thr	Gly	Ala
				200					205					210
Asp	Phe	Ile	Thr	Ile	Leu	Glu	Ser	Asp	Ala	Ser	Lys	Pro	Tyr	Met
				215					220					225
Gly	Asn	Asn	Asp	Leu	Thr	Met	Trp	Leu	Gly	Glu	Lys	Leu	Gly	Phe
				230					235					240
Tyr	Thr	Asp	Phe	Gly	Pro	Ser	Thr	Arg	Tyr	His	Thr	Trp	Gly	Ile
				245					250					255
Met	Ala	Leu	Ser	Arg	Tyr	Pro	Ile	Val	Lys	Ser	Glu	His	His	Leu
				260					265					270
Leu	Pro	Ser	Pro	Glu	Gly	Glu	Ile	Ala	Pro	Ala	Ile	Thr	Leu	Thr
				275					280					285
Val	Asn	Ile	Ser	Gly	Lys	Leu	Val	Asp	Phe	Val	Val	Thr	His	Phe
				290					295					300
Gly	Asn	His	Glu	Asp	Asp	Leu	Asp	Arg	Lys	Leu	Gln	Ala	Ile	Ala
				305					310					315
Val	Ser	Lys	Leu	Leu	Lys	Ser	Ser	Ser	Asn	Gln	Val	Ile	Phe	Leu
				320					325					330
Gly	Tyr	Ile	Thr	Ser	Ala	Pro	Gly	Ser	Arg	Asp	Tyr	Leu	Gln	Leu
				335					340					345
Thr	Glu	His	Gly	Asn	Val	Lys	Asp	Ile	Asp	Ser	Thr	Asp	His	Asp
				350					355					360
Arg	Trp	Cys	Glu	Tyr	Ile	Met	Tyr	Arg	Gly	Leu	Ile	Arg	Leu	Gly
				365					370					375
Tyr	Ala	Arg	Ile	Ser	His	Ala	Glu	Leu	Ser	Asp	Ser	Glu	Ile	Gln
				380					385					390
Met	Ala	Lys	Phe	Arg	Ile	Pro	Asp	Asp	Pro	Thr	Asn	Tyr	Arg	Asp
				395					400					405
Asn	Gln	Lys	Val	Val	Ile	Asp	His	Arg	Glu	Val	Ser	Glu	Lys	Ile
				410					415					420
His	Phe	Asn	Pro	Arg	Phe	Gly	Ser	Tyr	Lys	Glu	Gly	His	Asn	Tyr
				425					430					435
Glu	Asn	Asn	His	His	Phe	His	Met	Asn	Thr	Pro	Lys	Tyr	Phe	Leu
				440					445					450

<210> 29
 <211> 400
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 3703614CD1

<400> 29
 Met Asn Pro Phe Leu Gly Asn Leu Pro Ser Ala Pro Ala Met Gly
 1 5 10 15
 Cys Ser Asp Ala Ser Thr Leu Asn Pro Gly Ser Ala Ser His Val
 20 25 30
 Ser Thr Tyr Thr Glu Asp Ser Gly Ser Ala His Gln Ser Arg Asp
 35 40 45
 Gln Val Phe Leu Pro Ala Phe Pro Val Gln Val Arg Arg Cys Lys
 50 55 60
 Ala Leu Lys Glu Lys Asp Leu Ile Arg Thr Ser Glu Ser Asp Cys
 65 70 75
 Tyr Cys Tyr Asn Gln Asn Ser Gln Val Glu Trp Lys Tyr Ile Trp
 80 85 90
 Ser Thr Met Gln Val Lys Ile Thr Ser Pro Gly Leu Phe Arg Ile
 95 100 105
 Val Tyr Ile Ala Glu Arg His Asn Cys Gln Tyr Pro Glu Asn Ile
 110 115 120
 Leu Ser Phe Ile Lys Cys Val Ile His Asn Phe Trp Ile Pro Lys
 125 130 135
 Glu Ser Asn Glu Ile Thr Ile Ile Ile Asn Pro Tyr Arg Glu Thr
 140 145 150
 Val Cys Phe Ser Val Glu Pro Val Lys Lys Ile Phe Asn Tyr Met
 155 160 165
 Ile His Val Asn Arg Asn Ile Met Asp Phe Lys Leu Phe Leu Val
 170 175 180
 Phe Val Ala Gly Val Phe Leu Phe Phe Tyr Ala Arg Thr Leu Ser
 185 190 195
 Gln Ser Pro Thr Phe Tyr Tyr Ser Ser Gly Thr Val Leu Gly Val
 200 205 210
 Leu Met Thr Leu Val Phe Val Leu Leu Leu Val Lys Arg Phe Ile
 215 220 225
 Pro Lys Tyr Ser Thr Phe Trp Ala Leu Met Val Gly Cys Trp Phe
 230 235 240
 Ala Ser Val Tyr Ile Val Cys Gln Leu Met Glu Asp Leu Lys Trp
 245 250 255
 Leu Trp Tyr Glu Asn Arg Ile Tyr Val Leu Gly Tyr Val Leu Ile
 260 265 270
 Val Gly Phe Phe Ser Phe Val Val Cys Tyr Lys His Gly Pro Leu
 275 280 285
 Ala Asp Asp Arg Ser Arg Ser Leu Leu Met Trp Met Leu Arg Leu
 290 295 300
 Leu Ser Leu Val Leu Val Tyr Ala Gly Val Ala Val Pro Gln Phe
 305 310 315
 Ala Tyr Ala Ala Ile Ile Leu Leu Met Ser Ser Trp Ser Leu His
 320 325 330
 Tyr Pro Leu Arg Ala Cys Ser Tyr Met Arg Trp Lys Met Glu Gln
 335 340 345
 Trp Phe Thr Ser Lys Glu Leu Val Val Lys Tyr Leu Thr Glu Asp
 350 355 360
 Glu Tyr Arg Glu Gln Ala Asp Ala Glu Thr Asn Ser Ala Leu Glu
 365 370 375
 Glu Leu Arg Arg Ala Cys Arg Lys Pro Asp Phe Pro Ser Trp Leu
 380 385 390
 Val Val Ser Arg Leu His Thr Pro Ser Asn

395

400

<210> 30
 <211> 133
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 4000975CD1

<400> 30
 Met Pro Arg Lys Leu Pro Ser Trp Arg Glu Ser Leu Phe Leu Ser
 1 5 10 15
 Val Glu Leu Ser Pro Leu Ala Leu Ala Met Gly Ser Ala Pro Gly
 20 25 30
 Leu Gln Val Phe Ser Lys Thr Asn Pro Leu Phe Leu Ser Pro Pro
 35 40 45
 Leu Lys Ser Arg Ala Leu Gly Pro Ser Pro Gln Glu Gly Phe Trp
 50 55 60
 Pro Asn Leu Gln Arg Gln Val Arg Ala Val Ser Leu Gly Cys Glu
 65 70 75
 Ala Ala Gly Glu Gly Asp Phe Gly Gln Met Ser Leu Gly Cys Glu
 80 85 90
 Ala Ala Gly Glu Gly Asp Phe Gly Gln Met Ser Leu Gly Cys Glu
 95 100 105
 Ala Ala Gly Glu Gly Asp Phe Gly Gln Val Ser Pro Ala Leu Cys
 110 115 120
 Pro Ser Gln Val Gln Leu Arg Asp Gly Leu Cys Leu Leu
 125 130

<210> 31
 <211> 359
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 4598831CD1

<400> 31
 Met Arg Ala Ala Ala Met Thr Thr Ala Ile Leu Glu Arg Leu Ser
 1 5 10 15
 Thr Leu Ser Val Ser Gly Gln Gln Leu Arg Arg Leu Pro Lys Ile
 20 25 30
 Leu Glu Asp Gly Leu Pro Lys Met Pro Cys Thr Val Pro Glu Thr
 35 40 45
 Asp Val Pro Gln Leu Phe Arg Glu Pro Tyr Ile Arg Thr Gly Tyr
 50 55 60
 Arg Pro Thr Gly His Glu Trp Arg Tyr Tyr Phe Phe Ser Leu Phe
 65 70 75
 Gln Lys His Asn Glu Val Val Asn Val Trp Thr His Leu Leu Ala
 80 85 90
 Ala Leu Ala Val Leu Leu Arg Phe Trp Ala Phe Ala Glu Ala Glu
 95 100 105
 Ala Leu Pro Trp Ala Ser Thr His Ser Leu Pro Leu Leu Leu Phe
 110 115 120
 Ile Leu Ser Ser Ile Thr Tyr Leu Thr Cys Ser Leu Leu Ala His
 125 130 135
 Leu Leu Gln Ser Lys Ser Glu Leu Ser His Tyr Thr Phe Tyr Phe
 140 145 150
 Val Asp Tyr Val Gly Val Ser Val Tyr Gln Tyr Gly Ser Ala Leu
 155 160 165

Ala His Phe Phe Tyr Ser Ser Asp Gln Ala Trp Tyr Asp Arg Phe
 170 175 180
 Trp Leu Phe Phe Leu Pro Ala Ala Ala Phe Cys Gly Trp Leu Ser
 185 190 195
 Cys Ala Gly Cys Cys Tyr Ala Lys Tyr Arg Tyr Arg Arg Pro Tyr
 200 205 210
 Pro Val Met Arg Lys Ile Cys Gln Val Val Pro Ala Gly Leu Ala
 215 220 225
 Phe Ile Leu Asp Ile Ser Pro Val Ala His Arg Val Ala Leu Cys
 230 235 240
 His Leu Ala Gly Cys Gln Glu Gln Ala Ala Trp Tyr His Thr Leu
 245 250 255
 Gln Ile Leu Phe Phe Leu Val Ser Ala Tyr Phe Phe Ser Cys Pro
 260 265 270
 Val Pro Glu Lys Tyr Phe Pro Gly Ser Cys Asp Ile Val Gly His
 275 280 285
 Gly His Gln Ile Phe His Ala Phe Leu Ser Ile Cys Thr Leu Ser
 290 295 300
 Gln Leu Glu Ala Ile Leu Leu Asp Tyr Gln Gly Arg Gln Glu Ile
 305 310 315
 Phe Leu Gln Arg His Gly Pro Leu Ser Val His Met Ala Cys Leu
 320 325 330
 Ser Phe Phe Phe Leu Ala Ala Cys Ser Ala Ala Thr Ala Ala Leu
 335 340 345
 Leu Arg His Lys Val Lys Ala Arg Leu Thr Lys Lys Asp Ser
 350 355

<210> 32
 <211> 72
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 4992201CD1

<400> 32
 Met Lys Cys Ala Leu Ile Thr Gly Arg Leu Arg Arg Gly Asn Glu
 1 5 10 15
 Thr Ser Cys Ile Asp His Arg Ala Gln Ser Leu Ala Phe Arg Lys
 20 25 30
 Pro Ser Val Arg Val His Asp Ala Met Val Ser Val Ile Ile Leu
 35 40 45
 Phe Ile Leu Ile Ile Thr Phe Ile Ile Phe Leu Leu Phe Leu Glu
 50 55 60
 Asn Ser Leu Glu Gly Leu Ile Pro Cys Tyr His Gly
 65 70

<210> 33
 <211> 112
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 5441583CD1

<400> 33
 Met Asn Ser Ala Met Trp His Gln Thr Ala Thr Gln Thr Thr Val
 1 5 10 15
 Ser Ser Ala Thr Leu Val Thr His Ile Gln Ala Arg Phe His Leu
 20 25 30
 Gln Gln Ser Trp Met Arg Trp Leu Ala Glu Ala Asn Pro Leu Pro

	35		40		45
Ala	Leu	Gln	Ala	Lys	Ala
	50		55		60
Ser	Leu	Thr	Ile	Leu	Arg
	65		70		75
Arg	Cys	Leu	His	Ala	Pro
	80		85		90
Ala	Thr	Cys	Tyr	Ser	Ser
	95		100		105
Glu	Arg	Phe	Ser	Cys	Pro
	110				

<210> 34
 <211> 149
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 1639243CD1

	34				
Met	Leu	Gly	Cys	Tyr	Gly
	1		5		10
Pro	Pro	Ala	Ser	Pro	Ser
	20		25		30
Leu	Pro	Ser	Leu	Gly	Leu
	35		40		45
Leu	Pro	Phe	Arg	Thr	His
	50		55		60
His	Pro	Ser	Thr	Pro	Val
	65		70		75
Val	Phe	Val	Phe	Phe	Lys
	80		85		90
Glu	Cys	Ser	Gly	Arg	Ile
	95		100		105
Gly	Ser	Ser	Asp	Ser	Pro
	110		115		120
Thr	Gly	Thr	Cys	His	His
	125		130		135
Glu	Met	Gly	Phe	His	His
	140		145		

<210> 35
 <211> 97
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 1335166CD1

	35				
Met	Glu	Ser	Cys	Ser	Val
	1		5		10
Leu	Gly	Ser	Leu	Gln	Pro
	20		25		30
Cys	Leu	Asn	Leu	Leu	Ser
	35		40		45
His	Trp	Leu	Phe	Phe	Val
	50		55		60
Val	Gly	Gln	Ala	Gly	Leu
	65		70		75

Ala Leu Ala Ser Gln Ser Ala Gly Ile Thr Gly Val Ser His Cys
 80 85 90
 Ala Arg Pro Gly Arg Leu Leu
 95

<210> 36
 <211> 104
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 166894CD1

<400> 36
 Met Phe Ser Glu Ala Leu Leu Ile His Arg Thr Tyr Leu Ala Tyr
 1 5 10 15
 Leu Phe Ala Cys Leu Leu Leu Met Ser Ser Leu Thr Glu Ser Leu
 20 25 30
 Leu Gln Arg Thr Thr Pro Ala Ser Arg Pro Arg Asn Val Gly Lys
 35 40 45
 Gly Lys Ala Trp Leu Val Leu Val Glu Met Glu Met Leu Val Thr
 50 55 60
 Val Glu Glu Cys Pro Pro Ser Asp Ser Gln Trp Gly Gly Ala Leu
 65 70 75
 Gly Pro Cys His Cys Pro Arg Thr Ser Ala Phe Gly Cys Pro Ala
 80 85 90
 Glu Arg Met Arg His Leu Ser Ser Ser Phe Trp Ser Pro Glu
 95 100

<210> 37
 <211> 99
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 217969CD1

<400> 37
 Met Ser Ser Leu Cys Val Ser Val Thr Ser Lys Asn His Asn Met
 1 5 10 15
 Phe Met Ala His Asp Gly Tyr Cys Ser Phe Val Phe Cys Phe Phe
 20 25 30
 Phe Glu Thr Glu Ser Ala Ser Val Thr Gln Pro Gly Val Gln Trp
 35 40 45
 Tyr His His Ser Ser Leu Gln Pro Arg Pro Pro Gly Leu Glu Gly
 50 55 60
 Ser Ser His Leu Ser Leu Gln Val Ala Arg Thr Ile Gly Val Cys
 65 70 75
 His His Thr Gln Leu Ile Leu Phe Arg Trp Gly Leu Thr Met Leu
 80 85 90
 Pro Trp Leu Val Ser Asn Phe Arg Ala
 95

<210> 38
 <211> 80
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 335237CD1

<400> 38

```

Met Pro Asp Leu Ala Val Val Leu Phe Cys Ser Arg Val Pro Arg
 1          5          10          15
Ser Ser Ser Gly Thr Gly Ser Gln Gly Gln Leu Val Pro Arg Ala
          20          25          30
Ser Leu Ala Cys Pro Leu Gly Ser Ser Arg Asp Asn Leu Thr Cys
          35          40          45
Pro Ile Lys Ala Lys Gly Gln Asn Arg Arg Gln Asn Leu Ala Arg
          50          55          60
Pro Ser Ser Asn Ser Lys Gly Lys Pro Val Pro Trp Ile Leu Ser
          65          70          75
Glu Ile Lys Thr Lys
          80

```

<210> 39

<211> 96

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 938306CD1

<400> 39

```

Met Lys Cys Lys Gly Ile Leu Ser Val Pro Gly Trp Leu Pro Thr
 1          5          10          15
Val Leu Gly Lys Arg Val Ile Phe Gln Lys Gly Pro Glu Gln Ser
          20          25          30
Ala Cys Ile Leu Ser Pro Leu Leu Pro Val Ser Ser Lys Ala Ser
          35          40          45
Gln Lys Leu His Phe Pro Thr Ser Cys His Phe Gln Asn His Ser
          50          55          60
Leu Asn Leu Lys Asn Lys Trp Glu Ala Val Phe Leu Pro Leu Met
          65          70          75
Ile Ala Ala Thr Tyr Lys Pro Ala Arg Thr Glu His Ser Lys Gln
          80          85          90
Arg Arg Val Gln Ser Cys
          95

```

<210> 40

<211> 92

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 1448129CD1

<400> 40

```

Met Ser Ala Met Phe Asn Ala Pro Trp Trp Ser Leu Gly Lys Met
 1          5          10          15
Pro Thr Pro Tyr Leu Leu Ser Leu Met Asn Ser Gln Ala Ser Phe
          20          25          30
Gly Gln Thr Phe Gln Gln Ala Leu Glu Ser Arg Leu Ile Val Thr
          35          40          45
Arg Glu Arg Tyr Lys Leu Gly Glu Arg Lys Glu Pro Phe Leu Glu
          50          55          60
Glu Ser Ala Phe Glu Gln Phe Leu Lys Val Leu Val Gly Arg Gly
          65          70          75
His Ser Arg Gln Val Gly Leu Phe Thr Glu Trp Thr Ala Val Trp
          80          85          90
Val Ala

```


<210> 41
 <211> 77
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 1761049CD1

<400> 41
 Met Ile Asn Val Trp Tyr His Val Phe Leu Gln Asn Ile Glu Phe
 1 5 10 15
 Lys Glu Cys Ser Leu Gln Tyr Trp Gln Leu Ser Pro Asp Leu Leu
 20 25 30
 Phe Asn His Gly Val Ile Ser Glu Lys Tyr Leu Phe Tyr Phe Ile
 35 40 45
 Leu Phe Tyr Phe Ile Leu Phe Met Leu Phe Met Leu Phe Met Leu
 50 55 60
 Cys Tyr Val Met Leu Cys Tyr Val Met Leu Cys Tyr Val Met Leu
 65 70 75
 Phe Phe

<210> 42
 <211> 75
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 1959587CD1

<400> 42
 Met Lys Leu Pro His Leu Ala Gln Phe Leu Thr Ser Pro Leu Val
 1 5 10 15
 Leu Trp Ser Thr Gly Val Ser Gly Ser Ala Gly Phe His Gln Leu
 20 25 30
 Val Pro Gln Trp Glu Cys Glu Glu Val Pro Gly Cys Gly Lys Ser
 35 40 45
 Cys Leu Ser Lys Arg Gly Leu Ile Glu Met Leu Gly Lys Val Ala
 50 55 60
 Val Ser Leu His Tyr Gly Arg Glu Gln Ser Gly Arg Ala Cys Cys
 65 70 75

<210> 43
 <211> 85
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 2303463CD1

<400> 43
 Met Ala Ile Phe Ser Leu Leu Met Phe His Ile Tyr Ser Phe Met
 1 5 10 15
 Arg Ile Phe Ser Phe Ala Leu Met Ser Val Phe Ile Ile Ala Ala
 20 25 30
 Phe Lys Phe Leu Ser Ala Val Tyr Ile Leu Asp Ile Leu Glu Met
 35 40 45
 Ala Thr Ala Cys Phe Leu Ser Cys Val Phe Ile Thr Phe Ser Arg
 50 55 60

Val Phe Thr His Leu Leu Asn Trp Lys Leu Cys Pro Gly Asp Cys
 65 70 75
 Ile Gln Asp Trp Ile Lys Lys Thr Gly Phe
 80 85

<210> 44
 <211> 89
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 2512281CD1

<400> 44
 Met Ala Ala Ala Pro Ala Pro Lys Pro Ser Leu Ala Pro Val Leu
 1 5 10 15
 Gly Pro Leu Glu Val Leu Pro Ala Pro Leu Gln Ala Pro Thr Arg
 20 25 30
 Arg Ser Pro Gly Thr Glu Cys Ala Pro Pro Ala Thr Gly Lys Gly
 35 40 45
 Arg Leu Ile Arg Val Arg Ser Arg Asp Gly Ile Val Thr Met Lys
 50 55 60
 Ser Ser Arg Arg Ala Met Cys Leu Lys Pro Ser Val Thr Leu Pro
 65 70 75
 Asn Ser Gln Glu Ala Arg His Ala Leu His Pro Ala Glu Pro
 80 85

<210> 45
 <211> 123
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 2755924CD1

<400> 45
 Met Phe Ile Lys Ile His Asn Leu Phe Phe Cys Ile Cys Val Leu
 1 5 10 15
 Pro Thr Leu Ala Ile Ser Gly Trp Ser Cys Pro Ser Leu Leu Ser
 20 25 30
 Leu Ser Phe Phe Lys His Ser Ile Cys Ile Leu Phe Leu Phe Leu
 35 40 45
 Val Thr Gly Phe His Tyr Val Ala His Thr Gly His Glu Leu Leu
 50 55 60
 Ser Ser Gly Asp Leu Pro Thr Ser Ala Ser Gln Val Ala Gly Thr
 65 70 75
 Thr Gly Thr Cys His Cys Ala Gln Leu Val Thr Ala Asn Phe Asn
 80 85 90
 Leu Gly Met Phe Val Pro Leu Leu Tyr Cys His Val Lys Asn Phe
 95 100 105
 Ala Asn Ser Gln Glu Thr Ser Val Ser Ser Val Lys Leu Asn Leu
 110 115 120
 Ser Ser Leu

<210> 46
 <211> 159
 <212> PRT
 <213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 2796369CD1

<400> 46

```

Met Gly Ser Leu Cys Ser Glu Asn Gly Arg Val Trp Asp Gly Leu
 1          5          10          15
Ser Phe Leu Leu Val Gly Pro Gly Ser Gly Ser Gly Ala Ala Pro
 20          25          30
Phe Leu Trp Ser Thr Gln Arg Glu Gln Glu Gly Leu Asp Leu Gly
 35          40          45
Lys Glu Ala Ile His Arg Ala Pro Gln Lys Pro Gly Pro Pro Gly
 50          55          60
Ala His Cys Cys Ala Glu Ala Thr Arg Leu Gly Tyr Phe Leu Pro
 65          70          75
Glu Ala Gly Asn Arg Glu Cys Arg Glu Ala Arg Gln Gln Gln Glu
 80          85          90
Ala Pro Asn Ala Gly Val Ser Lys Pro Glu Pro Pro Pro Asp Phe
 95          100          105
Thr Pro Val Cys Pro Ala His Ser Arg Leu Ser Leu Gly Gly Pro
 110          115          120
Trp Gly Leu Asp Leu Pro Asp Leu Trp Pro Gln Lys Gly Leu Ser
 125          130          135
Pro Glu Ser His Gly Met Glu Pro Gly Met His Arg Pro Ser Gly
 140          145          150
Leu Cys Leu Gly Ser Arg Pro Gly Ile
 155

```

<210> 47

<211> 77

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 3010920CD1

<400> 47

```

Met Ala Leu Arg Lys Ser Ser Cys Leu Pro Leu Lys Leu Gly Thr
 1          5          10          15
Leu Ile Thr Tyr Ser Leu Ile Phe Leu Ala Trp Phe Leu Leu Lys
 20          25          30
Ser Ala Thr Phe Asn Gln Val Ile Met Pro Arg Glu Leu Cys Gln
 35          40          45
Asp Leu Ile Tyr Val His Ser Tyr Asp Lys Tyr Leu Leu Ile Phe
 50          55          60
Gln Ile Asn Ser Cys Gly Cys Cys Asn Thr Tyr Ile His Tyr Arg
 65          70          75
Lys Leu

```

<210> 48

<211> 130

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 3360955CD1

<400> 48

```

Met Leu Phe Val Phe Ser Phe Cys Pro Gln Gln Ala Val Thr Ser
 1          5          10          15
Asp Gln Glu Val Ser Lys Ser Thr Glu Thr Leu Arg Arg Leu Met

```

	20		25		30
Leu Ser Ala Lys Ile	Met Asp Gly Glu Asp Thr Gly Leu Tyr His				
	35		40		45
Gln His Phe Ser Trp Tyr Leu Thr Ile Asn Arg Met Met Ala His					
	50		55		60
Arg Ser Lys Gly Thr Ser Phe His Ala Leu Pro Ser Leu Pro Ile					
	65		70		75
Leu Ala Asn Pro Ser Ser Trp Pro Pro Asp Tyr Asp Thr Thr Gln					
	80		85		90
Met Ser Ile Phe Ser Ala Arg Lys Ser Leu Leu Gly Thr Lys Leu					
	95		100		105
Leu Thr Ser Cys Leu Ser Ser Leu His Phe Arg Lys Cys Pro Val					
	110		115		120
Leu His Cys Asn Leu Leu Lys Ala Gly Lys					
	125		130		

<210> 49

<211> 97

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 3409459CD1

<400> 49

Met Asn Phe Tyr Arg Ala Ser Cys Leu Ser Leu Trp Val Phe Ala					
1	5		10		15
Gly Gly Gly Phe Gly Leu Asn Ala Ala Asp Met Ser Asp Ser Pro					
	20		25		30
Leu Ala Ala Ala Gly Glu Val Ala Ile Val Val Pro Leu His Pro					
	35		40		45
Gly His Leu Arg Cys Trp Tyr Leu Leu Asn Gln Gly Ile Trp Pro					
	50		55		60
Gly Arg Ala Ser Ser Pro Ala Pro Pro Ala Trp His Cys Pro Leu					
	65		70		75
Pro Val Leu Gln Arg Ala Ile Arg Lys Ala Gly Leu Pro Thr Leu					
	80		85		90
Leu Pro Arg Pro Ala Gly Pro					
	95				

<210> 50

<211> 74

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 4102938CD1

<400> 50

Met Pro Ser Leu Leu Asp His Pro Phe Ala Glu Lys Pro Phe Leu					
1	5		10		15
Leu Leu Ala Leu Phe Gln Leu Asn Phe Leu Ala Pro Leu Ser Gln					
	20		25		30
Val Ala Gly His Ala Ala Glu Gly Asn Trp Gly Asp Ser Arg Thr					
	35		40		45
Ala Asn His Phe Ser Lys Leu Arg Phe Gln Phe Glu Thr Arg Leu					
	50		55		60
Ala Asn Met Val Lys Pro Arg Leu Tyr Lys Lys Tyr Lys Asn					
	65		70		

<210> 51

<211> 74
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 4124601CD1

<400> 51
 Met Leu Gly Leu Gln Gln Gly Gln Gly Ser Ser Glu Arg Gln
 1 5 10 15
 Lys Trp Val Gly Pro Arg Gly Trp Arg Ala Ala Glu His Lys Ser
 20 25 30
 Arg Leu Lys Gly Ala Ala Thr Ala Gln Ser Pro Leu Thr Ala Ala
 35 40 45
 Gly Trp Asp Cys Lys Pro Arg Val Ala Arg Ser Val Ser Phe Phe
 50 55 60
 Gln Asp Lys Leu Glu Ile Arg Phe Ser His Gly Ile Val Ser
 65 70

<210> 52
 <211> 151
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 4180577CD1

<400> 52
 Met Ser Ser Ser Thr Ser Phe Ile Leu Ser Ala Ile Ala Ser Gly
 1 5 10 15
 Phe His Tyr Ser Leu Ser Ala Val Thr Ala Cys Gly Gln Leu Leu
 20 25 30
 Leu Leu Thr Ala Cys Arg Glu Leu Pro Asn Phe Ser Ser Gln Phe
 35 40 45
 Phe Leu Arg Ser Trp Leu Phe Trp Pro Gln Leu Lys Gly Val Leu
 50 55 60
 Leu Ser Ser Leu Arg Val Leu Ser Leu Phe Asp Pro Ile Val Val
 65 70 75
 Phe Ser Ser Phe Glu His Val Phe Gln Tyr Ser Tyr Phe Asn Leu
 80 85 90
 Leu Arg Thr Leu Lys Gly Asn Asp Lys Leu Val Val Gly Ile Trp
 95 100 105
 Gln Thr Gly Ala Cys Leu Phe Glu Arg Ser Ser Arg Arg Asp Lys
 110 115 120
 Ile Gln Ser Ala Ile Cys Phe Ser Trp Arg Gly Lys Arg Glu Asn
 125 130 135
 Leu Leu Asp Tyr Ile Leu Val Pro Trp His Thr Thr Tyr Met Phe
 140 145 150
 Lys

<210> 53
 <211> 137
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 5265807CD1

<400> 53

```

Met Leu Thr Tyr Ser Ser Phe His Phe Leu Leu Phe Tyr Leu Leu
 1          5          10          15
Leu Pro Leu Ser Leu Leu Ser Pro Ala Pro Gln Gln Lys Val Leu
          20          25          30
Gly Leu Leu Leu Ala His Ser Ala Asp Val Asn Ala Arg Asp Lys
          35          40          45
Leu Trp Gln Thr Pro Leu His Val Ala Ala Asn Arg Ala Thr
          50          55          60
Lys Cys Ala Glu Ala Leu Ala Pro Leu Leu Ser Ser Leu Asn Val
          65          70          75
Ala Asp Arg Ser Gly Arg Ser Ala Leu His His Ala Val His Ser
          80          85          90
Gly His Leu Glu Val Arg Thr Val Pro Ile Gln Ala Gln Leu Gly
          95          100         105
Leu Ser Leu Phe Leu Pro Ser Tyr Ser Arg Phe Pro Ala Ser Gly
          110         115         120
Pro Ser Ser Leu Lys Glu Lys Gln Pro Gly Trp Leu Tyr Lys His
          125         130         135
Leu Ser

```

<210> 54
 <211> 137
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 5405979CD1

```

<400> 54
Met Gln Ser Phe Thr Phe Tyr Leu Val Leu Pro Ser Pro Val Val
 1          5          10          15
Leu Ala Pro Pro Val Pro Ser Ala Ala Gly Pro Val Phe Ser Phe
          20          25          30
Gln Pro Arg Ser Ser Gln Pro Leu Leu His Gln Trp Cys Leu Leu
          35          40          45
Trp Ala Ser Pro Arg Leu Arg Cys Phe Arg Leu Ser Leu Leu Arg
          50          55          60
Gln Gln His Ala Ser Arg Trp His Ala Cys Pro Leu His Ala Ser
          65          70          75
Leu Gly Leu Pro Leu Leu Ala Gly Gln Gln Pro Ala Glu Pro Arg
          80          85          90
Tyr Leu Pro Phe Pro Cys Cys Ser Ser Leu Ser Pro Leu Ser Ser
          95          100         105
Trp Ala Cys Leu Gly Gln Lys Gly Gln Val Ser Gly Thr Ser Gln
          110         115         120
Glu Thr Leu Gly Arg Glu Val Ser Leu Ser Leu Glu Thr Val Asp
          125         130         135
Lys Leu

```

<210> 55
 <211> 205
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 7481109CD1

```

<400> 55
Met Met Arg Thr Leu Ile Thr Thr His Pro Leu Pro Leu Leu Leu

```

```

1           5           10           15
Leu Pro Gln Gln Leu Leu Gln Leu Val Gln Phe Gln Glu Val Asp
20           25           30
Thr Asp Phe Asp Phe Pro Glu Glu Asp Lys Lys Glu Glu Phe Glu
35           40           45
Glu Cys Leu Glu Lys Phe Phe Ser Thr Gly Pro Ala Arg Pro Pro
50           55           60
Thr Lys Glu Lys Val Lys Arg Arg Val Leu Ile Glu Pro Gly Met
65           70           75
Pro Leu Asn His Ile Glu Tyr Cys Asn His Glu Ile Met Gly Lys
80           85           90
Asn Val Tyr Tyr Lys His Arg Trp Val Ala Glu His Tyr Phe Leu
95           100          105
Leu Met Gln Tyr Asp Glu Leu Gln Lys Ile Cys Tyr Asn Arg Phe
110          115          120
Val Pro Cys Lys Asn Gly Ile Arg Lys Cys Asn Arg Ser Lys Gly
125          130          135
Leu Val Glu Gly Val Tyr Cys Asn Leu Thr Glu Ala Phe Glu Ile
140          145          150
Pro Ala Cys Lys Tyr Glu Ser Leu Tyr Arg Lys Gly Tyr Val Leu
155          160          165
Ile Thr Cys Ser Trp Gln Asn Glu Met Gln Lys Arg Ile Pro His
170          175          180
Thr Ile Asn Asp Leu Val Glu Pro Pro Glu His Arg Ser Phe Leu
185          190          195
Ser Glu Asp Gly Val Phe Val Ile Ser Pro
200          205

```

<210> 56

<211> 199

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 6247114CD1

<400> 56

```

Met Glu Thr Phe Pro Leu Leu Leu Leu Ser Leu Gly Leu Val Leu
1           5           10           15
Ala Glu Ala Ser Glu Ser Thr Met Lys Ile Ile Lys Glu Glu Phe
20           25           30
Thr Asp Glu Glu Met Gln Tyr Asp Met Ala Lys Ser Gly Gln Glu
35           40           45
Lys Gln Thr Ile Glu Ile Leu Met Asn Pro Ile Leu Leu Val Lys
50           55           60
Asn Thr Ser Leu Ser Met Ser Lys Asp Asp Met Ser Ser Thr Leu
65           70           75
Leu Thr Phe Arg Ser Leu His Tyr Asn Asp Pro Lys Gly Asn Ser
80           85           90
Ser Gly Asn Asp Lys Glu Cys Cys Asn Asp Met Thr Val Trp Arg
95           100          105
Lys Val Ser Glu Ala Asn Gly Ser Cys Lys Trp Ser Asn Asn Phe
110          115          120
Ile Arg Ser Ser Thr Glu Val Met Arg Arg Val His Arg Ala Pro
125          130          135
Ser Cys Lys Phe Val Gln Asn Pro Gly Ile Ser Cys Cys Glu Ser
140          145          150
Leu Glu Leu Glu Asn Thr Val Cys Gln Phe Thr Thr Gly Lys Gln
155          160          165
Phe Pro Arg Cys Gln Tyr His Ser Val Thr Ser Leu Glu Lys Ile
170          175          180
Leu Thr Val Leu Thr Gly His Ser Leu Met Ser Trp Leu Val Cys

```

41/91

<400> 58														
Met	Pro	Ser	Gly	Cys	Arg	Cys	Leu	His	Leu	Val	Cys	Leu	Leu	Cys
1				5					10					15
Ile	Leu	Gly	Ala	Pro	Gly	Gln	Pro	Val	Arg	Ala	Asp	Asp	Cys	Ser
				20					25					30
Ser	His	Cys	Asp	Leu	Ala	His	Gly	Cys	Cys	Ala	Pro	Asp	Gly	Ser

<400> 59														
Met	Ser	Arg	Leu	Glu	Ser	Ser	Glu	Ala	Ala	Cys	Arg	Ala	Val	Pro
1				5					10					15
Ser	Ala	Trp	His	Thr	Phe	Leu	Leu	Ser	Pro	Leu	Cys	Leu	Leu	Leu
				20					25					30
Ile	Gln	Val	Trp	Ala	His	Gly	Pro	Ser	Leu	Gln	Val	Val	Thr	Lys

	35		40		45									
Val	Ala	Pro	Pro	Ala	Leu	Thr	Ser	Ser	Met	Ser	Asp	Ser	Leu	Val
				50					55					60
Phe	Thr	Lys	His	Phe	Ser	Leu	Cys	Lys	Val	Ile	Asp	Ser	Ala	Asn
				65					70					75
Val	His	Arg	Gly	Cys	Thr	Thr	Cys	Gln	Ala	Leu	Val	Lys	Ala	Arg
				80					85					90
Asp	Val	Glu	Thr	Ser	Cys	Cys	Arg	Phe	Ser	Ala	His	Ala	Leu	Ala
				95					100					105
Gly	Glu	Ala	Val	Ser	Gln	Gln	Asn	Lys	Gln	Arg	Gly	Gly	Glu	Ala
				110					115					120
Ala	Ser	Cys	Leu	Leu	Arg									
				125										

<210> 60
 <211> 137
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 2414185CD1

	<400>	60												
Met	Met	Gly	Lys	Leu	Ser	Pro	Thr	Phe	Ile	Leu	Gly	Ile	Cys	Trp
	1			5					10					15
Val	Pro	Ala	Gly	Leu	Gly	Tyr	Gln	Gln	Gly	Lys	Lys	Thr	Trp	Pro
				20					25					30
Leu	Ser	Ser	Ser	Pro	Tyr	Asn	Leu	Gln	Asp	Lys	Met	Tyr	Ala	Leu
				35					40					45
Glu	Lys	Ala	Gly	Asp	Pro	Ser	Lys	Ala	Arg	Ser	Met	Gly	Pro	His
				50					55					60
Lys	Ser	Pro	Glu	Thr	Gln	Arg	Gly	Gln	Pro	Met	Glu	Met	Ser	Gly
				65					70					75
Leu	Lys	Gly	Gln	Val	Thr	Ser	Thr	Ala	Leu	His	Thr	Leu	His	Phe
				80					85					90
Pro	Arg	Arg	Pro	Pro	Ser	Gly	Cys	Gln	Thr	Asp	Gln	Ala	Gly	Asp
				95					100					105
His	Glu	Pro	Gly	Gly	Arg	Phe	Leu	Ala	Gln	Pro	Gln	Arg	Leu	Arg
				110					115					120
Glu	Leu	Ser	Leu	Met	Ile	Ser	Pro	Leu	Gln	Leu	Leu	Pro	Phe	Gly
				125					130					135
Ser	Arg													

<210> 61
 <211> 77
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 5266594CD1

	<400>	61												
Met	Leu	Glu	Gly	Thr	Leu	Pro	Leu	Pro	Thr	Val	Leu	Leu	Val	Gly
	1			5					10					15
Arg	Pro	Val	Leu	Leu	Leu	Ala	Leu	Gly	Ala	Ala	Val	Pro	Gly	His
				20					25					30
Leu	Ala	Ala	Pro	Thr	Asp	Val	Glu	Leu	Pro	Glu	Leu	Leu	Leu	Asn
				35					40					45
His	Cys	Ala	Gly	Arg	Val	Val	Ala	Leu	Ile	Val	Gly	Ala	Arg	Val
				50					55					60

Leu Leu Leu Phe His Tyr Ile Pro Val Gly Ile Ile Ile Pro Gly
 65 70 75
 His Ile

<210> 62
 <211> 110
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 7610617CD1

<400> 62
 Met Ser Asn Leu Trp Leu Leu Val Gly Ala Arg Ala Cys Ser Leu
 1 5 10 15
 Ser Leu Leu Thr Tyr Ser Phe Leu Gly Asp Leu Ile Pro Ser His
 20 25 30
 Cys Leu Lys His Leu Pro Gly Thr Gly Val Ile His Leu Cys Ser
 35 40 45
 Ser Ser Ser Glu Ile Pro Ser Ala Pro Phe Ile His Leu Phe Ile
 50 55 60
 His Ser Ala Asn Ile Cys Gly Ile Ser Val Pro Gly Thr Ala Leu
 65 70 75
 Gln Pro Gly Cys Thr Ile Gly Thr Gln Thr Asp Thr Pro Phe Pro
 80 85 90
 Met His Ser Leu Leu Thr Asp Thr Pro Ala Trp Gln Cys Leu Gly
 95 100 105
 Val Phe Thr Ala Pro
 110

<210> 63
 <211> 103
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 1902436CD1

<400> 63
 Met Thr Phe Ile Tyr Thr Phe Ile Leu Ser Phe Phe Leu Gln Leu
 1 5 10 15
 Cys Cys Ser Phe Met Lys Leu Ile Leu Leu Ile Ser Asn Thr Asn
 20 25 30
 Ala Val Ser Phe Ile Leu His Arg Pro Cys Thr Leu Cys Ser Asp
 35 40 45
 Phe Tyr Ser His Ile Cys Met Leu Leu Thr Val Ser Val Asn Phe
 50 55 60
 Leu Ser Phe Trp Asn Asn Phe Gln Thr Ile Leu Thr Trp Ala Asp
 65 70 75
 Leu Phe Ser Met Leu Leu Ala Tyr Glu Tyr Arg Phe Thr Arg Leu
 80 85 90
 Phe Ser Val Leu Pro His Thr Ser Val Met Leu Cys Phe
 95 100

<210> 64
 <211> 192
 <212> PRT
 <213> Homo sapiens
 <220>

<221> misc_feature

<223> Incyte ID No: 2310369CD1

<400> 64

```

Met Ala Pro Lys Pro Gly Ala Glu Trp Ser Thr Ala Leu Ser His
 1          5          10          15
Leu Val Leu Gly Val Val Ser Leu His Ala Ala Val Ser Thr Ala
          20          25          30
Glu Ala Ser Arg Gly Ala Ala Ala Gly Phe Leu Leu Gln Val Leu
          35          40          45
Ala Ala Thr Thr Thr Leu Ala Pro Gly Leu Ser Thr His Glu Asp
          50          55          60
Cys Leu Ala Gly Ala Trp Val Ala Thr Val Ile Gly Leu Pro Leu
          65          70          75
Leu Ala Phe Asp Phe His Trp Val Asn Gly Asp Arg Ser Ser Ala
          80          85          90
Asn Leu Leu Leu Gly Gly Gly Met Val Leu Ala Val Ala Gly Gly
          95          100          105
His Leu Gly Pro Glu Gly Arg Ser Val Ala Gly Gln Ala Met Leu
          110          115          120
Leu Val Val Ala Val Thr Ile Leu Ile Val Ala Val Phe Thr Ala
          125          130          135
Asn Thr Tyr Gly Met Trp Gly Gly Ala Met Leu Gly Val Ala Gly
          140          145          150
Leu Leu Ser Arg Leu Glu Glu Asp Arg Leu Leu Leu Leu Pro Lys
          155          160          165
Glu Asp Val Cys Arg Trp Ala Leu Ala Val Gly Ser Trp Ala Tyr
          170          175          180
Cys Arg Ala Leu His Thr Gln Arg Leu Gln Trp Glu
          185          190

```

<210> 65

<211> 310

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 6180576CD1

<400> 65

```

Met Asn Gly Leu Ser Leu Ser Glu Leu Cys Cys Leu Phe Cys Cys
 1          5          10          15
Pro Pro Cys Pro Gly Arg Ile Ala Ala Lys Leu Ala Phe Leu Pro
          20          25          30
Pro Glu Ala Thr Tyr Ser Leu Val Pro Glu Pro Glu Pro Gly Pro
          35          40          45
Gly Gly Ala Gly Ala Ala Pro Leu Gly Thr Leu Arg Ala Ser Ser
          50          55          60
Gly Ala Pro Gly Arg Trp Lys Leu His Leu Thr Glu Arg Ala Asp
          65          70          75
Phe Gln Tyr Ser Gln Arg Glu Leu Asp Thr Ile Glu Val Phe Pro
          80          85          90
Thr Lys Ser Ala Arg Gly Asn Arg Val Ser Cys Met Tyr Val Arg
          95          100          105
Cys Val Pro Gly Ala Arg Tyr Thr Val Leu Phe Ser His Gly Asn
          110          115          120
Ala Val Asp Leu Gly Gln Met Ser Ser Phe Tyr Ile Gly Leu Gly
          125          130          135
Ser Arg Leu His Cys Asn Ile Phe Ser Tyr Asp Tyr Ser Gly Tyr
          140          145          150
Gly Ala Ser Ser Gly Arg Pro Ser Glu Arg Asn Leu Tyr Ala Asp
          155          160          165

```

```

Ile Asp Ala Ala Trp Gln Ala Leu Arg Thr Arg Tyr Gly Ile Ser
170 175 180
Pro Asp Ser Ile Ile Leu Tyr Gly Gln Ser Ile Gly Thr Val Pro
185 190 195
Thr Val Asp Leu Ala Ser Arg Tyr Glu Cys Ala Ala Val Val Leu
200 205 210
His Ser Pro Leu Thr Ser Gly Met Arg Val Ala Phe Pro Asp Thr
215 220 225
Lys Lys Thr Tyr Cys Phe Asp Ala Phe Pro Asn Ile Glu Lys Val
230 235 240
Ser Lys Ile Thr Ser Pro Val Leu Ile Ile His Gly Thr Glu Asp
245 250 255
Glu Val Ile Asp Phe Ser His Gly Leu Ala Leu Tyr Glu Arg Cys
260 265 270
Pro Lys Ala Val Glu Pro Leu Trp Val Glu Gly Ala Gly His Asn
275 280 285
Asp Ile Glu Leu Tyr Ser Gln Tyr Leu Glu Arg Leu Arg Arg Phe
290 295 300
Ile Ser Gln Glu Leu Pro Ser Gln Arg Ala
305 310

```

<210> 66
 <211> 135
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 2274523CD1

```

<400> 66
Met Phe Ala Gln Pro Phe Ser Pro Ile Arg Ala Ser Lys Arg Met
1 5 10 15
Ala Lys Val Ser Ser Asn Asn Phe Ala Ser Leu Pro Arg Gln Ala
20 25 30
Pro Met Leu Leu Phe Cys Pro Leu Trp Met Pro Val Thr Ser Val
35 40 45
Pro Gln Glu Ala Lys Leu Leu Arg Gln Leu Lys Phe Ser Gln Gly
50 55 60
Thr Gly Val Cys Val Leu Ile Tyr Thr Pro Leu His Thr Tyr Phe
65 70 75
Phe Lys Leu Ser Pro Thr Leu Gly Thr Pro Val Leu Glu Tyr Pro
80 85 90
Gln Gln Phe Lys Gly Lys Lys Arg Leu Lys Gln Lys Asp Phe Phe
95 100 105
Leu Pro Lys Leu Cys Leu Leu Ala Trp Gly Pro Arg His Ala Asp
110 115 120
Leu Lys Ile Asn Gln Ala Trp Val Gly His Gly Gly Ser Arg Leu
125 130 135

```

<210> 67
 <211> 205
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 1801820CD1

```

<400> 67
Met Val Asn Leu Ala Ala Met Val Trp Arg Arg Leu Leu Arg Lys
1 5 10 15

```

Arg Trp Val Leu Ala Leu Val Phe Gly Leu Ser Leu Val Tyr Phe
 20 25 30
 Leu Ser Ser Thr Phe Lys Gln Glu Glu Arg Ala Val Arg Asp Arg
 35 40 45
 Asn Leu Leu Gln Val His Asp His Asn Gln Pro Ile Pro Trp Lys
 50 55 60
 Val Gln Phe Asn Leu Gly Asn Ser Ser Arg Pro Ser Asn Gln Cys
 65 70 75
 Arg Asn Ser Ile Gln Gly Lys His Leu Ile Thr Asp Glu Leu Gly
 80 85 90
 Tyr Val Cys Glu Arg Lys Asp Leu Leu Val Asn Gly Cys Cys Asn
 95 100 105
 Val Asn Val Pro Ser Thr Lys Gln Tyr Cys Cys Asp Gly Cys Trp
 110 115 120
 Pro Asn Gly Cys Cys Ser Ala Tyr Glu Tyr Cys Val Ser Cys Cys
 125 130 135
 Leu Gln Pro Asn Lys Gln Leu Leu Leu Glu Leu Phe Leu Asn Arg
 140 145 150
 Ala Ala Val Ala Phe Gln Asn Leu Phe Met Ala Val Glu Asp His
 155 160 165
 Phe Glu Leu Cys Leu Ala Lys Cys Arg Thr Ser Ser Gln Ser Val
 170 175 180
 Gln His Glu Asn Thr Tyr Arg Asp Pro Ile Ala Lys Tyr Cys Tyr
 185 190 195
 Gly Glu Ser Pro Pro Glu Leu Phe Pro Ala
 200 205

<210> 68

<211> 2569

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 3211795CB1

<400> 68

ctccggaccctc ttaaggcgc gccagagtc ctcccgagcga aaaacgactt aaaggagacg 60
 cgtggcgcgca tggcgggcgc cccacgcgcg ggccggcgcc gcgggcagcc gctcctggcg 120
 ctgctgcttc tgctgctggc gccactgccg cggggggccc cgccggcgcc cgacgcctac 180
 ttcccgagg agcgtggag cccggagtcg cccctgcagg cgccgcgctg gctcatcgcg 240
 ctgttggcgc gaaacgcggc ccacgcgttg cccaccacgc tgggcgcact cgagcggctg 300
 cggcaccgcg gggagcgcac ggcgctatgg gtggctacgg accacaacat ggataaacag 360
 tcaactgtgc tgcgggagtg gctgggtggc gtgaagagtt tgtaccattc cgtggagtg 420
 cggccagcag aggagcccag gtactaccg gacgaggaag gcccgaaca ctggtctgac 480
 tcacgctacg agcatgtcat gaagtgcgc caggcagccc tgaaatcagc tcgagacatg 540
 tgggctgatt acatcctggt tgtagatgcg gacaacctga tcctcaacct tgacacactg 600
 agcctgctca tcgctgagaa caagacggtg gtgcgcccga tgctggattc ccgggctcg 660
 tactccaact tctgggtgtg aatgacttcc cagggctact acaagcgcac acctgcctac 720
 atccctatcc gcaagcgaga ccgcccgggc tgctttgcag ttcccatggg gactcagacc 780
 ttccctgatcg acctgcggaa ggccggcgctc aggaacctgg ccttctaccc acctcaccct 840
 gactacacct ggtcctttga cgacatcatc gtctttgcct tctcctgcaa gcaggcagag 900
 gttcagatgt atgtgtgcaa caaggaggag tacggattct tgccagtgcc attgcgcgcc 960
 cacagcacc tccaggatga ggccgagagc ttcatgcatg tgcagctgga ggtcatggtg 1020
 aagcaccgc cgcagagacc ctcccgttc atctcggctc ccaccaagac accggacaag 1080
 atgggcttcg acgaggtctt catgatcaac ctgaggcgcc ggcaggaccg gcgggagcgc 1140
 atgtgcggg cgctgcaggc acaggagatc agtgccggc tggtgaggc cgtggacggc 1200
 aaagccatga acaccagcca ggtggaggcg ctggggatcc agatgctgcc tggctaccgg 1260
 gaccctacc acggccggcc cctcaccaag ggtgagctgg gctgcttctt gagccactac 1320
 aacatctgga aggaggtggt ggaccggggg ctgcagaaat cgcttggtgt tgaggatgac 1380
 ctgctgtttg agatcttctt caagagacgt ctgatgaacc tcatgcggga tgtggagcgg 1440
 gagggcctgg actgggacct catctatgtg gcccggaagc ggatgcaggt ggagcacc 1500
 gagaaggctg tgcctcgcgt gaggaacctg gtggaggccg actattccta ctggaccctg 1560

```

gcttacgtga tctccctgca aggcgcgcgc aaactgctgg ctgctgagcc gctctccaag 1620
atgtgcctg tggacgagtt cctgcccgtc atgttcgaca aacaccagc gtccgagtac 1680
aagcccact tctccctccg caacctgcat gccttctctg tggagccgct gctcatctac 1740
cccacacact acacaggaga cgatggctat gtgagtgaca ccgagacctc agtcgtatgg 1800
aacaatgagc acgtcaagac cgactgggac cgcgcgaagt ccagaagat gcgggagcag 1860
caggcactga gccgtgaggg caagaactcg gacgtgctcc agtccccact ggacagtgtc 1920
gcccgggatg aactctgagg ggtagcagcc agaaagccaa agcagccatc ggtggcccag 1980
gctccacgtg cttactgagg acatcagggc cacctctgga ccccttgga gccacagag 2040
ggctctcgtg tgggggtggg tccagccagc tcttgctaag caatcacgtg cacacaggca 2100
gcattaatgg agtgcctact gcagccagc aacagggctt ggccctgggg aattgggagg 2160
aaccaagccc tcttcatctg ttcattgtgc cagcatttat taagcacctg ctgtatgcaa 2220
ggttcccatg ttacggcagt gaatgaggca taattgttcc ctccatcagc gattgattca 2280
gtcatcaagc agttactgat cagattaaga atcaggcact agtgatacac attcatTTTT 2340
aaaattcatt caaggattta ttgagtgcct actgtgtgtt gggtgccatt ccaggctctg 2400
ggattttttt tttttttttt ttttaagagt agagtctgtc tctgtcacc aggctggagt 2460
gcagtgtgtg gacggctcac tgcagcctgc gcctccagc gtccagcaat tcttgtttct 2520
cggcctccca agtagctggg actataggtg cgtgccatca catctggct 2560

```

<210> 69

<211> 2387

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 6813464CB1

<400> 69

```

atggcggggc cccctcccc cgccttgetg ctgccttgca gtttgatctc agactgctgt 60
gctagcaatc agcgacactc cgtggcgta ggaccctccg agctagtcaa gaagcaaatt 120
gagttgaagt ctcgagggtg gaagctgatg cccagcaaaag acaacagcca gaagacgtct 180
gtgttaactc aggttgggtg gtcccaagga cataatatgt gtccagacc tggcataccc 240
gaaaggggca aaagactagg ctcgatttcc aggttaggat ccagcgcca gttcacctgc 300
aacgagggct atgacctgca aggggtccaag cggatcacct gtatgaaagt gagcgacatg 360
tttgccgctt ggagcgacca caggccagtc tgccgagccc gcatgtgtga tgcccacctt 420
cgaggccctt cgggcatcat cacctcccc aatttcccc ttcagtatga caacaatgca 480
cactgtgtgt ggatcatcac agcactcaac cctccaagg tgatcaagct cgcctttgag 540
gagtttgatt tggagagggg ctatgacacc ctgacggctg gtgatgggtg tcaggatggg 600
gaccagaaga cagttctcta catctgaca ggtacatcgg tcccggtatc cattgtcagc 660
accaatcatc aaatgtggct cctcttccag actgatggca gtggcagttc cctgggattc 720
aaggcttctt atgaagagat cgagcagggc agttgcggtg accctggcat acctgcata 780
ggccggaggg aaggctccc gtttcaccac ggtgacacac tcaagtttga gtgccagccc 840
gcctttgagc tgggtggaca gaaggcaatc acatgcaaaa agaataacca atggctcggc 900
aagaagccag gctgcgtgtt ctctgcttc ttcaacttca ccagcccgtc tggggttgc 960
ctgtctccca actaccaga ggactatggc aaccacctcc actgtgtctg gctcatcctg 1020
gcccaggcctg agagccgcat ccacctggcc ttcaacgaca ttgacgtgga gcctcagttt 1080
gatttctctg tcatcaagga tggggccacc gccgagggcg ccgtcctggg cacttctca 1140
gaaaaccagc ttccctctc catcacaagc agtggccacg tggcccgctc cgagttccag 1200
actgaccact ccacaggga gaggggcttc aacatcactt ttaccacctt ccgacacaac 1260
gagtgcctgg atcctggcgt tccagtaaat ggcaaacggg ttggggacag cctccagctg 1320
ggcagctcca tctccttct ctgtgatgaa ggcttctctg ggactcaggg ctccagagacc 1380
atcacctgct tctgaagga gggcagcgtg gtctggaaca gcgctgtgct gcggtgtgaa 1440
gtccctgtg gtggtcacct gacttcgccc agcggacca tctctctcc gggtggcct 1500
ggcttctaca aggatgcctt gagctgtgcc tgggtgattg agggccagcc aggetacccc 1560
atcaaaatca ccttcgacag gtgcttgtga ccacagcctt gggggagagg gcaaggggga 1620
gatttttaca gagccacgtg ggagggaatg cagggtggat aggggaccag taccaacgct 1680
gaaatgtcag caaggttggg gaacactgga ccaaatggcc ctgggtgggg cagacctcac 1740
tggctctgcc cttgggctt gtttgtgctg tttaggggac tatggcagga ggggtgaaga 1800
gaaggcttgg gtaccatttg ggatgttaga gctccccct cccagcccc caaggagggg 1860
gagtgggcag atagacacca tcagatgtta aagcatgact ctggtatctg acgagcattc 1920
tcttgaaaat gtccaacgtg ggaaggcagc cccccacctc tgcccaaat cctgtctatg 1980
tgcccactt ctagtccca cctggctat tggagacgt agccacctaa 2040
atgtgaaaac aaacttcttc cggacaaagt gttcctggcc tgggaatagca gtctgcaaac 2100

```



```

ttttgtgtga agagccagat ggtaaattatt ttaggctttg tggggcactt gcctacttgc 2160
atatctctga tgcattgctct gctcttggcg cctttttttt ttcttaagat ctcttttaca 2220
atgtaaaaag catccttagc tctccaggct atagaccaga tctgggccct gaggtcttta 2280
gtgtgctgac ccctagccta gaatgttctg acctccatga ggctagtcat tgtgcctcta 2340
ggtcttaagc tgggctcaca gatcaggcca cagagaggat gagagct 2387

```

<210> 70
 <211> 1959
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 2156540CB1

```

<400> 70
ggccggctgg gcggtgcgtc gctccccgaa gccggggctg ggccggagcc gggcgagggc 60
tgggagctgg gccgggtccg gggacagcgg gcgaggggca gctgccggag ccgggcagcc 120
aggccgctca gggcagggga cagctggcgc cggttctgcg gtctccgggg cccagatgtg 180
aggcggcggc gcccccgcc cgagagcgca cgatgggggc cccgctcgcc gtagcgctgg 240
gcgcccctca ctacctggca cttttcttgc aactcgggcg cgccacgcgg ccgcgcggcc 300
acgcgccctg ggacaaccac gtctccggcc acgcccgtgt cacagagaca ccccatgaca 360
tgacagcacg gacgggcgag gacgtggaga tggcctgctc ctcccgggc agcggtctcc 420
cctcctactc gctggagatc cagtgggtgt atgtacggag ccaccgggac tggaccgaca 480
agcaggcgtg ggcctcgaac cagctaaaag catctcagca ggaagacgca gggaaggagg 540
caacccaaat aagtgtggtc aagggtggtg gcagcaacat ctcccacaag ctgcgcctgt 600
cccgggtgaa gcccacggac gaaggcacct acgagtgcg cgtcatcgac ttcagcgacg 660
gcaaggcccg gcaccacaag gtcaaggcct acctgcgggt gcagccaggg gagaactccg 720
tctgcatctc gccgaagcc cctcccgcgg cgcccggccc gccgcccccc aagccaggca 780
aggagctgag gaagcgctcg gtggaccagg aggcctgcag cctctagact gatgccctg 840
ccccgcacca tccgccccca cgctgtacag agtgcattag gagccgcccg accaccgggg 900
accgactgcc tgcgtccagc cgtgccccat cccgaggcc gcctgtggcc accatgtcgg 960
cctcttttcc accaccctt gctcagcatg taagccccac ccaccctgc cttttcagac 1020
ccctgcgggtg acctggctcg gagaagggtg ccctgggcac caaggggcca accgcctga 1080
acactggggc agggaccatg ctggggcccg gggccacccc ctctctgtca ccagcttctg 1140
tggagtccag tgttttgcct tgcctgtctg tccccatcc tgcctgagc cggggccccc 1200
cagcctcgcc tccctcctcc taccatccct cacttggacc tgggggtgtg gacagtgacc 1260
cctccctgaa tcttgacttg aatcttctga gcagaactag ggcctctccc ctggtgaaga 1320
cccagggaac ccaggagggc ccttctgggg cagtggctct gcagggtcac tcatggaggc 1380
ctaggggaac agcgagatgc cccaccacct cctggcgagt ccttctgtt cagctccctg 1440
tgcgaccctc cagggatgca ggggatccag gattctctgc cctgtcacac ggcgagtcag 1500
aaggaggggg cctttccctc ggaccatgg cccaggcag agttttgcac cagcaggacc 1560
cctttgaggg ccttcaaggc tctcccagga gtccccctct gccggccccc caatgcccc 1620
gtcctctctt gggctcctgt ccaagtccgc cccagggcct ggggctgttg ggagccaagg 1680
gccccctggt actcagttcc ctcacgattc ccgatcacgg gcacacctgc cccctgggta 1740
tttgtaaata tttctatttg acccaattct cctcggaatt ggctggcacc tctggctgcc 1800
acagctcagt gatgacgtg gggaggtggg agaggccgag ggctttgcct aggggtgggt 1860
tgccctgtat acatgatcca gtctgtgact accagccaac ctgaataaag cggtttttaa 1920
aaaaaaaaa aaaaaaaaaa atgcggcgca agcttattc 1959

```

<210> 71
 <211> 1562
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 894939CB1

```

<400> 71
cctcttccgt cggctgaatt gcggccgtat gcgcggctct gtggagtga cctgggggtg 60
ggggcactgt gccccagcc cctgctcct ttggactcta cttctgtttg cagccccatt 120
tggcctgctg ggggagaaga cccgccaggt gtctctggag gtcattcccta actggctggg 180

```

```

ccccctgcag aacctgcttc atatacgggc agtggggcacc aattccacac tgcactatgt 240
gtggagcagc ctgggggcctc tggcagtggt aatgggtggcc accaacacccc cccacagcac 300
cctgagcgtc aactggagcc tcctgctatc ccctgagccc gatgggggcc tgatgggtgct 360
ccctaaggac agcattcagt tttcttctgc ccttggtttt accaggctgc ttgagtttga 420
cagcaccaac gtgtccgata cggcagcaaa gcctttggga agaccatata ctcactactc 480
cttggccgat ttctcttga acaacatcac tgattcattg gatcctgcca ccctgagtgc 540
cacatttcaa ggccacccca tgaacgaccc taccaggact tttgccaatg gcagcctggc 600
cttcagggtc caggcctttt ccaggtcag cggaccagcc caacccccct gcctcctgca 660
cacagcagac acctgtcagc tagaggtggc cctgattgga gcctctcccc ggggaaaccg 720
ttccctgttt gggctggagg tagccacatt gggccagggc cctgactgcc cctcaatgca 780
ggagcagcac tccatcgagc atgaatatgc accggccgtc ttccagttgg accagctact 840
gtggggcttc tcctccatg gctttgcaca gtggcgacca gtggcttact cccagaagcc 900
ggggggccga gaatcagccc tgccctgcca agcttcccc ctccatcctg ccttagcata 960
ctctcttccc cagtcaccca ttgtccgagc ctctcttggg tcccagaata acttctgtgc 1020
cttcaatctg acgttcgggg ctccacagg ccctggctat tgggaccaac actacctcag 1080
ctggctgatg ctccctgggtg tgggcttccc tccagtgga ggcttgtccc cactagtctc 1140
gggcatcatg gcagtgccc tgggtgcccc agggctcatg ctgctagggg ggggcttggg 1200
tctgctgctg caccacaaga agtactcaga gtaccagtc ataaattaag gcccgctctc 1260
tggagggaag gacattactg aacctgtctt gctgtgcctc gaaactctgg aggttggagc 1320
atcaagttcc agccggcccc ttcactcccc catcttgctt ttctgtggaa cctcagaggc 1380
cagcctcgac ttccctggaga cccccagggt gggcttccct catactttgt tgggggactt 1440
tggaggcggg caggggacag ggctattgat aaggtccctt tgggtgtgcc ttcttgcatc 1500
tcacacatt tcccttggat gggacttgca ggcctaaatg agaggcattc tgactgggtg 1560
gc 1562

```

<210> 72

<211> 3425

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 4620890CB1

<400> 72

```

tcccagggtg atccccagca cggccgggtg ctgcgggcgc ggcagcccca gaagacggga 60
aagtctcgcg cgggagcgcg gagatgcgag gtagcgacac ggcgctcacc gtggaccgga 120
cctactcgga ccccgccgg caccaccgct gcaagagccg ggtagaacgt catgacatga 180
ataccttaag cctgccccctg aacatacgcc gaggggggtc agacaccaac ctcaactttg 240
atgtcccgga tggcatcctg gacttccaca aggtcaaaact cactgcagac agcctgaagc 300
aaaaaattct aaaggtaaca gagcagataa aaattgagca aacatcgcgc gatgggaatg 360
ttgcggagta tctgaaacta gtgaacaacg cggacaagca gcaggcggga cgtatcaagc 420
aagtctttga gaagaagaat cagaaatcag ctactccat cgcccagctg cagaagaagt 480
tagagcagta tcatcgaaag ctcagagaga tcgagcagaa tggagcctct aggagctcaa 540
aggacatttc caaagaccac ctgaaggata tacatcgctc tttgaaagat gccacgtga 600
aatctcgaa cgtcccccat tgcatggaga gcagcaaatc gggcatgcca ggggtctcac 660
ttactccacc tgtgtttgtt ttcaataagt ccagagagtt tgccaacctg atccggaata 720
agttttggcag cgcgcgacaac attgtctact tgaaaaatc cttagaggag tttaggccag 780
aggcgagtgc cagggcctac gggggcagcg ctaccatcgt gaacaaacc aagtatggca 840
gtgatgatga atgttcgagt ggcacgtcag gctcggccga cagtaacgga aaccagtcgt 900
ttggggcttg tggagccagc acactggaca gccagggcaa gctcggcgtg atcctggagg 960
aactgagggg gatcaaggat acccaagctc agctggctga ggacatcgag gcaactgaag 1020
tgcagtttaa gagagaatat ggttttattt ctacagacct gcaagaggaa agatacaggt 1080
atgagcgact ggaagaccag ctgcatgacc tgacggacct gcacagcat gagacagcca 1140
acctgaagca ggagctggcc agcattgagg agaagggtgg ctaccaggcc tacgagcgct 1200
cgcgggacat ccaggaagcc ttggaatcct gccagactcg catttctaag ctggagctcc 1260
accagcaaga gcagcaagct ctgcagacag acaccgtgaa tgctaaagtt ctctgggga 1320
ggtgcatcaa cgtgatcctg gccttcatga ctgtcatctt agtgtgtgtg tccaccatcg 1380
cgaagtctgt ctaccccatg atgaagagtc gctgccacat tcttggcacc ttctttgccg 1440
tgactcttct tgctatattt tgtaaaaact gggaccatat cctgtgtgcc atagaaagga 1500
tgataatacc aagatgaagc cactggttcc tgccttcaag ttctttcaag tttttatttt 1560
aaagaaaact ctgtgcatac taccaaattt tacagtgaat gattgtgagg actcgtgtgt 1620
aagaaaaact aggactgtgt ggtgtaaata actacaattc tcttaactcg gtagcagttg 1680

```

```

ccaactcagt ccttgtaactt cgttaacacg aatctgtttc agagctctcc taccttgctc 1740
actgccttaa tcagaccgat ttcctgcccc cctgaccagc ccagcgtggg aaacctctgt 1800
atattgagac cttggcataa ttggtgatcc tgaagaaaaga ggtctctctc ctaagtctct 1860
gtcagaattg agcttcacaa ttgctaattg ttgttttctg tgagtcctat aaaaagcaag 1920
gatatgcattg attcagggaa tgaagaatca caggcttggg cagtgtttaa cactgtggcc 1980
tatggtcccc gtgtgatcca cctgtcttct ctccagggga ccatagggtcc cgtcatgtac 2040
tcagtgtcca cagcagtcag tcgtgtatga ccctgtaacg tggaaatctt atcacacacc 2100
tgttatccaa caagtctacc tgagggggtt ttgttacctt taaatgggaa ggcataggga 2160
tttatgaatg gggctttcac cttctcatal ccaggcaacc aacacctgat tttgtctcaa 2220
ctggctagca aatgccacgc cttcagagtg tgcaggaatg ttttcaaata cctcatcaga 2280
ctgtgacttt aacattaatt tggaaatctg tgagcactac tctgaagggt tgtgttttgg 2340
caaactctttt ttcttttttg agacagggct ctgctaaata ttgctcaggc tgggttgcaa 2400
ctccttgctt caagggatcc tcccacctca gcctcccaag cagccgggac tgcaggcaca 2460
agccaccatg cctggctgtt ttttggcaaa tcttgattgt gataagcccc cctggaggat 2520
atgattcact ttatgtgatt catcttattc acaggtctgt gagggactgc gaagcttact 2580
caggaaatga aaacaaatga tggatcatgt gcagtttttt ccttgaagga caaccgaacc 2640
atagcctcta aggttcaagt gcactgaggt gtcggaacgc tgaaagcatg aggaaacgag 2700
gacgtagggt gtgactgaat ggtggctaga ttagtgggag cagttcacct ggatgaagat 2760
tgagagcatc gtcttttgaga agtgaaagac tagcaagaat aaaataaatt aagtccagt 2820
tttgagccaa ggttgccacc tgtctcttaa catctcactg aacataagtc ctgaggtatt 2880
aggacgacca tactgcctct gagctgaaaa cattcaaaag ttcacatccc tgtttggggg 2940
ataccattca ccgccttcag cccagatgat actttccttt aaatctgtgt ctctgtgtgt 3000
atacaaaaga ggaagatgga aacaatgttc atggaaactg ctgttgagcc ccttgtccca 3060
ccactccccg catctgtctg aggcaggaag gcatgtgagt gtacgttttc ttccaggaga 3120
catcaggtcc ccttgattc aaattaagt caatattttg caaacagctc ttcttaggga 3180
aatctcctga agggaaaaaa tgtgacagaa tgttccatag tctgagagaa tggaaatcgt 3240
gagcatttag tacaagtcca gtgtgtgtga gcgggactta ggcagctcaa gcttgctttt 3300
ttttttaagc gtacaattga gtgggttttag taaattcaca aacttggtca accatcacca 3360
ctatctaatt ccagactcac gcttttttaa acaataaatg tcatttcatg aaaaaaaaaa 3420
aaaaa

```

<210> 73

<211> 3130

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 5514146CB1

<400> 73

```

gggacagtca aagaatttta aaaacaggga actttgtaac tgtgaaatac tctccaggat 60
ttaaaaggct gtggagctcc agataaagaa tcgtttatct ttcttctgaa gaaattcctt 120
tggttacaag ttaccctcat aaacggcaac aactcacct ccatccaaga cagactcaag 180
gtggagggaag cgtggaaatg tgcttccgga caaagctctc agtatcctgg gtgccattgt 240
ttcttctact cagccgtgtt ttttctactg agacagacaa accctcagcc caggacagca 300
gaagccgtgg gagttcaggc caaccggcag acctgctaca ggttctctct gctggtgacc 360
accaccccca caaccactca agaagcctca tcaaaacatt gttggagaaa actgggtgcc 420
cacggaggag aaacgggaat caaggagatt gcaatctgtg ctttgaacca gatgcactat 480
tactaatagc tggaggaaat tttgaagatc agcttagaga agaagtgggt cagagagttt 540
ctcttctcct tctctattac attattcatc aggaagagat ctgttcttca aagctcaaca 600
tgagtaataa agagtataaa ttttacctac acagcctact gagcctcagg caggatgaag 660
attcctcttt cctttcacag aatgagacag aagatatctt ggctttcacc aggcagtact 720
ttgacacttc tcaaagccag tgtatggaaa ccaaaacgct gcagaaaaaa tctggaatag 780
tgagcagtga aggtgctaatt gaaagtacgc ttcctcagtt ggcagccatg atcattactt 840
tgtccctcca ggggtgtttgt ctgggacaag gaaacttgcc ttccccagac tactttacag 900
aatatatttt cagttccttg aatcgtacga atacactccg cctatcagaa ctagaccaac 960
tcctcaacac tctctggacc agaagtactt gtatcaaaaa tgagaaaaat catcaatttc 1020
aaaggaaaca aaacaacata ataaccatg atcaggacta ttctaatttc tcttcatcca 1080
tgaaaaaaga gtctgaggat ggtccaattt cctgggatac gacctgcttc tctgctaggc 1140
agctgggtga gatatttcta cagaagggcc tctcactcat ttctaaggag gactttaagc 1200
aaatgagtc agggatcatc cagcagctcc tcagctgctc ctgccactta ccaaggacc 1260
aacaagcaaa gctgccacct accactctgg agaaatacgg ctacagcacg gtggctgtca 1320

```

```

cccttctcac actgggctcc atgctgggga cagcgctgggt ccttttccat agctgtgagg 1380
agaactacag gcttatctta cagctgtttg tgggcttgge cgtcgggaca ctgtctgggg 1440
acgctctgct ccaccttate cctcaggttc ttggtttaca taagcaggaa gccccagaa 1500
ttgggcattt ccatgaaagc aaaggtcata ttggaaact gatgggatta attggaggca 1560
tccatggatt tttcttgata gaaaaatgtt ttattcttct tgtatcacca aatgacaagc 1620
agggcctgtc attggttaat gggcacgtgg gtcattccca ccatcttgca ctcaactctg 1680
aattaagtga ccaggcaggc agaggcaaat ctgcttcaac tatccagttg aaaagcccag 1740
aagattcaca ggcagctgaa atgcctatag gcagtatgac agcctccaac agaaaaatga 1800
aagccattag cttggttagca atcatgatc tgggtgggga cagcctgcat aattttgcag 1860
atggcctagc cataggagca gccttctcat catcatccga gtcaggagtg accactacga 1920
ttgctatctt gtgtcatgaa atcccacatg aaatgggaga ctttgccgtg ctcttaagct 1980
ctggactttc tatgaagact gccatcctga tgaattttat aagctcccta actgccttca 2040
tgggattata cattggcctt tccgtgtcag ctgatccatg tgttcaagac tggatcttca 2100
cagtcactgc tgggatgttc ttataattat ccttggttga aatgcttctt gaaatgactc 2160
atgttcaaac acaacgacct tggatgatgt ttctcctgca aaactttgga ttgatcctag 2220
gttggtcttc tctcctgtc ttggctatat atgagcaaaa tattaataa taagtggagg 2280
tcttcaacat ctttcaaaaa tgcatttata tagtcttact ttgtttcttt cattgcactc 2340
tataatgatt tttaaattaa gaatttttta tcttaggcaa agtgtgtctc tttcaattca 2400
ttaacttatt aattttataa tgcagtttta tttttggaaa catataaata tcagactgtc 2460
cttaattgaa attttgtctt tggtttccaa caccatgatg aagctcttgc tttttaaaaa 2520
gtagtttagta aattctgcat gaattttagt aaactttaaa aaatagattt tttccctaag 2580
aaagaattgt tgtagaattt aaagtggaca gatgctgtt ggggtaaaat caactgcaac 2640
tttttgatgt taattttttt cctgttgcaa ttataaacta taagcaagtt aagtgcagag 2700
caaatgtaat aaagactagt tttaaaaaaa aaaaaaaggg gggttttctc cccggccctg 2760
ggctatattt gggcaaaaatt ttaaaatata agtgggggtt tcaacattt tcaaaaaagc 2820
cattaatagg gtctaacttg gtctcttcca tggcactcta aaaggttttt aaatttagga 2880
tttttttacc ttgggcaaaag tggggctctt ccaatcctta acctttttaa ttttataagg 2940
cggtttttatg tttggaaaaac cttttaaatt ccgagcgggc ctaaatggaa atttggcttt 3000
gggttcccaa ccccggtgtg aacccttgc ttttaaaaag tagtgggggc ccgggcccga 3060
atccgaaaca tgctaaaggg gttcccccgg gaatttttta cccgcaaaa tttcccaaaa 3120
tatggggcgg                                     3130

```

<210> 74

<211> 3172

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 7474769CB1

<400> 74

```

ggctcctgta gacaatatac ccgtgaatgg tgtagttagt tctaataaaa ctttatctac 60
aagaggagac tgaccaggtt tggcctgggg gccacagtg gtagacctct ggaaagatac 120
atcctgagaa gaaaaaaaga atatatgcag gaatgtttaa ctttgtgggt tttctctcct 180
cttgcctca ctgactcagg atacacaaag acctatcaag ctacagcaa gcagaaattc 240
agccgcttat ggtccagcaa gtctgtcact gagattcacc tatactttga ggaggaaatc 300
aagcaagaag aatgtgacca tttggaccgc ctttttgctc ccaaggaagc tgggaaacag 360
ccacgtacag tgatcattca aggaccacaa ggaattggaa aaacgacact cctgatgaag 420
ctgatgatgg cctggtcgga caacaagatc tttcgggata ggttctctga cagttctat 480
ttctgtgca gagaactgag ggagttgccc ccaacagatt tggctgactt gatttcaga 540
gagtggcctg accccgctgc tcctataaca gagatcgtgt ctcaaccgga gagactctg 600
ttcgtcatcg acagcttoga agagctgcag ggcggcttga acgaaccgga ttcggatctg 660
tgtggtgact tgatggagaa acggccggtg caggtgcttc tgagcagttt gctgaggaag 720
aagatgctcc cggaggcctc cctgtctate gctatcaaac ccgtgtgccc gaaggagctc 780
cgggatcagg tgacgatctc agaaatctac cagccccggg gattcaacga gattgatagg 840
ttagtgtatt tctgtgtgtt cttcaaagac ccgaaaagag ccatggaagc cttcaatctt 900
gtaagagaaa gtgaacagct gttttccata tgccaaatcc cgctcctctg ctggatcctg 960
tgtaccagtc tgaagcaaga gatgcagaaa ggaaaagacc tggccctgac ctgccagagc 1020
actacctctg tgtactcctc tttcgtcttt aacctgttca cacctgaggg tgccgagggc 1080
ccgactccgc aaaccagca cagctgaag gccctgtgct ccctggctgc agagggtatg 1140
tggacagaca catttgagtt ttgtgaagac gacctccgga gaaatggggg tgttgacgct 1200
gacatccctg cgctgctggg caccaagata cttctgaagt acggggagcg tgagagctcc 1260

```

```

tacgtgttcc tccacgtgtg tatccaggag ttctgtgccg ccttggttcta tttgctcaag 1320
agccaccttg atcatcctca cccagctgtg agatgtgtac aggaattgct agttgccaat 1380
tttgaanaag caaggagagc acattggatt tttttggggt gttttctaac tggcctttta 1440
aataaaaaag aacaagaaaa actggatgcg ttttttggtc tccaactgtc ccaagagata 1500
aagcagcaaa ttcaccagtg cctgaagagc tttaggggagc gtggcaatcc tcagggacag 1560
gtggattcct tggcgatatt ttactgtctc ttgaaatgc aggatcctgc ctttgtgaag 1620
caggcagtga acctcctcca agaagctaac ttcatatta ttgacaacgt ggacttggtg 1680
gtttctgcct actgcttaaa atactgtctc agcttgagga aactctgttt ttccgttcaa 1740
aatgtcttta agaaagagga tgaacacagc tctacgtcgg attacagcct catctgttgg 1800
catcacatct gctctgtgct caccaccagc gggcacctca gagagctcca ggtgcaggac 1860
agcaccctca gcgagtgcac ctttgtgacc tgggtgaacc agctgaggca tcccagctgt 1920
cgcttccaga agcttggaa aaataacgtt tccttttctg gccagagtgt tctgctcttt 1980
gaggtgctct tttatcagcc agacttgaaa tacctgagct tcacctcac gaaactctct 2040
cgtgatgaca tcaggtccct ctgtgatgcc ttgaactacc cagcaggcaa cgtcaaagag 2100
ctagcgctgg taaattgtca cctctcacc attgattgtg aagtccttgc tggccttcta 2160
accaacaaca agaagctgac gtatctgaat gtatcctgca accagttaga cacaggcgtg 2220
ccccttttgt gtgaagccct gtgcagccca gacacgggtc tggatatacct gatgttggct 2280
ttctgccacc tcagcgagca gtgctgcgaa tacatctctg aaatgcttct gcgtaacaag 2340
agcgtgcgct atctagacct cagtgcgaat gtctgaagg acgaaggact gaaaactctc 2400
tgcgaggcct tgaaacatcc ggactgctgc ctggattcac tgtgtttggt aaaatgtttt 2460
atcactgctg ctggctgtga agacctcgcc tctgctctca tcagcaatca aaacctgaag 2520
attctgcaaa ttgggtgcaa tgaaatcgga gatgtgggtg tgagctgttt gtgtcgggct 2580
ctgacgcata cggattgccg cttagagatt cttgggttgg aagaatgtgg gttacagagc 2640
acctgctgta aggatctcgc gtctgttctc acctgcagta agacctgca gcagctcaac 2700
ctgaccttga acaccttgga ccacacaggg gtggtgttac tctgtgaggc cctgagacac 2760
ccagagtgtg ccctgcaggc gctcgggctg agaaaaactg attttgatga ggaaccagc 2820
gcacttctga cggctgagga agagagaaat cctaacctga ccatcacaga cgactgtgac 2880
acaatcacaa gggtagagat ctgattgcga ggaacctggg ctctgactcg aacacctgca 2940
aaggacaggg actgggaccg ttacttacat gacactgcac ccaggagata caaatcattg 3000
acactctgag ttgtgagatt tctggcacc cattcataga tttgatatga tacacgtggt 3060
ttttatgtgc tctgtggcct tggatgagtc actgaaaggc cttcatggtc tctcggcttc 3120
acaaggacct cttaaccctt caataaagtg ttacatttct aaacattgga aa 3172

```

<210> 75

<211> 2094

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 065296CB1

<400> 75

```

tgtgtgctga ttttttaatg aacaacttat aatcttacaa aataactggc tgtagctgtt 60
ttacattaca atacagtaga tcagctcctt tcaactacatt cccaatcaag ccatccctt 120
gggtgttaac taatgcactc aatagtgatc acaagaata caggaagtgc caagttcaat 180
gccttgtaat agaaaaaaga gaagatgaga atcaggaaga gctctttact ttcacaatcc 240
ctttacttta atttctgtca gtatctgccc attctctca cctgtcccc ttacctgcct 300
ttctgattgg aggctgtcat gctagtgttt agagccagct agccctgggc cacctgggca 360
caatcaaaca cacagaagag tctctgagaa aggtctotcaa tgaccacatg ggtggataag 420
agtagcagaa tgtggaccac cacatggatt gagtgtgagt tgatgttgct gggatctgtg 480
accacagtgt cttagtcctt ccactttatc agtgacctga gttgctcaaa acatcaggtc 540
ccaggaacaa agactggaag aacctgctgt ctctgaatag ctatagctga ctatggctgg 600
gaagccttat agcctaagag gctcttcaca cacaacagga actttccttt tgetttccca 660
gagcagtggt gaattgcaga taataaaata ttttaaaatg aaatttaaaa cagaaatgtt 720
cttactgttg ctcttggtga gagactgcac gaagacacac actggcatga accacaggct 780
ccatgttctc gagctgagta atgccagga taacaattcc tctgcctcaa tctcagacaa 840
agtaggattc tccaaggcag aacttagaat gtgtctagca atttggacat tttccccaat 900
aaaacaagtt tataaaattt tgaaaataga gtgtctaaac ttttccattg tcttagcgt 960
attgaagcct atcagaatcc caagaataaa tatgtttgtc ttcttggggg cactaagcat 1020
gaccaagac aatgaatggt atctgaacta cattttcttt acactagaga ttagtagaca 1080
aaagggtttt tttgaatggg tgaactctgc tctttccttt tcccaataac tttgcaacct 1140
tcattccaca aaccacaga ttaatgtggg tacatcacct tattcgtgtt ttattctctt 1200

```

```

tcaggatatt gttcttccct acgtgatagt gtggataata tttctatcat tgtggtagta 1260
tttctatctt ctatttgcac agccattgct ctagtggtga tgggaagagt agaaaaactg 1320
tgggtcaaagc aaaagagcag aagatgagaa gagaccacag ctgagctcta aattttgtct 1380
gttcttgggt tccattggag aagaaacttt aaaggagaca agcctcactc acagccttcc 1440
taaaatgagt aaagctagaa tgtaaacag aaactggctc cttactcact ttccctaggg 1500
tctaattccag aaaccgttac aagaaaatgt aactgcagta gaccacatcc cagttagcct 1560
gggatgtaaa tagagagagg cttaatatgg tcatttcatg tgcattccat ctattggata 1620
agaaaaggta aaaacaacaa aatggacatt aaatagtga aattcttctg tctcccaca 1680
gagtgtcacc cttgtccat gcatttccat ggcttgcac agtcaaaatc tctggctcac 1740
agattttgta catgatactt gaagagaatt tatctcagat tcttcattgc tcaagggacc 1800
cactttgggt ttctttggag acttggaaac gaggaccaag tctccctgat cacttccatt 1860
ctcatgcccg ggccctggat ttaaccctcg acaggtctt ggccgtgtct ttcattctgt 1920
tccagctttt ccctattcat ttccatgaaa ttttctcca gaaaattaaa gttggtctgt 1980
gactggcgac ttgggacttc tacagaagtt ggatcgggca gtttgtggtt gtgtggtcct 2040
ggatcggggt caaaggggtg tcatttgtgg ggccctgggt ggtttttga accg 2094

```

<210> 76

<211> 1119

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 231994CB1

<400> 76

```

ggcgagctg agggaccg cggctggtga gcgcccgtg gaggctggag cttccgggcc 60
ctggaaagg ggtccccgc cgccccgggt cggaggcaga cccctgggtt tgggggacat 120
gggcatttgg ggccctgaa cccaagacct ctggatgagc tgccccgttc agaccatgga 180
tcttgaggtg accttgcgtc tgcagtgcct tggcgggggc ctgccccagg agcagatata 240
ggccgagctg agccccgccc atgaccgtcg cccactgcca ggtggggacg aggccatcac 300
tgccatctgg gagaccggc taaaggccca accctggctc ttcgacgccc ccaagtccg 360
ctgcactca gccaccctgg cgcctattgg ctctcgggg ccacagctgc tcttgcgcct 420
gggccttact tctaccgag acttccctgg caccaactgg tccagctcag ctgcctggct 480
gcgacagcag ggtgccaccg actgggggtga cagcaggcc tatctggcgg acccactggg 540
ggtgggctg gactagcca cagccgatga ctctctgtc ttctgccc gctcccggca 600
ggtggctgag gcccttggc tgggtggacgt acctgtggg caccctgagc ctgagccct 660
gtgccctggg ggcagcccc agcaccagga cctcgctggg cagctggtgg tacatgaact 720
cttttccagt gtccctcagg agatctgtga tgaggtgaac ctgcccgtgc tcacctgag 780
ccagcccctg ctgttgggca tcgcccgaaa tgagaccagt gctggccgag ccagtgccga 840
gttctatgtc cagtgcagcc tgacttctga gcaggtgagg aagcactacc tgagtggggg 900
acccgaggcc cagcagctca caggaatctt ctttgtggg acacagaacg tgcggagatt 960
gcccgagacg gagatgtggg ctgaactctg cccctcgccc aaaggcgcca tcacctcta 1020
caaccgggtt cagggaagtc ccaactggagc ggccctaggg tccccagccc tactcccgc 1080
gctctgaaaa taataaacga ctttattctt ggaaaaaaa 1119

```

<210> 77

<211> 3321

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 538054CB1

<400> 77

```

gtcgccgggt gaggaagcgg ggggctagcg gcctggcgct gcggcgaccg accggggcgt 60
caggatccct ggcccccgga gagcgaaggc cgggcgggtc ccggagcaag aagaattgta 120
tagaaatgta aaaaacattt atgtgtgctt cacatgtggc atgaaaaaaa tgaggacctc 180
tggtcaaaat tgcctgagtt cagaatttca tgcagggaag aaagtttagac gtaacttcta 240
atttctagtt ccagtaagca tctcttcttg gaaaatttcc aaaaaagagt cctgctctgt 300
tgccaggct ggagtacatt ggtgtgatga tcatgggtca ctgcagcctc gacctcccag 360
gctcaagcaa ttctcttacc tcagcctcct gagtaactgg gactatagga tttctttgaa 420

```

```

gaaaaatgtg cttcttgaat caactgcagt ctctcctctg cataagagct cttaagggtg 480
ctagcaggat .agaagcaaat gaaactgaaa gctcatctgc agctcagaaa agcaaagaca 540
tggaatttta aagagtgaag gtagcatggt gtcggccatg ggtgaacaag acacagccag 600
acaatgtgga ccaatttctt caaactacgg cttttctgct gtctgtctgc agtgttgatg 660
gtggtggtgc tggatcaaa tgttactcag gtagagtact tggaccatga gactgtttca 720
gccactttca tcgacagcag tggacagttt gtttctctcc aggtgacagg aattagccga 780
aatccctact gtggctatga tcagcagacc ctgtccagcc aggagcgcat ggaggaggac 840
tccttgctgg ctgacctgca ccggcagggt cctgatgtgg gccagtcctt ctttgtgaag 900
agcactgacc cttcttccag ctactttgtc atcttgaact ctgctgcctt ctttaagggtg 960
ggaagccagc ttgagggtgt ggttcatgtg caggattttc aaagaaagcc caagaagtat 1020
ggtggagact acctgcaggc cagaattcac tccctcaagc tgcaggctgg ggctgtgggc 1080
agggtggtgg attaccagaa tgggttttac aaagttttct ttactttgtc atggccaggc 1140
aaagttaaag tatccgtatc tctggtccac ccagtgaaag ggatcagagt tcttcagcgc 1200
ttacaggaag ataaaccaga cagggtctat ttcaagagtc tcttccgttc aggaagaatt 1260
tctgaaacta ctgagtgcga cgtgtgtctt cctgggaatc tgcccctgtg taactttaca 1320
gacctctaca ctggggagcc ctggttctgc ttcaaaccac agaagctccc ttgcagcagc 1380
agaattaccc atttcaaagg tggatacctg aaaggtctcc taaccgctgc agagagtgtc 1440
ttcttccaga gtggtgtcaa tatcaaatg ccagtcaact ccagtgacc tgattgggta 1500
actgtgattc ccaggagaat aaaagaaact aacagtctag aactatctca aggtcaggga 1560
acttttctt ctgggtatta ttataaagac cagtggaggc ccagaaagt ttatagtcgt 1620
cagtttaagt accctgacaa cattacagag tgcttcaaaa gaaaagtggg gcattttatt 1680
ggtgactcaa caatcaggca atggtttgaa taccttacta catttggtcc agatttagtg 1740
gagtttaact tgggtagtcc caagaatgtg ggtcccttcc ttgcagtggg ccagaagcac 1800
aacatcctgc tcaaataccg ctgccatggt ccaccatccc gcttcacgac tgtcttttagc 1860
aatgagctcc attatgtggc gaatgagctg aatggcattg tgggagggggaa gaacacagtg 1920
gttgccatag ctgtatggtc tcacttcagc accttccctt tggagtgta catccggcgg 1980
ctcaggaaca tccgtcgagc agtgggtcgg ctctcgcagc gaagcccaaa gaccgtgggtg 2040
gtcatccgga cggccaacgc ccaagagctg ggacctgagg tgagcctttt caacagcgac 2100
tggtacaact ttcaagctgga caccatcctt cggaggatgt tctcaggggt tggagtatat 2160
ctcgtcgatg cctggggagat gaccttgccc cattatctac cgcacaagct gcacccagat 2220
gaagtatttg tgaagaacca gctggacatg ttcttgcctt ttgtgtgccc cttggaaacc 2280
tagcctgtct tgggaaggac tggaggaatc atattcaatg accttctcaa ttgacctgag 2340
ttacagaaag tggccccagt gagagatgac tgcccttaat aagtataaaa tttcaaaaag 2400
atctggactt aatatgatga cttataagga gcttagaaaa tgcagggttac atttatatct 2460
acctatagga ttttatccaa tgttgactta gccatggtag aactcttaac tgcacttaca 2520
cactatattg ctcttgaac caaagatgct aatgagtgtg tttgaattag cttctcctag 2580
gaggggtgac tactttgcta aagagtatga aaaatgtttt gatggaaggg acaagtttgg 2640
ttggttagtag agtgtttgta gcttatctgg taaaggaaat tgagtgcac tcgcatgcata 2700
gcacaagtaa tccatcttct ggccgttaac tagagaaagc ttatagtcag aggtctctgt 2760
gaatgcactg aagactgctg tgtctcagga gtttctcttg atgatctacc ttttctgaat 2820
tacttgaaaa ggccaatgtg cttgggtcga gttttttagt gaattttac aatttctgc 2880
ttttatctag aaaagaagca aaagggaat atgaaagcag tatatataaa atctgtgctt 2940
tggaattgat atttgcattt ggtgattgtt ttttatttga ttcagctaaa ttttatgtgt 3000
tgaatactta ctctagaata actacttttg gaactggaag ggggaaatct gtaaatggaa 3060
atattttttt atgttctgaa atttgagggt ttaggaattg acttttgtgt aaagaatgct 3120
acggatttat aaagctgtgg aagttttttc cactagtttc aggtgttttt tttgttgtt 3180
gttggtgttg ttatttttga ctacttacca aaattagaag ttaaaatcta attagaaatt 3240
aataagatac taaaattcat gacaaaaaaa aaaacaaaaa acacaaaaaa aaaaaaaa 3300
aaaaacaaaa aaaaaaaaaa a 3321

```

<210> 78

<211> 2646

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 1259305CB1

<400> 78

```

gtcccttccc ctacccgct ccacgccctc ctgggccgag tggagttggg tgggtgtcggg 60
agcctctccc tgaggggcac cgcgtcttca ggagctgggc ctccagtgcg gcgcgatgtc 120
aggcgcggtg acagctctgt gagtccgagg ccgcggccgt ggcgctgggc ggctgcgggg 180

```

```

cctgaccggg cccgtcatgg tgcggccacg acgccatcgc ggggcaggaa ggccaggggt 240
gctgagttct tcacctcctt ttagactgag atctgccaa gtttccggca ttgctcttga 300
ggatctcaga agggctctta agacaagact gcaaatgggtg tgtgtatttg tcatgaaccg 360
aatgaattcc cagaacagt gtttcaactca ggcgaggcga atggctcttg ggattgttat 420
tcttctgctt gttgatgtga tatgggttgc ttcctctgaa cttacttcgt atgtttttac 480
ccagtacaac aaaccattct tcagcacctt tgcaaaaaa tctatgtttg tttgtacct 540
tttgggcttt attatttgga agccatggag acaacagtgt acaagaggac ttcgcggaaa 600
gcatgctgct ttttttcag atgctgaagg ttactttgct gcttgacaa cagatacaac 660
tatgaatagt tctttgagt aacctctgta tgtgcctgtg aaattccatg atcttccaag 720
tgaaaaacct gagagcaca acattgatac tgaaaaaacc cccaaaagt ctcgtgtgag 780
gttcagtaat atcatggaga ttcgacagct tccgtcaagt catgcattgg aagcaaatgt 840
gtctcgcatg tcatactctg tgaaagaaca agaattcata ctgaaaactg tggggaaact 900
tactgcaact caagtgcga aaattagctt ttttttttgc tttgtgtggt ttttggcaaa 960
tttgcataat caagaagcac tttcagacac acaagttgct atagttaata ttttatcttc 1020
aaactccgga ctttttacct taactcttgc tgcagtattt ccaagtaaca gtggagatag 1080
atttaccctt tctaaactat tagctgtaat ttttaagcatt ggaggcgttg tactggtaaa 1140
cctggcaggg tctgaaaaac ctgctggaag agacacagta ggttccattt ggtctcttgc 1200
tggagccatg ctctatgctg tctatattgt tatgattaag agaaaagtag atagagaaga 1260
caagtgggat attccaatgt tctttgggtt tgtagggttg tttaatctgc tgctcttatg 1320
gccaggtttc tttttacttc attatactgg atttgaggac ttcgagtttc ccaataaagt 1380
agtattaatg tgcattatca ttaatggcct tattggaaca gtactctcag agttcctgtg 1440
gttgtggggc tgctttctta cctcatcatt gataggcaca cttgcactaa gccttacaat 1500
acctctgtcc ataatactg acatgtgtat gcaaaaagggtg cagttttctt gggtattttt 1560
tgcaggagct atccctgtat ttttttcatt ttttattgta actctcctat gccattataa 1620
taattgggat cctgtgatgg tgggaatcag aagaatattt gcttttatat gcagaaaaa 1680
tcgaattcag agagttccag aagacagcga acagtgtgag agtctcattt ctatgcacag 1740
tgtttctcag gaggatggag ctagttagct gtctgttgtc tgtagcccag cttgataatg 1800
gaactataca gcgaagagac aatctctggc aagtttttgt agaaaaaatg tttcagtgcc 1860
tagtctgaaa aataacagtt tgagttcttt gaaactctaa aatataattt tctcatacct 1920
gttttcttca ttttcataat gaagcacttt gctatgtagc tgtgtacata tcaactacag 1980
tataggaagt ttcagtctac agtccatcca aaggaccaac ctgccttaca catctcaagg 2040
aattcagctg ttgaaatcat ttgaaactaa caaggaataa atcctaattg tctgggactt 2100
tattttcaca tgttaaatgc tggaatataat tatgaaaatg ttttcaagaa atcacttaag 2160
tgttcataga ccagtatttc tgacaggtaa aatgctaaaa taagctacct gtaataagtg 2220
tggattatat ttttgggttt tgtagaatat tgcaaatata ccacacaaaa aatgtttaat 2280
ttatgcaaca agcatgtttg tgcaaatttc atgggacttt aaaaagaata agtatttgag 2340
aaaatatctg gttcacttac actacattta ctgtattatt cttttatagc attagtgcc 2400
ttgtatttta aatctgtgac aaaccatggc aaatttttaa aggggaagta ttattataaa 2460
atgaagaaat atgtatttct aaaggctata ttgctgtaaa cttaattgat aaagctctgt 2520
ttaatttaga gttttgaaga aatagtctcc cttcaattaa gaaattttca taatggaatg 2580
atttaaatg aagtgacaaa gagtattatt aaaatacaat gtttatacgt gaaaaaaaaa 2640
aaaaaa

```

<210> 79

<211> 1749

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 1483702CB1

<400> 79

```

tcagcctggc tggcagcagc cttggactcc gcccgaggag ccctgggcct gttgaccac 60
cagcttagga gcaccacca agctctgggt caacgtggag gtaccaggcc accatgctca 120
gtctcaagct gccccaactt cttcaagtcc accagggtccc ccgggtgttc tgggaagatg 180
gcatcatgtc tggctaccgc cgcccacca gctcggtttt ggactgtgtc ctcagctcct 240
tccagatgac caacgagacg gtcaacatct ggactcactt cctgccacc tggtaacttc 300
tgtggcggct cctggcgctg gcgggcggcc ccggcttccg tgcggagccg taccactggc 360
cgctgctggt cttcctgctg ccgcctgccc tctacccctt cgcgtcgtgc tgcgcgcaca 420
ccttcagctc catgtcgccc cgcagcgccc acatctgcta cttcctcgac tacggcgccg 480
tcagcctcta cagctgggtt tcctggagct ggaaagccct gggctcagta aggtcctccg 540
cacaggagcc ttcgcctatc cattcctggt cgacaacctc ccactctttt atcggctcgg 600

```



```

gctgtgctgg ggcagggggc acggctgtgg gcaggaggcc ctgagcacca gccatggcta 660
ccatctcttc tgcgcgctgc tcaactggctt cctcttcgcc tcccacctgc ctgaaaggct 720
ggcaccagga cgcttttgatt acatcgccca cagccaccag ttattccaca tctgtgcagt 780
gctgggcacc cacttccagc tggaggcagt gctggctgat atgggatcac gcagagcctg 840
gctggccaca caggaacctg ccctgggcct ggcaggcaca gtggccacac tggctctggc 900
tgcagetggg aaactactca ttattgctgc tttcacagcc accctgcttc gggccccag 960
tacatgccct ctgctgcagg gtggccact ggaggggggt acccaggcca aacaacagt 1020
aggccccatc cctgacctg tcttgagggt ggcagaggcc agggccaggt gctgacgagg 1080
agccagatt tgggcctaatt cagggtgggga cgcattctag cctggaacca acaggggctg 1140
aggagagagg gcacaggaga gagggcagag aagaggagggt gtgtctagggt ggactggcag 1200
agtgtgagag ggacctgtgag ggggctcttg atgggagtggt aagaagtgtt gagggtctga 1260
gaggggagat gcatgcgtgt ccaggctgaa gatgccccta tattctgtca aaggttggcg 1320
gggggagggt tgggggtcct tcatctggc tccgtttctg gtgcttctgg aagtctctgc 1380
tcagcacagg gaagaactaa cagactaac ctaggcctac cctgaatgct tcttgctaac 1440
cagggccgaga ggccacacac ttgccccccc atccccacaa accaggtaat gccagtttgc 1500
cagcagctat ttgcctatag agatgagttc gtcctgggtc taactgtgtg ctcaagggtg 1560
ccaggctttt gggggtgggc ctatctgggt gcattatgga tgggttgggt gattgagggt 1620
tggggaggag ggtccctagg tagagggggt atccctagtt agactttggg aagccacctt 1680
caatgttttc tggaacaagg caggtacaaa taaaaaata aaactttgga aagcacaaaa 1740
aaaaaaaaa

```

<210> 80

<211> 2339

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 1519324CB1

<400> 80

```

ctcttcatta gttgaaagg atagtaatac ctacttgag gttgttgtca tctgagttga 60
gcactgggtca cattgaagg gctgggtaag tggtagctct tgttgettcc cgttcagcgt 120
cacatctgca gtggagcctg aaaaggctcc acattagggt acctgtgcac agccatggct 180
ggaatgatga aggggatacg ctggagttgc cctgccatcg cctccatcag ccagacgagg 240
tcctcacagg agaaggacag ctcttcccca ccctgggatc tcaggagggc agccacggag 300
tggggaggcc ccagatgcgc tgtgcccagg ccaggtccga ggccaaagtt ctccctgcca 360
tccttggtgc cgtcctgccc ctctctcctt catgcctggg cctgcaggcc caccacagcc 420
accactgagt ccactcggag tgcctgtgt tcttgagaa ggcattccag ggttgaatct 480
tgtccagcc tcagcctggg acacctaggt ggagagagtg gtctccgctc tgaattggat 540
ccaggggacc tgggtcatt ctctctggct caccaaccct gcaggcctca tctttcccaa 600
aaccacttt gtcttggtgg gagtgggtcc gcgctgctct gcagcagggg ctggggagtg 660
gacagcatca ggtgggaaag tggagtcac cctcatgttt ctgtaggatt ctcaccgtgg 720
ggctggaaga aaagagcatc gacttgattt ctccaaccac tcatccctct ttttctttct 780
tccaccactc cccacccag ctgtagttaa tttcagtgcc ttacaaatcc taagctcaga 840
gaaagttcca tttccgttcc agaggggaagg gaacctccct aggtccttcc ctggcttgtt 900
ataacgcaaa gcttggttgt ttatgcaact ctatcttaag aactgccag cctcagctga 960
aaacccgaat ctgagaagga attgcgtcat gtaagggaag ctggaattaa gggagctgag 1020
ccagtcattg ttgtggcgtg tgagtcagga gacctagggt tcagccctc tctactgtca 1080
gagagctgtg caacgtgggc aagtcattgt cctctgagct gcagtttctt catctgtcac 1140
atcgctacag acaagacctc cctggaaccc ttctgattgt cttagacact gtggttgcaa 1200
aaccacgga aagcctcatt tgtgtgaaa gtcagaggaa aaatgatcca gtggacactt 1260
ggggattatc tgtcattcaa gatccttctt tcaaccccaa ggtcagctcc catctcattt 1320
ccagaaaggc tcatacctgg cttgcaggga agcatctgtc ttgtcattcc aggtgccaga 1380
atcctctcag agtcattgaa ggggtttcac ccatccacc caaggcttgg cacttgcca 1440
gtgtcttagc agggctctgt gagggtggg ggcattccag cactcagaag gcaaaggaa 1500
caccctaccc atttggcctc tggagggggc agaagaaaga aagaaacctc atcctatatt 1560
ttacaaagca tgtgaattct ggcattagct ctcataggag acccatgtgc ttccttgtct 1620
agtgcacaaac tgatgattct acttgctgta gatgaatggt taacacgagc tagttaaaca 1680
gtgccattgt tttgccagtg aagcctccaa ccctaagcca ctgggacggg ggccagagat 1740
gccagcagcc tctgtgcgcc ttagtcatat aaccacaaatc cagaccttat ccacaacccg 1800
gggcttgaaa aggaaggat tttggaatca caccctccgg ttatgttgct ccagtaaaat 1860
cttgctgga aagaggcagt cttcttagca tggtagctg agttcatggc tttttttt 1920

```

```

agccagtcct gtccctggcc atccatgtga tggtttttga tggagttaaa cttgatgcc 1980
gtgggcagtg catgtggaaa gtatcagagt aagcctctcc cctccagagc cctgagtttc 2040
ttggctgcat gaagggtttc tttagaatca gaattgtagc cagtttcttt ggccagaagg 2100
atgaatactt ggatattact gaaagggagg ggtggagatg ggtgtggcag tgtatggtgt 2160
gtgattttta ttttcttctt tggatcatggg ggccaaggag aaaggcatga atcttccttg 2220
tcaggctctt acagccacag gcactgtgtc tactgtctgg aagacatgtc cccgtggctg 2280
tggggccgct gcttctgttt aaataaaagt ggccctggaag ctggaaaaaa aaaaaaaaa 2339

```

<210> 81

<211> 1006

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 1630169CB1

<400> 81

```

ctggcccaag cgtccctctg tctctctgat ctggcccatc cggtctcgga gggaggcgag 60
ggtgggtgag caaagggatt ggtctctgtg ggtccaggcc cgaaccctg aagacgggct 120
ccgcccccg caccgcctcg cgccccgccc ccggctggag gactctctcc tggaccatcc 180
gaacctagcc tgtcccgccc cgcagcctct atggaggctc ctgccgggccc gtagagccct 240
tcgccccctg gggaccaccc cgtctataag gtccgttttg cctgcagcag cctgagtcgg 300
taatgctggg cactgttcat gggatcgccc ccctatggag ccctgtgtct ataggggact 360
cctacgggtc ctaggggttc gccccgtcca taatgactcc atatacaggg ccttccatcg 420
ctctataggg ctacagccctc ggcttccaga gcctgtcagc agtggccgta cccttcgccg 480
ggactgccgg gtctccggga cctctctgat ctacaaggct atgttatgcc tatagaggtc 540
gcatttgcag ggcctcacc cgggtagagg gtccctctcca ggtttttacg gcctgtcccc 600
gctctaaggg tcagtgcagg aggcggcgat ggccctgggc aaggttcttg ccatggcact 660
ggttttggcc ttggccgtgc tggggtcgct gtccctggg gcccgggcg gggactgcaa 720
ggggcagcgg caggtgctgc gggaggcgcc aggtctctg acgatgggtg cgggcaacta 780
cagcgtcaat ggcaactgcg agtggctcat cgagggtgag tggggccgag tgggtcactc 840
actaattcgc tggtagttca ttcattgtgt cattcatcca ctaccacat tcccagagct 900
cggttctggt ccaggccttt tgctctaaaa tgcctgacac ccagggttgc agccaacctg 960
ggggggatcc actagttcag agcgccgtcc cgcgtggctg cagctt 1006

```

<210> 82

<211> 1050

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 1664253CB1

<400> 82

```

cgcttccggc acagcgggaa tccgggtgcc ggttgaggtt gctggtggac ctgctctggt 60
ggtcttggat gaggcccat gagcgcgcg cccctggtgg gctacagcag cagcggctcc 120
gaggatgagt ccgaggacgg gatgcccacc aggcggggg atgggagcca ccgtcggtgc 180
cagagcccc ttcccaggca gagatttcca gtacctgaca gtgtgctgaa catgttccc 240
ggcaccgagg aggggcctga agatgacagc aaaaaacac ggggacgggt gcgcacctc 300
ccccacgagc gaggaactg ggccaccac gtctatgtac catatgaagc caaggaggag 360
ttcctggatc tgcttgatgt gttgctgccc catgccaga catatgtccc ccggctggta 420
aggatgaagg tgttccacct cagcctgtcc cagagtgtgg ttctgcgcca cactggatc 480
ctccccttcg tgcaggctct gaaagcccg atgacctct tccacagatt cttctttact 540
gcccaaccag taaagattta caccaatcaa gagaaaacca ggaccttat tgggcttgag 600
gtcacttcag ggcattgccc gttcctggac ctggtttcag aggtggacag agtcattggag 660
gaattcaacc tcaccacttt ctaccaggat ccttctttcc acctcagcct ggctggtgt 720
gtgggtgatg cactgtctca gctggagggg cagtgcctgc aggaactaca ggcaatcgtg 780
gatgggtttg aagatgctga ggtgctgctg cgcgtgcaca ctgagcaagt ccgctgcaag 840
tctgggaaca agttcttctc gatgcctttg aagtgagcac cagaggcctt cctcctccag 900
ggccctctgc agaccagggt gagatggagg aacctgctaa aatcgatgga gatgcttcta 960
gcctcccagt aggaggcccc agccatgcct tcaacctggc agggaggtgta gccactcctc 1020

```

atcctccctg agtgetgata ttctctctct

1050

<210> 83

<211> 1774

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 1864715CB1

<400> 83

```
gggccttgge cgggtgggtcc cgggcgggac cggggaaggc ggaaggcttc ggcagagctg 60
cgccgcgag gctgagcggg ccttctcgc tgcggccgcc caggtgcccg cgccgtggc 120
gctatggagg cggcgctgct ggggctgtgt aactggagca cgctgggctg gtgcgccgcg 180
ctgaagctgc cgcagatctc cgctgtgcta gcggcgcgca gcgcgcgggg cctcagcctt 240
ccgagtttac ttctggagct ggcaggattc ctgggtgttc tgcggtagca gtgttactat 300
gggtatccgc cgctgacctc cctggagtag cccatcctca tcgcgcaaga tgcatcctc 360
ctgctctgta tctttcattt taacgggaac gtgaagcagg ccactcctta catcgctgta 420
ttgggtgtctt cttgggtcat ccttgccctg cagaagtgga tcatagacct ggccatgaat 480
ctatgtactt tcatcagcgc ggccagtaag ttgacacagc tccagtgtct gtggaagacg 540
agagactcag gaactgtgag tgcgctgact tggagcctct cttcctatac ctgtgcaaca 600
agaataatca caaccttaat gaccaccaat gattttaca ttcttctacg ttttgtgac 660
atgctggcctt taaatatatg ggtaacagt acagtacttc gctaccggaa gaccgctata 720
aaggctgaat gatggatata ttattccttc acacagtggg ttttgagtaa ctgaaccaa 780
ggaaaaagaa gctctttgct aaattaaggt cttttataaa tttagtaa atcagtttataa 840
tctttaaagc caaagggttt tttagacttg aaagaaagag ccacttaa tcttggttaa 900
aaataccaat ttgcctcctc cttcctcact tcgttaggtt atggtagtgc tcagacatct 960
gcagtgttga gaaggccagt cactgttgga agtcatccaa gaagccattt ttgaggccat 1020
tttgagcctt actcttaagt tctctatgaa gaactacatt gatgtgttg ctttcagaat 1080
cttttaggaa ataatcctc tccaggacaa aaatgaacat gaatggagtg gcattttgtt 1140
ccaagtcaga ggtgggcacc tataataaat gactagggtt cactttctgg gactgatgtt 1200
taattgtaac acagatacaa cagggtggcc ttgtgtata taatacggta ttatacctgc 1260
atgtgctcta gcaaggatac caaggcaagc atacatgtag ctggcttgag tttgtaccaa 1320
aacagtcctt caactttgca ctgtgcctta agtaattact aacaaaaggt actaggatta 1380
gctgcaatct ctactttcga tgaggaaatc ccagtaagct ttctgattca agtacaatgc 1440
tgccattttt taaagggcca caactataga attaccactg ttggaatttg gtacaaaata 1500
tgttttgtct attgaaaaca tacacggtaa atgggtgtgt taggtagggt ctgtccagtt 1560
cttagggact tttttcacat tatagcattt ttaccctaaa catgatgttg agattattat 1620
atactgtatt ttcttctaaa ttaaccctaa tgtttaaaaa ctcactttcc ccctttaatt 1680
gaaggcattg ttttggttaga tgcagtaatg atgtttacca gagattattg tttcctatgc 1740
aaaataaatt ttcatatttt gaaaaaaaaa aaaa 1774
```

<210> 84

<211> 2608

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 1929395CB1

<400> 84

```
cgtacggctc gactctagct tcagactcat gtcatttcca gtctgtcagc gttagacatg 60
gtcctgttcc agtttccaga cattcatttc ttactataga gaagagtact cccctggcgt 120
cttaacctat ggaaaacatg cacggatagg atatatattga ttgcctcctc tcccttca 180
gtatatgtat tattaatatt attattatta ttattattat tagttcatca gtttgcgtgt 240
ctctgcagtg agcagaatca aatgggcaat atttgcctg ggagacctgt gccgcaccca 300
gggtccccgtg ttaacgtgtg cctgcggttg tggttggcac cctcggttg tagctcttct 360
actgtaatga gacaagcctt tcttctgtca ctgcagaatt tagaaggggc cgtgagcagt 420
cttcccaagc atggtccagg agcagctcag agaggggtgga gtgaggatgc atgcatccag 480
ggaacaggca tcaagaagcc agggccgtgc ctggagcact gcagagatga ctttgagaa 540
agcaaacgag tgaccacagt accaggcaca ttactcgtga gcgaggtatt cccagtcctc 600
```

```

ctgctaagcc taatccaagc cttaccacgc tgtgctactg tcatttgcaa agcaaggaaa 660
tactactact ggtaataaaa ctacacatgc aagatgtcag ataagaaata ggttttatatt 720
taccttcctg cgatgtgggg ttggtgcttg aagataaatg tgctgtttct agtttcttaa 780
atagcccaca cccctggatg caaaaaggca ggggtatttt taccatgag catcctgtgc 840
aggcaataga ctccctcac tgtctccgcc tgcaggagg agaatgtacc ccactgcacc 900
tgagctcctg gtccctcagc caagacctca ggggtcccca gccagcctgc tgctggggac 960
ccctgtgctt gctgcagtgt atggagcctc ttgccttccc ctggggaggc acccctgtac 1020
cccagcttcc ttcccttggc cttttctggc ccagtgctc ctcccttaca tagacttggt 1080
cacacagaaa cgtgcacgcc cctttttctc cgccacttca ccagtttctg aaatccaacc 1140
tcccagactt cacaggaaga tagatattct tgagataatg aaaagtgata tcttcgcata 1200
cgaaaggaaa aaaggttgag gtatatatga tttttaactg tattaggggt gtatgaacca 1260
gtttaaaaac gaggttttat ttactgtaga gatgaatgca aatcagaacc aatgatccct 1320
tggcctactt agttaaaacc agttcataca tcccttaggg tttttattat tattattatt 1380
attattacag ttgttattgt tgtttttgtt gttattatta tttgggggtt cttgtgtttt 1440
ttctttgcga ctctccacac taaacttgca atattgtggg gagaagctgt gactaaactc 1500
tacgtgcgg tgagatgtag cagtaatcag ctcccagcga cgtgtgtagc tggggctgcc 1560
gctcgcaata atcactattg atttaaagct ttacttagcc ttgatctgta ccctcgtagt 1620
caataaggtc ttgccacatt ttattagtga ggtggagaaa cgtattattt gtttgttgtt 1680
tttgcccttc cccaccccc caatattaaa ctgtgaaatt tgtgatttgt ttaaactctg 1740
ggatgaatcat agcttagttt gcatgtccag ctaatttggt tctatacatt ttgtttgatt 1800
ctctttctcc ttctctcagg gcttttacia aaaaatatat atatatggat cttctgaaaa 1860
gttttttgag gtgcaagttt ttctcttgtt ttttttttct ttttttttct cattgattaa 1920
tggacatgat gctgagattc aatcactaca tgaaacacct ggctgtgaaa acaaaacaac 1980
ccagagggct gtgttccaag cagcgtggg gaagctacgt aacagtcgga tgccagtttt 2040
ggaagattca ccatgcgttc tgacctctg ttcgtctctt tctctcctc tttcttcaag 2100
aaggaaattg atcctagtga ttccagccca tgcattaaac aggaacaat aataaatttg 2160
tagaattcat atttttctaa agggaaacta aaaactgctg ctacatgta tgtacaaaac 2220
tggtttatgc cacatgaaca gagaatcaca agtttggtt ttgtactttt tgttctctt 2280
tgtattcagt tgtatagaac ttccaaattc agaagagaa gaaagctgtt ctgtatcaaa 2340
ccatttaagc aataaatgtt atattttaaa agctgcagaa tactggtaga tgctgccatt 2400
gttgatatgg ttcttgcaa acccatgtga tttgagagct tctcagacta tccacctttg 2460
ggtcgctttg ctgttcgtga tatgagacag acagttgcgg tgggtgtcat caaagcagtg 2520
gacaagaagg ctgctggagc tggcaaggtc accaagtctg ccagaaaagc tcagaaggct 2580
aatgaatat tatccctagt acctgccc 2608

```

<210> 85

<211> 1336

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 1987737CB1

<400> 85

```

cccacgcgtc cggtgacttt caaatccctc ttgcagtgaa aagaaacagc aaacacttaa 60
gattcagcat ctgttctcca gttgactga ggaatgcact gtctgcagc accagctctg 120
cagagccctt gcccagact ctttgcggtt ttatttatat gtatttccat atttcattcc 180
tgtgtgtcac tgctgcattg gtgtggcagc aagtgaccaa atgctacagg tcttactatg 240
gacaccaggt caggtgcaac cacacaaaac aaagccagtt ccatgagctg cctatgatat 300
gcattgcgga agtaacattt taccagggtt gtgccattgc agtgatataa atatatattt 360
ttcttagact aaatatgagc tgactatctc ttttgatgtg tgtacatagg tgtgagtgtg 420
tctgtatgcg tgccgtgtctg tgtgcggtgt tgtgtatgtg cgtagcctca tgcttaggac 480
taccatgaa tgttgtggaa tgctacacct ggagagttct ggttttccac cagtttcaag 540
atgaagaact acatgataca gtggacctgg agaccatccc cttggaaga caaccagag 600
atgttcagca tctgtatct acacgcattc tgtatctaca cgtgtatttt gtagctgtca 660
cactaacctt aataagaatt ctacagcttt ggacagaggc attttcacct taatggtgaa 720
gtaatttaaa atataaatcc attcaggtga caaccatca tcaaaattac aaattttctg 780
attgaactca tctgaatcat cagttccttg atggagagag agaaggagat ggaatgtgtc 840
tggttaacccc aaatggagta caagtagcct ttgttttctt gcataaatgg actgttgtaa 900
tgcgaacgaa tatatgcaat tcatatactt ttggagatga acgtagatat gtgtgtcagc 960
tttgagatgg ttgttctgtg attaatactt tgtctcccaa tatcacagaa aaatacatgc 1020
cagtgactct tgagggttaag gtagttggga tgaaatggcc tcaggcaatt tcacattccc 1080

```

taattacctg gaaagttcta cagtaattaa tatgcagcta actcctgttg ccctcacaag 1140
agcatcagcc ttctagaatc ggagctccgg agtgtgaaga ttcagtattg atatgatatg 1200
tataccaaac tccagccaac ttactgccat ttttcataat ctgagtggtg gccttgctta 1260
tcctaagctg tgggtgcaga aaccgtggcc atttatataa gctataacat caaatcaggg 1320
aaaaaaaaa aaaaaa 1336

<210> 86

<211> 3062

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 2122866CB1

<400> 86

gtccccgccag ctgtgaggag ccccgagcac ggctgtcgcg gcctgaggca gagcgcctct 60
gctctggccc tgtccctggc tccctcatta ccgtcgccag cggagcgaac ccgagagcgt 120
ggagcgggct gggccagctg ttgagtggaa tgtcatggcc agctcctcgg acagtgaaga 180
tgacagtttc atggctgtgg accaggaaga aactgtgctg gaagggacaa tggatcaaga 240
tgaggagccc caccagtat tggaggctga ggagactaga cataataggt ccatgtcggg 300
gctgccagaa gaggttttgg agtatatcct gtcctttctc tcaccgtatc aggaacacaa 360
aactcgggcc ctgtgtctga aacagtggta tgcacttatc aaaggtgtag cccatcagtg 420
ttatcatggg ttcattgaagg ctgtccagga aggaacatt cagtgggaga gccgtacct 480
tccttatcct ggaaccccaa tcaactcagc cttctcgac agtgcatgct attatgatgc 540
taatcagctc atgtatgtgt ttggaggctg taccagagc agctgcaatg ctgctttcaa 600
tgacctctgg agacttgacc taaacagcaa agagtggatc cgacctttgg cttcagggtc 660
ctatccttcc cccaaagctg gagcaactct ggtcgtgtac aaggacttgc tagtgctgtt 720
tggtggctgg acgcgcccaa gcccttatcc cctacaccag ccagagagat tctttgatga 780
aatacacact tactcaccct ctaaaaattg gtggaactgc attgtgacaa cccatggggc 840
acctcccatg gctggccact cctcctgtgt gatagatgat aaaaatgattg tctttggttg 900
ctcttttaga tccccgcaaa tgagcaatga tgtctgggtc cttgaccttg agcagtgggc 960
tggttccaag cgaacatct ctggccccag tccctcatct cgagggtggc aatctcagat 1020
tgtcatagat gatgcaacta tcttaatcct cggagggtgt ggcggtecca atgctctatt 1080
caaggatgct tgggtgttgc acatgcattc tggctccttg gcctggcagc cactcaaggt 1140
agaaaatgaa gagcatgggg cccagaact gtgggtccat ccagcttgcc ggggtgggaa 1200
gtgtgtgggt gtcttcagcc aggtccttag tgggagagcc ccactcagcc ccagtttgaa 1260
ctctcgccca tccactatca gtgccactcc tccagctctc gttcctgaaa cccgagagta 1320
ccgctctcag tctccagtaa gaagcatgga tgaagctcct tgtgttaacg gccgctgggg 1380
aacactgaga cccagggctc aaaggcagac tccctcaggt tccccgggag ggagcctttc 1440
cccagccaga ggagacggct cctcctatcct caatgggtgg agtttgtctc caggaaacggc 1500
agctgtgggt ggctcttctt tggacagtcc tgtacaggcc atatctccaa gtactccatc 1560
tgctcctgaa ggatacgacc tgaaaatagg actttctttg gcccccgac gaggatcact 1620
accagatcag aaagatctga gattaggatc catagatctg aattgggagc tgaaacccgc 1680
ttccagtagt aatcccatgg atggcatgga caataggaca gttgggggaa gtatgagaca 1740
ccctcctgaa cagacaaatg gtgtgcatac cccacctcac gtggccagtg cccttgcaag 1800
ggcgtctcct ccaggtgccc tgcgtcggag tctggaagcc atcaaaagcg tgcctccaa 1860
aggccctcct gccctctgcag cactaagctc tctccttggg tcttctccag gctctcctgg 1920
gagccagagt ttgagcagtg gagaacagct gcccatccct cgcccagggc ctgcccagg 1980
agatggacat tccttacctc ccattgctcg ccgctgggc caccacctc cacagctcct 2040
aaatgttggc aaaccttat accagagtat gaactgcaag cccatgcaga tgtactgtgt 2100
ggacattaaa gacaccaagg agaaggggag ggtcaaatgg aaagtattta atagcagttc 2160
tgtggttggg cctcctgaaa ccagcctgca taccgtggta caaggcaggg gtgaactcat 2220
catatttggg ggactcatgg acaagaaaca gaatgtgaag tactatccaa aaacaaacgc 2280
cttgactttt gtacgagcaa agagataatg tgttctaaac ccctttcctt ttctgtggct 2340
tttaatttgg aattttccag tgtgtaagca tttggactga gaattgggaa aacaaaatta 2400
ctcccagaag ccaaaactct ttaattccca accgaagtca ctccaggctg ggatcaaatc 2460
tccattaaag aaaaaatta tatataaata tatatatata tattatatag ccaactctgt 2520
tgacaaaaaa agggagagat ttccatcctg gttcagataa agttgttgct gtgttttaac 2580
aggggctggg ctgccttttt ctaccttgct ggtaactaga ccaagaagtt agagaataga 2640
ctaaccatcag taacttccca aaagaaactg aagagccccc tgtaaatctt tatgtggcct 2700
tcttgaggtt aaaaaatgaa agggcatatg taagttgcaa aggtggaggg ttttagactc 2760
tcatgcttca ggtgctgtcg gggtaaaagt aactgttttt ccccttctct taaaaccaca 2820

gaggacctgt gacagctctg cagaaatgcc agtgcctggc cccctcttgc cttttatggc 2880
 tgaggaaagt taccoaacia aggattttat tccacatttg tgtgccgggt cattgtgaaa 2940
 taatgtttat gcagccaaca tctgaccgcc tagtagtgc cattggtctt tggagtgcctt 3000
 cttgtgtgtc tcagaaaaca ttttgtgtct gattgtggaa atttctgaca atcaatttct 3060
 tc 3062

<210> 87

<211> 2543

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 2123981CB1

<400> 87

gcacctgcga gcctggaggt ggcgggtcgt gcgaaccgg cggcgcggcc aaacctgtgc 60
 tggctgcctg ggccttcttg gcctcctggg ccagaagctg gcgctggggg cggcgccggg 120
 gtgggaaagc ggcgcgcggg tggggcctcc tggacgcgtc ggcggaggtg gcttcgtccc 180
 gcggagtcca ggcttcaggc tttcaccagt tctcaggatg cccataggga tgggtgaagc 240
 ctgcctggcc tgtgtgtgctt cccagtggcc gtcatctcat tagggcccca cagtggcatt 300
 aggatgcacc tctcggcggt gttcaacgcc ctccctgggt cggtgctggc agcggctcctg 360
 tggaagcatg tgcggctgcg tgagcatgca gccacactgg aggaggagct ggccctcagc 420
 cgacaggcca cagagccagc cccagcactg aggatcgact acccgaaggc actgcagatc 480
 ctgatggagg gcggcacaca catggtgtgc acgggcccga cgcacacaga ccgcatctgc 540
 cgcttcaagt ggctctgcta ctccaacgag gctgaggagt tcatcttctt ccatggcaac 600
 acctctgtca tgcctgccaa cctgggctcc cggcgcttcc agccagccct gctcgacctc 660
 tccaccgtgg aggaccacaa cactcagtac ttcaacttcg tggagctgcc tgcctgctgc 720
 ctgcgcttca tgcaccaagg ggtgttctgt ccagacgtgg ccctcatcgc caaccgcttc 780
 aaccccgaca acctcatgca cgtcttctcat gacgacctgc tgccactctt ctacaccctg 840
 cggcagtttc cgggcctggc ccacgaggca cggctcttct tcatggaggg ctggggcgag 900
 ggtgcacact tgcacctcta caagctgctc agccccaagc agcctctcct gcgggcacag 960
 ctgaagacct tgggcggct gctgtgcttc tcccatgctt ttgtgggctt ctccaagatc 1020
 actacctggt accagtatgg ctttgtgcag cccaggggcc cgaaggccaa catcctcgtc 1080
 tcaggcaatg agatccggca gtttgacagg ttcatgacag aaaagctgaa cgtgagccac 1140
 acaggagtcc ccctaggcga ggagtacatt ctggtcttta gccgaacca gaacagactc 1200
 attctgaatg aggcagagct gctgctggca ctggcccagg agttccagat gaagacagat 1260
 acagtgtccc tggaggacca caccttctgt gatgtcgtgc ggctggtcag caatgcctcc 1320
 atgctggtca gcatgcatgg ggcccagctg gtcaccaccc tcttctctgc ccgtggggca 1380
 actgtggtag agctcttccc atatgctgtc aatcccgaac actacactcc ctataagacg 1440
 ctggccatgc tgcctggcat ggacctccag tatgtagcct ggccgaacat gatgccagag 1500
 aacacagtca cacacctga gggccctgg gatcaggggg gcatacccca tctggaccgg 1560
 gctgagcaag ccggtatcct gcaaagccgt gaggtcccac ggcatctctg ttgccgggaa 1620
 cccgagtggc tcttccgaat ctaccaggac accaaggtgg acatcccac cctcattcaa 1680
 accatacggc gcgtggtgaa gggccggcca ggaccacgga agcagaagtg gacagtcggc 1740
 ctatatccag gcaaggtgcy ggaggcacgg tgccaggcgt cagtgcattg cgcctccgag 1800
 gcccgctca ctgtctcctg gcagatccca tggaaacctta aatacctgaa ggtgaggagg 1860
 gtgaagtacg aggtgtggct gcaggagcag ggggagaaca cctacgtgcc ttacatcctg 1920
 gctctgcaga accacacctt cactgagaac atcaagccct tcaccaccta cctggtgtgg 1980
 gtccgctgca tcttcaacaa gatcctcctg ggacctttg cagatgtgct ggtgtgcaac 2040
 acgtagcgag caggccacag cctggcctcg ggaaggtggc tctgcagtt cagcgtccct 2100
 gggccatta atcccactgt ggagacttct gggaactatt tattgagcag gcctgtgcct 2160
 ccacatcatc ttgttgtctc tgggggtgtg tgccacagca ctctctttg ccctagagat 2220
 aagggacctg acttcccctt ctcccatcct gaacatttgt acccctggag aagttcctta 2280
 gcaggaggga ggaagaggag agggaggaagc aaagaatcac aaggaacctc tggctagggtg 2340
 atcctgatgt ttctactga gtttttcttg tatccagatt tctggaaacc gagtaatcat 2400
 gtactgtttg attgggtggt tcatctgctt ccactccagt gaaatttacc tgtagcccag 2460
 tgaagggtgt gtttgggaaca ttcaaaatat cacttcaaca tcttatcaaa ataaaaatta 2520
 ttaatgagaa aaaaaaaaaa aaa 2543

<210> 88

<211> 873

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 2200177CB1

<400> 88

```
tacatgcggtt aatcactttt gcacacctgg ataatatctg cttgctttat gatctgtttg 60
aagaaataaa cgttacagta acaagaaagt ctgggaaggt ggtacctaaa aagaacaaa 120
acagaaaaca aaaaacaaaa acctcagaaa cttgttttgt aagtaacatc cttttatata 180
acattgataa tctctattct tgtctgtctc tgtgtcacaa aagaagataa ttgtagagct 240
tgaatgatat atcaaaagatt atctagtaca tttttcacag gtacagaaac agatgctcag 300
tgaggataaa ttatttcaaa taatacacag tctgttaata acacagctag catcagaaac 360
taagatctct gcaaccatct gtctgccttt actcttccat tgcttgtttc tgttagtttt 420
gagctttcct atcacactct gtataaggca ctctggcccc tatcatattt atcctctgtt 480
acagggtgtc aatctcattt tcctccagac ccatttcttc agctacattg ctggcattat 540
gcaaaaagctc ctttccaatg ttgtgcactc tcaaaaaatt catccagaaa tactaagggtt 600
tgggaaggtc tgtgcccata gcactataag caaaaagttc aaagaagaaa aatacaaaaac 660
tcctcataca ataagctcaa tatcacaat tcatgaaacg gcaactataa agagcaaagt 720
attcagaaaa ttaagtacat atttcagcat tgtcttaaag ctaaaagaaa ttaaaatagc 780
tggtttttaa tatctatgga gttcaaaacta gattagtgtt gaactcttca ccttatcagt 840
atatatataa taaaactttg gttatgaatt tat 873
```

<210> 89

<211> 2134

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 2319255CB1

<220>

<221> unsure

<222> 713

<223> a, t, c, g, or other

<400> 89

```
ctctgcaggc acccaccggt tctgttttct tctgttatcc ttccagagat tttctatgcc 60
taaacaccat ttgaagccat gaatgtatag tgatgagggg ttttttacag ttgcacatga 120
ctgggaatag aaatgcaatt agatattgag aatattgatg agccaatagg agctgatggg 180
ttagaaaaat aaggaggtga agagaatcca ggattttaag gtttaaatga agttagagac 240
tagatttgga gcgagtaaga gaagatacag aattctattt aaaagattca gaactgggaa 300
aattagaact agcattagtt ttaactgaaa tcatataatg catgtgttca tapcaagtag 360
aacaanaatg gccttttttt ctctcatatt taaactccac attatctact gtaatgttcc 420
acaaagagca catatttagt aaatattgcc tttagtgtcc aattatgttt gagcttgctg 480
tagagttagt atttctaatt gctaactcat tttcattttt aaaataatct ataattgtat 540
ttgcaagaat ttcaatccat agcctcctag aaaaaaatc aataccatag gcagaaaaat 600
aatatttcag gtaatagagc atatcactgc atacaaatcc attcttactt caatcaaaaa 660
agaatatgat gcctttattg agacaataaa gaaagaccga agaactacat ttngtcttca 720
tggaanaact aaagggttgg cagcagagcc tacagctttg gtatattaca ggaaaagaac 780
aatccaactt gaagcaaagc atgactcttc aagaatccat gaatctagat gttctcacta 840
aatacatgaa acatcttgaa gataaatatg cagaaattaa acaagctatg ttgataaaat 900
atgtgccagc tcagaggaag gctgatttag atgaagaaat gatttgttta ttaaacgggc 960
gagatgtagc tgaaaatctg aacaaaaaat tgcagttttg tcatcaaaaga ctgcagataa 1020
tttcacaggc acttagttca tgggttaaat ctgatatgag cagcccattt caagactttg 1080
tggagcaaat tcagaaaacc aaatatttac aagggtttaat gagttaatct cacttgggtg 1140
atatgaaaag gcagctgtt atgcagcaaa cagtcctaga agaattcttc gaaacattgg 1200
tacaatgaat acatttaagg ctgttggaat aattagagga aagcctcttc cattactctt 1260
attttttgag gccctcttta tcacaagtca tgcttttcca tgcctgtttg atgcagctct 1320
aaccttgga ggaatcaaat gtggattatc agaaaaacgg ttagatttag ttaccaattg 1380
ggttacacag gaaagactga cttttctga ggaggctggg gatgtgattt gtgattatgg 1440
ggagcaggat acttataaca aggccaagtg cctggcttta gctcagatta tctacagtga 1500
```

```

atgtggcctg cacaagaaag ctattctttg cttgtgtaaa cagggtcaga ctcatagggg 1560
catggagtag atacagcagt tgaaggactt tactaccgat gacctgttgc agctattaat 1620
gtcatgtccc caagttgaat taattcagtgt tctcactaaa gagttgaatg agaaacaacc 1680
atctttatct tttggctctg ctatacttca tctgttctct gcagacatga aaaaagttgg 1740
cattaagcta cttcaagaaa tcaataaagg tgggatagat gcagtagaaa gtcttatgat 1800
aaatgattcc ttttgctcca tagaaaagtg gcaagaagtg gcaaataat gttcacagaa 1860
tggctttgac aaattatcta atgacatcac gtctattctt cgatctcagg ctgcagttac 1920
agaaatttct gaagaggatg acgcagtcac cctaattgaa catgtgtttt ggtagttcta 1980
tatcttaacc agctgaggga gctgtacaa caccttatgt atgctggttg ggaaaaactg 2040
attttaacgt aactaactta taaataaatc tagagtaaga agatctcaga aataaatcac 2100
acttttggtt tgatgacaaa aaaaaaaaaa aaaa 2134

```

<210> 90

<211> 1886

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 2792452CB1

<400> 90

```

tggctgtccc caccagctcc aaagtctcca cctctcaaac accatgtctt ctttgggtgc 60
tgggacacag gctgttcttg gccaggccac agcctcatca tgcattggctc tgcccgggtg 120
ctcacgggct ctgtccacaa caccggggga caggtgctct tatctctccc aggagtcagg 180
atgagacgtg catgcatgta ttgccagctt ctcttgggat cctgggtgct tcctcagccc 240
ttggccgtcc cttgggcttg gctccccacc tctctggcac ccaagactgg tggaaaagga 300
aagccctctc gttctgagtt gagatctcag aacccttggc ctttctgcag ggccctccaa 360
ggcgtctggt ggaccctgca cctgctctgc agcgccccct ccaggccttc ttgcacggga 420
cagtgcaggg gagttctgtc tcagagcgag tccagcaggg cctccacaga gtgcaagggt 480
gcctgccctg cctgtggaag agcaaacctt gtccctgtt tcccagaccc cctactcagt 540
ggcctctgaa tgtecccaat gtgggcccct cctgggtgc cctcttgcct ctccaagtcc 600
gagcccaccc agagcccaag ccctcggagg cccttgccac ctggaaaagt gacactgggg 660
cagggttctc ttctgatgtc tgtgttttgt cttgtggggt ttggggtgct cctgcccctg 720
ataagaaggg gtttcagagc agaaataaag cctcagacag gtgagcctct gtggcacatg 780
gccctctcag ccagccatgc atctgggttc agcccctgct aggacacttg accaggcccc 840
ttaccttctc cgagctggga tcatgtggg aagactgaat gggacgcagc cggtgaaaac 900
gcctggaaga gagtaaacgc cttttctct aagcatattt atttcttta caacagtgc 960
gaccatttgt gagccatctc tctgtctcag gcacggtgct acatgccaac gaaacctgct 1020
cccattgaac cctggccagc cagtgaagaa aggggttggc ctgggagggt ccactttaca 1080
gacaggggca ccaaggggca ggggtggcag agggccaccg gacgttcccc atgaagtagc 1140
agtcccagca tccacaccca gcaggcaca cgctggccc cagcctccct gccagcacgc 1200
ctggcttccc gcctctggaa cttgatctgc tccctcttcc ggacactggg gctcctgcca 1260
agtctgtggc tgggcagcaa ctgctgaaca ttctaagaaa tccctcccag ggttttctca 1320
ggagcccggg tggggcagga agtccccagg ggctgagggg accgtggcgg cagggtggc 1380
ccagagcagc actctcctgg ggcccaggct gttgggccc aggcaggact gtgaggccta 1440
gtgtaggggc tcctgccagt ggccggcacc tactgtggg gctgggggtt ccccagcag 1500
gttgggctcc ccacctgaca cactcacaga ccttgtgcct tggagagcca gtgttcccgg 1560
ggccacatag ctatgccgcc caggggctgg gcctgtccca gctctggtcc cccggcccc 1620
ggctcctggg gctggctccg cgcagcagca ggcgccctcc ggaggacacg atgtgactgg 1680
ctgccgtac gtcgcactca gatgagctct cgccggatcg acctgtgcc gactcctgcc 1740
ggacaggcac aggcaggag tgaaaattat ctaccctttt ttatttctta ataactgaat 1800
gaaaataaac attggtggtt tgacaaataa ctacatattt tcaaacccag ccagtcagg 1860
ggtgagcagtt tccaggtgag ttatgc 1886

```

<210> 91

<211> 928

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 2853088CB1

<400> 91

```
ctacaagaag ctattttggc actaagagag gagagcctgc ctgaggggtga agccaaatgc 60
atagggaagg cagaaccaga gatggtgagc cacagggttcc tgatatcttc tggccctgt 120
gttcacctgt ctgaatctgt ccgctctgga cttttcgatt acttgggcca tagcttcctt 180
tactgttttag gccagctgga gccgggtcaa tcagacactg gatgggtgga ggatctcact 240
gtgctcatgc tttcaactac cctccgtact ttaattgttc tcgatttctc taactggcct 300
tgatgtctct tcttagctcc agactatatg cccacagacg tcccaaaagc acgtctggaa 360
ttgacctcat tattattatt gttattatgt ttgagatgga gtctcgctct tttgcccagg 420
ctggattgca gtggcgcagt cttggctcac tgcaacttcc gactctgggg ttcaagtgat 480
tcttctgcct cagcctcctc tcaagtagct gggagtacag ggcctgcca ccaagccgg 540
gcaaaagagc gagactccat ctcaaaaata ataacaataa taataatgag gtcaattcca 600
gacgtgcttt tgggacgtct gtgggcatat agtctggagc taagaagaga catcaaggcc 660
agttagagaa atcgagaaca attaaagtac ggagggtagt tgaaagcatg agcacagtga 720
gatectccac ccatccagtg tctgattgac ccggctccag ctggcctaaa cagtaaagga 780
agctatggcc caagtaatcg aaaagtccag agcggacaga ttcagacagg tgaacacagg 840
ggccagaaga tatcaggaac ctgtggctca ccatctctgg ttctgccttc cctatgcatt 900
tggttcacc ctcaggcagg ctctcctc 928
```

<210> 92

<211> 962

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 2949004CB1

<400> 92

```
ttggacctag caatgccatg aaattaccgc gagaggaagc aagcgcaagt ttcgtgagga 60
gggcagatct cagcagagag gatttggcgc cctcctctgt ggattctggc caggccgggt 120
tcggcgggtg ctgtgagagc gggtctccca acaccatgcc gtccgccttc tctgtcagct 180
ctttcccgct cagcatccca gccgtgctca cgcagacgga ctggactgag ccctgggtca 240
tggggctggc cactctccac gcgctctgcy tgctcctcac ctgcttgctc tcccgaagct 300
acagactaca gatcgggcac tttctgtgtc tagtcatctt agtctactgt gctgaatata 360
tcaatgagggc ggctgcatg aactggagat tattttcgaa ataccagtat ttcgactcca 420
gggggatggt catttctata gtattttcag cccactgct ggtgaatgcc atgatcattg 480
tggttatgtg ggtatggaag actttgaatg tgatgactga cctgaagaat gcacaagaga 540
gaagaaagga aaagaaaagg agaaggaaag aagactgagg ggcagcagct gcttggagtt 600
tgcgctcctc ccgtccacc agtgcagctc ccagtgcctg agtgtgcgtg gcgtgggcat 660
ccttcagctg gactcatggt ttgaaaaacc gttgttttat ttaaataatc acagtggtag 720
ggcacacact gaagtgggt tttcagccag cactgaatgt atccatcagg acatgcgtct 780
tcaggtgcct gatctttgta gtcaggctgt gggaacggtc tctgcagagc ttcataactg 840
ggaatttgat ttgaagaagt ccatgtcata tgtgtaacta gtactaatta taaatataaa 900
atacacaata taaaatatga aactcaataa taaacagtgc cacctgtaaa aaaaaaaaaa 960
aa 962
```

<210> 93

<211> 2644

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 3011670CB1

<400> 93

```
ccctctcttt cgctgtttga gagtctctcg gctcaaggac cgggaggtaa gaggtttggg 60
actgccccgg caactccagg gtgtctggtc cacgacctat cctaggcgcc atgggtgtga 120
taggtataca gctgggtgtt accatggtga tggccagtgt catgcagaag attatacctc 180
actattctct tgctcgatgg ctactctgta atggcagttt gaggtggtat cagcatccta 240
cagaagaaga ttaagaatt cttgcaggga aacaacaaaa agggaaaacc aaaaaagata 300
ggaaatataa tgggtcacatt gaaagtaagc cattaacat tccaaaggat attgaccttc 360
atctagaaac aaagtcagtt acagaagtgg atacttttagc attgcattac tttccagaat 420
```

```

accagtggct ggtggatttc acagtggctg ctacagtgtg gtatctagta actgaagtct 480
actacaattt tatgaagcct acacaggaaa tgaatatcag cttagtctgg tgccactctg 540
ttttgtcttt tgcaatcaaa gtctctattt cattaactac acactatttt aaagtagaag 600
atggtgggtga aagatctggt tgtgtcacct ttggattttt tttctttgtc aaagcaatgg 660
cagtgttgat tgtaacagaa aattatctgg aatttggact tgaaacaggg tttacaaatt 720
tttcagacag tgcgatgcag tttcttgaag agcaaggttt agaatctcag agtcctgttt 780
caaaacttac tttcaaattt ttcctggcta ttttctgttc attcattggg gcttttttga 840
catttctctg attacgactg gctcaaatgc atctggatgc cctgaatttg gcaacagaaa 900
aaattacaca aactttactt catatcaact tcttggcacc tttatttatg gttttgtctt 960
gggtaaaacc aatcaccaaa gactacatta tgaaccacc actgggcaaa gaaagtatcc 1020
ctttaatgac agaagccaca ttcgatactc tgcgactctg gtttaataatc ctgctgtgtg 1080
ctttgcggtt ggccatgatg cgtagtccac tgcaagctta tttaaattta gcccaaaaat 1140
gtgtggatca gatgaagaaa gaagcggggc gaataagcac gggttgagcta cagaaaatgg 1200
tggctcgagt cttttattat ctttgtgtca ttgcaactgca gtatgtggcg cctctggtaa 1260
tgctgcttca cacaactctg cttttgaaaa cactaggtaa tcattcctgg ggtattttatc 1320
cagaatctat ctctacctta ccagtggata atagtctact gtccaattct gtttactctg 1380
aattaccatc agctgaaggg aaaatgaagg taactgttac acaaataaca gtggcactga 1440
gcagcttaaa aaatattttt actcctcttc tttttcgagg acttctgtct tttctgacct 1500
ggtggattgc tgcttgctc ttttctacaa gccttttttg gcttttctat caccagtatc 1560
tgactgtggc atgaatctca gttaacaaaa aagcatatcc aaatcacctt ttaaattaaa 1620
atatctgtgc ccttaaaggg ctgatgaaaa ccagaagaaa gcaaatacaa tgggaaaaaa 1680
aaaacatata agaattgtct gtattaaatg tttcctctgt attctcaggg tgaattaatg 1740
tagtaaatatt taaaattaca aaatagattg ttaactgtta cactgtggca ttggaatttt 1800
aactctttgt atttactggt atgagagggc tatctacaag ggtaaatatt ctgattacc 1860
tggtttacag aaacctccag cagtctttga aacatctcac atgactctag ttattgattg 1920
cttttaatgg ttttatggta ctgttgatag tcatagtggc tgcctataga acaatcttca 1980
aactgagcca tgctttaggg gagggaaaagg ggctaaagtc tcttctgttg gtaattttat 2040
agttactctt gaaacagtaa aatccaacag aaaggaagag atagctactg tatattacag 2100
taaagaaagc tgcatagtta ttttaaattt aatggagatg aatatgggta aaatatataa 2160
ctactgtctg ttgagaatag caagagtatt gttttaaacc atattccacc caacttgaga 2220
gttcttttaa aatgattggc catatgaaca tttgtaatct tgccattagg tttggacctg 2280
ccatattttt ttttattctg tgatcctaac tagttccttt taataggcta aaatatttat 2340
caatactgat cagactttta agaaattact ttgtaaacct gctgactacc tgtatgtatt 2400
gtatatatat tatatatata atatataata tattgagatt ataaaagatg aaaatattga 2460
atccttataa tatttttaagt tgcagaatgt atgttaaaaa gtgacttgaa tgagatgtat 2520
ttgtatctag aaattttatt tcttttttga atgagattaa aatacatatt gaaagttcaa 2580
caaacagaaa agaaaagaca gtagcagaag tcggtgaagc ggagtgtgcg cgagagtagc 2640
ccct

```

<210> 94

<211> 1875

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 3242083CB1

<400> 94

```

cgctcgccaa ggtcgcgggc cgcttgggga gtcagcagcg cgccaggccc ctteggggccc 60
cacacgcatt aggtgccttc ttgatgggta cggagtgaac gcgggcggcg gcgggaccga 120
ggcagcgccc agtttgtaac cgccgcgcgc cccgtgcccg cgcgcgccac accccagcgc 180
gcttcgggcc ggccacagtg accgcgcgtg cacgtgttcc ggctctcccg cttegcgcgt 240
ccgaacctcc tcttggtcgt cccggcattc gtccacgcgg agccggcttg ggcgggggccc 300
gggaggcggc ggccggagaa gccgcggaga cgcgagcgcc gagcgtcgcg agggagcagg 360
cccgggcagg caagcgggcg cctccgccat gaacccagg ggctgttcc aggaactcaa 420
cccagtaag tttctcatc acactgcct cgtgctcttc tcgggtctgc tgccctccg 480
cctggacggc atcatccaat ggagctactg ggccgtcttt gccccatat ggctgtggaa 540
gcttctagtc gtcgcaggcg cctccgtggg cgcgggcggt tgggcccgc accctcgcta 600
ccgcaccgag ggagaggcct gtgtggagtt caaagccatg ctgatcgctg tgggcatcca 660
cctgctgctg ctcatgttcg aagtcctggt ctgcgacagg gtggagaggg gcaccactt 720
ctggctgctg ctcttcacgc ctctcttctt cgtgtccccc gtgtccgtgg ctgcctgcgt 780
ctggggcttt cgacacgata ggtcgctgga gctggagatc ctgtgctcgg tcaacatcct 840

```

```

gcagttcatc ttcacgcgcc taaagctgga caggattatt cactggccgt ggctgggtgt 900
gtttgtgccc ctgtggatcc tcatgtcggt cctttgcctg gtcgtcctct attacatcgt 960
ctggtccttc ctgttcctgc ggtccctgga tgtgggtgccc gagcagcggg gaacacacgt 1020
gaccatggct atcagttgga taacgattgt cgtgcctctg ctcaactttg aggtcctgct 1080
ggttcacaga ttggatggcc acaatacatt ctctacgtc tccatatttg tccccctttg 1140
gcttttcctta ctaactttaa tggccacaac atttaggcga aaggggggca atcattgggtg 1200
gtttggcatt cgcagagact tctgtcagtt tctgcttgaa attttcccat ttttaagaga 1260
atatgggaac atttcatatg atctccatca cgaagatagt gaagatgctg aagaaacatc 1320
agttccagaa gctccgaaaa ttgctccaat atttggaaaag aaggccagag tagttataac 1380
ccagagccct gggaaatacg tccccccccc tccaagtta aatattgata tgccagatta 1440
aactcctaga gaggaccag gcacacacag actccacttg gccttcgcct cttgttcatt 1500
catcccaaac ctggaaatgg aaacaggctt caaacactcg tctcacgccg tgtttgagat 1560
caccgcctca tcagatgca tcatagatgg aggtgggttc agtatgtggg tgtgtgtggt 1620
gtgtacctgg gtaagagact tgctttccag gttcgcactt tcaggtgtag ctgggggag 1680
taagtcaaat tgttttagta ggtcctcaaa aggaataacc acacagctgt ttgtttaaat 1740
gctactgtac ctatcaaaac tattgtttta aaagtatttt tatacactgc taatctaaaa 1800
ttgtatttca gattgtgcct gtcataacaa tagcaaatgt aaaaagtctt ctttcccacc 1860
aaaaaaaaa aaaaa 1875

```

<210> 95

<211> 2378

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 3363391CB1

<400> 95

```

aaagtgcagc agatgttgcc ttgctagttc aattcactgg gcggccgtaa tatggcggtt 60
gacgcgtgcg gtaaacgacc gtcgtagtag agtcagagca catgacgggc gaggcaagga 120
atgtgttctt ggtctctcta ccatgacctg ggaccgatga tctattactt tcctttgcaa 180
acactagaac tcactgggct tgaagggttt agtatagcat ttctttctcc aatattccta 240
acaattactc ctttctggaa attgggttaac aagaagtgga tgctaaccct gctgaggata 300
atcactattg gcagcatagc ctctctccag gctccaaatg ccaaactctg actgatgggt 360
cttgcgcttg ggtgtgtctt ctactgata gtgcaagctg tgacttggtg gtcaggaagt 420
catttgcataa ggtacctcag aatttgggga ttcatcttag gacagattgt tcttgtgtgt 480
ctacgcataa ggtatacttc actaaaccca atctggagtt atcagatgtc caacaaagt 540
atactgacat taagtgccat agccacactt gatcgattg gcacagatgg tgactgcagt 600
aaacctgaag aaaagaagac tggtaggta gccacgggga tggcctctag acccaactgg 660
ctgctggcag gggctgcttt tggtagcctt gtgttcctca cccactgggt tttcggagaa 720
gtctctctgt ttccagatgg gcagttagtg ggcattccca tccagggcca gatcctaacc 780
catttggagg tgcagtactg ctgtgcttgg caagtggatt gatgcttcca tcttgtttgt 840
ggtttctgtg tactggtttg atctggtggg ttacaggaac agcttcagct gcggggctcc 900
tttacctgca cacatgggca gctgctgtgt ctggctgtgt ctctgctatc tttactgcat 960
ccatgtggcc ccaaacactt ggacacctta ttaactcagg gacaaaccct gggaaaacca 1020
tgaccattgc catgatattt tatcttctag aaatatcttt ctgtgcctgg tgcacagctt 1080
ttaagtttgt cccaggaggt gtctacgcta gagaaagatc agatgtgctt ttggggacaa 1140
tgatgttaat tatcgggctg aatatgctat ttggctctaa gaaaaacctt gacttgcttc 1200
ttcaaacaaa aaacagttct aaagtgtctt tcagaaagag tgaaaaatac atgaaacttt 1260
ttctgtggct gcttgttggg gtgggattgt tgggattagg actacggcat aaagcctatg 1320
agagaaaact gggcaaagtg gcaccaacca aagaggtctc tgctgccatc tggcctttca 1380
ggtttgata tgacaatgaa ggggtgtcta gtctagaaag atcagctcac ctgctcaatg 1440
aaacagggtc agatttcata acaatttttg agagtatgc ttctaagccc tatatgggga 1500
acaatgactt aacctatgtg ctaggggaaa agttgggttt ctatacagac tttggtccaa 1560
gcacaaggta tcacacttgg gggattatgg ctttgtcaag ataccaatt gtgaaatctg 1620
agcatcacct tcttccgtca ccagaggcg agatcgacc agccatcaca tgaccgtta 1680
acatttcggg caagctgggt gattttgtcg tgacacactt tgggaaccac gaagatgacc 1740
tcgacaggaa actgcaggct attgctgttt caaaactact gaaaagtagc tctaatacaag 1800
tgatatttct gggatatatc acttcagcac ctggctccag agattatcta cagctcactg 1860
aacatggcaa tgtgaaggat atcgacagca ctgatcatga cagatggtgt gaatacatta 1920
tgtatcgagg gctgatcagg ttgggttatg caagaatctc ccatgctgaa ctgagtgtt 1980
cagaaattca gatggcaaaa tttaggatcc ctgatgacct cactaattat agagacaacc 2040

```

```

agaaagtggg catagaccac agagaagttt ctgagaaaat tcattttaat cccagatttg 2100
gatcctacaa agaaggacac aattatgaaa acaaccatca ttttcatatg aatactccca 2160
aatacttttt atgaaacatt taaaacaaga agttattggc tgggaaaatc taagaaaaaa 2220
agtatgtaag ataaaaagaa gagattaatg aaagtgggaa aatacacatg aagaacctca 2280
acttaaaaaa cacatgggat ctatgcagtg ggaaattacc tccatttgta aactatgttg 2340
cttaataaaa acatttctct aaaaaaaaaa aaaaaaaa 2378

```

<210> 96

<211> 1597

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 3703614CB1

<400> 96

```

ttccagccct catgaacccc ttcttgggaa acttgccctc agcacccgcc atgggctgtt 60
cagatgcctc cactctgaac cctggaagtg cctcacatgt cagcacctat actgaggact 120
ctggatcagc acaccagagc agggatcaag tcttcctccc tgcattcccg gtacaagtgc 180
gtagggtgtaa agctttgaag gaaaaagatt taattagaac gtctgagtca gactgttact 240
gctacaatca aaattcccaa gtggagtggg aatacatatg gtcaactatg caggtgaaaa 300
ttaccagtc caggcctgttc agaattgtat atatcgaga aagacataat tgccaatatc 360
cagaaaacat tctatctttt atcaaatgtg tgattcataa cttttggata ccaaaggaat 420
ctaaccgaat aaccataatc atcaatccat acaggagagc tgtgtgcttc tctgtggagc 480
ctgtcaagaa gatatttaac tatatgatac atgtgaatcg aaacatcatg gatttcaaac 540
tcttccttgt gtttgtggca ggagtttttc ttttctttta tgcaaggacc ctgagtcaaa 600
gccctacttt ctattactcc tcgggaactg tgctagggtg tctaatagaca ttagtctttg 660
tcttgctgtt ggtgaaaaga ttcatccga agtatagcac cttttgggct ctaatgggtg 720
gttgttgggt tgccctcagtt tatattgtat gccagttgat ggaagatctg aagtggctgt 780
ggatatgaaa caggatatat gtattaggct atgtcttgat agttggattt ttcagctttg 840
ttgtttgtta caagcatggg ccccttgagc acgacaggag cagaagtctt ctgatgtgga 900
tgctgcgact cctctccctg gttctggtct atgtcgtgtg ggctgtgcct cagtttgcct 960
atgcagccat aatcctcctc atgtcctcct ggagtctgca ctaccactg agagcatgca 1020
gttatatgag gtggaaaatg gagcagtggt ttacatcaaa agagctgggtg gtgaaatatc 1080
ttacggaaga cgagtacagg gagcaagctg atgtctgaaac gaacagtgtc ctggaggagc 1140
tacgccgggc ctgccgaaaa cccgactttc cctcatgggt ggtcgtctcc agactccaca 1200
ctcctagcaa ctgatgaaac caaggccctg aaaggttcac caagttgcca gggctcgtgca 1260
gatcaacaaa ctgcagaagc aagactcaac caggtcttct gctgtcaaat cctgtgccct 1320
tcccatggcc agcagagcct cccccagctg gatcagaaac ctttcaggga aatttagaag 1380
gtgctttctt tcatacacac aaaaagtagt gatcacttaa ccttaaaatc ttgtttcaca 1440
aagatgttgt aatattatac ctatacttaa aacgttagat ttctaataaa atttaattaa 1500
tagagctatg aataacttaa tagtaataat gaaagtcata gtaattaata gtaacttaat 1560
aaatagcaat aaaatagcaa tgaaaaaaa aaaaaaa 1597

```

<210> 97

<211> 653

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 4000975CB1

<400> 97

```

aaaacaaaaa aaaactgtgg gctctaccaa tagtcccaaa tgccaagaaa gcttccttca 60
tggagggaaa gcctttttct ctccgtggaa ctccagccac tggccttggc aatgggcagt 120
gctcctggcc tgcaagtctt ctccaagaca aatcctctgt ttcttagccc acccttgaaa 180
agttagggctc tgggcccctc ccccaggag gggttctggc ccaatctgca acgtcaagta 240
cgcgccgtgt ccttgggttg tgaggcagct ggggaggggtg actttggaca gatgtccttg 300
ggctgtgagg cagctgggga ggtgtacttt ggacagatgt ccttgggctg tgaggcagct 360
ggggaggggtg actttggaca ggtgtccccg gcactgtgcc ccagtccagt gcaactaagg 420
gatgggctgt gcctactttg aaggtaacag agtctgcccc gggccacagt ccaagggcga 480

```

ggcccttttg gggcgccctgt catggcatct ggtcatcagc caccttggcc cctgcaagct 540
 ggggccaccc atgggcactc ctgagcaagg cctgggaggg acctgccagt gtcctgggag 600
 ccactctgcc aaggctcagt ggcgatgtca tactttggaa aagaaaaaa aaa 653

<210> 98

<211> 3090

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 4598831CB1

<400> 98

ggctgggccc ggctgggcta cgcgcacggg ctcggccggc gcccctgccg gttgcatacc 60
 ctgtcctgag ggcgcggcac ggagtgcatt cgggcccgtg ccatgacgac cgccatcttg 120
 gagcgccctga gcaccctgtc ggtcagcggg cagcagctgc gccgcctgcc caagatcctg 180
 gaggatgggc ttcccaagat gccttgact gtcccagaaa cggatgtgcc ccagctcttc 240
 cgggagcctt acatccgcac cggctaccgc cccacggggc acgagtggcg ctactacttc 300
 ttcagcctct ttcagaaaca caacgaggtg gtcaacgtct ggacccattt actggcagcc 360
 ctggccgttc tcttgcgatt ctgggccttt gccgaggtg aggccttgcc atgggcgtct 420
 acccactccc tgcctctgct cctcttcac ctgtcgtcaa tcaactacct cacctgcagc 480
 cttctggccc acctgtgca gtccaagtca gagctctccc actacacett ctactttgtg 540
 gactatgttg gcgtgagcgt ttaccaatat ggcagtgctt tggctcattt cttctacagc 600
 tctgaccagg cctggtatga ccggttcttg cttttcttct tgcacgagc tgccttctgt 660
 ggctgggttat cttgtgctg ctgttcttat gccaaatatc gttaccggag gccttatcca 720
 gtcattgagga agatctgtca agtggtgcca gcaggtctgg cttttatcct agacatcagc 780
 cctgtggcac accgtgtggc gctctgtcac ctggctggc gccaggagca agcagcctgg 840
 tccacacccc tccagatcct cttcttctg gtttagcgtt atttcttctc ctgccccgtg 900
 cctgagaagt acttcccggg ttctgtgac atcgtggggc atgggcatca gatcttccat 960
 gcatttctgt ccatctgtac actctcccag ctggaggcca tcctcctgga ctaccagggg 1020
 cggcaggaga tcttctctga gcgccatgga cccctatctg tccacatggc ctgcctctcc 1080
 ttcttcttcc tggctgcctg cagtgtgccc accgcagccc ttctgaggca caaagtcaag 1140
 gccagactga ccaagaaaga ttctgagggc tggcaagtgg ggcaacgtgt ggaggaagcc 1200
 cctcataatt tggagaaaac ttgatacaat agaagctgac ttttaaggca ttggctttta 1260
 aattaatata tatatccaag gatattgtat agctgcagtg tttgaaagcc aaaggattta 1320
 agagttttgt tgttggttaat aaaaggaata ctcttttcc ttttgatca tagcttaaca 1380
 aggcacagga aggggaaggga tcttgactaa gattcatgag acattgaatt aaggagaatc 1440
 atcttcatgc ctgaaaattt agcaaaattc cgactatggc ctccaggggc aattcctaaa 1500
 agctgaatgg ataataaaat tggactggaa agtaagtagg tggctggctc tcaccctgtt 1560
 ggaatggcta tctactatg ctgttctttg gtaatggaat aaattgaccc aaggaccgaa 1620
 tttcatttgg atttcaaatt gtccagagtg gaaaagcctt caagatgaca tgatgaatta 1680
 ctcagttcat ctgattctg gtccctcctt tctcgacaac tataatacta acccttttct 1740
 caggataact gtctacacct ggcagtttct tctgacgtgc tgttactca catccctacc 1800
 ttgcatggta atataaagga ctaggaagca gtcatacttc caggaaatgc ttggattcat 1860
 gtggacattc aggaagctta ttctcatata atactaatct aaacagtact agaaattaca 1920
 gtgccaagag ccaccaggag gccagcccaa taagcataga tactatatgg tatcatggga 1980
 cccatctatt ttttaccagt ggactacagg attacttgag agttatcagg gctgcctaac 2040
 agaccaggag atctgggggt tgcaccaggg aatcgccata tttgaccagc atgttttaaa 2100
 agctcttggg aggtattagt ggttctaagg atccctctag ggacctcatt atttcaagag 2160
 gaacccaaag tccagcctcc tacatagatg ctgccccacg aaggacccac aaaactaacc 2220
 tagttcaggg ttctcaggca ggcagttctg cttcagctta gagcagaacc cataaaatac 2280
 tcaagtactg ggataggcaa gcatgtgtgt ttactgtgga ttggtccctg aaggctcctt 2340
 tgggtgagaa catgtgaacc aggcaccctg gtttgtttgg agcattgtcg cccagaagct 2400
 tctatgggat agtggtgtct tgggattgat gtgttgtggc catgcagccc tccctgagga 2460
 ttgacttctg cactaatcca gtgaaggagg ctgtgtcaaa agaagggtc agaagccctc 2520
 ttttcagagg caatgattcc tgtcagtatg aggtccctta gttactaaa agggacatga 2580
 ttttaactcca gtttgatgaa cctctccga gtttacttta ttgtctcaa atcttttgtt 2640
 ttcttctctt ttgtgagatt tgtgggtttt gtgccttata aatggaaatg tatgaacaca 2700
 atatattgct atgtagaatt ttctgttctg ggttattggg ataagaaaa atatatattg 2760
 ctcttcaact agtgaatgaa agaaacttca gaaagctaga attgcttatt aatcaaaaga 2820
 ctttctcaat ctattttggc cacaacaaaa catattcaac tgaagcttct caataatctt 2880
 tataatcaaga aagcatgcgt cttgtcagct acattgtttt cttagatgga tttctcctgt 2940

taatcctcaa atatctgaac ttctgtgtta cccaagtgtc ttatacaagc ttctgggtgc 3000
taggacaaat ttatggcaaa taaaattagc aaaactgaaa aaaaaaaaaa aaaaaaaaaa 3060
aaaaattctc ggcgcaagaa ttagctggcc 3090

<210> 99
<211> 1274
<212> DNA
<213> Homo sapiens

<220>
<221> misc_feature
<223> Incyte ID No: 4992201CB1

<220>
<221> unsure
<222> 1191, 1250
<223> a, t, c, g, or other

<400> 99
cgccacgagc tggctcttgaa ctcccgacct caggatgatcc gcccgcttcc gccccccaaa 60
ggactgcgat tacatgcgtg agccaccgag cccagcttcc aaaagtttta agcagagctc 120
agaggtctta agcacaggca catcagagga gcatttttga aatgctttcc agcttcctca 180
ataggaatgg aagccaaatt ccgaatttat gactcctttg aggaagtcga gagctgtaag 240
gaaagccagg aacaggggca agggagagat gcatcccgaa tgatcctgtg ccaattcttt 300
ctggaatctt tgatgtgatc tcagctgccc ttctataact tgacacagtg attgtggcac 360
ccactggtct agctgtggtc tacaaggaac ccccaaaagg aagggcacag tgagcagggg 420
catcggcctg agtgacaagg atttgagagg gcaggttgga tgcagggaga ggactggcca 480
aatgccatgt gtctggactt aggcctgcctg gttcaaaactg gacttcaccc tttttgactt 540
catgatctgg tacaagttat atgaaaatgc attgctcctt ttctagtctg taaaataata 600
atgaaatgtg cactaataac tgggagacta cgcagaggaa atgaaacaag ctgcatagac 660
cacagagctc agagcctggc ctttaggaag ccctcagtaa gggttcatga tgccatgggtg 720
tctgtcatca tctcttttat cctcatcatc accttcataa tctttttatt gttcttagag 780
aatagtttag agggactcat tccctgctat catgggtgag atgtctatga aaaggacaac 840
cagtggggga ggaagcaaa attttgaata agatttctga gaccgcccc ccccgaccat 900
aaccaagaac agaaactcca cagtctgctg agccgacagt ttgcacgttg gtctcctccc 960
atctgcccac cgcactctcc tgtttttttc ctgaggatga ggatacaaaa caaggctccc 1020
aaccgtccct cagcactcac tgaactgccc ttccctctg ctgggccatg accacggaga 1080
acaagtccac tgtcctccct gcgtgggtgca cgatggaggc tcagactcca tcctcaaggc 1140
tggcaagaag acaggtgaga catgagcctc ctgatacagt gatggttctg ncacagatgg 1200
actcacatga ggtgagccaa tattacatca tgctggagtc agccaagctn tagcaatgca 1260
gtccacttag atca 1274

<210> 100
<211> 1514
<212> DNA
<213> Homo sapiens

<220>
<221> misc_feature
<223> Incyte ID No: 5441583CB1

<400> 100
aatctgcaaa gttcctgtga gcgctgtcat tttgtcactc tgggtttttca gattcttccc 60
ctggaggctg gagtttccag gatgtcaaaa ttacctctgc ttgggtgagc tatttcaagc 120
agctgggata cctgtgtcac tcctgtgtgc tgccagtgac tgcccagggtg tctgtctggtt 180
cctccccagg agtagggagg aaccagggtgg gctggctggg atgggtggat atttaaagac 240
caggccttgg acgctgcagc acttctatct ctgcttgatg cctgctgcca cgtggctgggt 300
cctcctcctc tctctgtggc tgagccttgg ggtgaagaca ggcagctgct cccaacccca 360
gaacctttgc tgtcttggga cggatcacca ctgcaagagg ggaagttgct actgtgatga 420
attctgccat gtggcaccag actgccaccc agaccacagt gtcctctgca accctggtaa 480
ctcacatata ggcccattc cacctacagc aaagctggat gcgatggctg gcagaggcaa 540
accctttgct tgcacttcag gccaaagccg ggatgtggcc tagatgggtc ctaagggtccc 600
tgacaatcct gagatcttgc atcttgtcta tttcagggtc aagggtgccta catgctcctt 660

```

ctagctttgt ttccttgatg ttccttgcca cctgctactc ctctctgagc tacttttcca 720
gggtccacag ggagagggtc agctgtccgt ggtagacatg agggtagaga gtgagggtgt 780
tgggttccac ttacctttct atcatcttgc agtatggatg tctcttgact ggtaccgtgt 840
agacttttga gggcacacag gagtctgcag agagatactg gggacattga gcagcagcag 900
tggggttggg aggcagaaat gaggacagga acatttacct tgtgtttctc tcagcttctc 960
agatgaccaaa gatgggtgctg cagatgggtgc tgaggatgga gaaccaccca agccccgcta 1020
ggagccacct agactggatg cagagcatgg ttcagcgagt tctccaaacc agcctcccag 1080
gaatgccatt ctccatggct gtgaggagaa taaagaagag agcctgactc ctctcctgag 1140
gccccctccc caccctgagc cagcaggatc cacggagcag aggtcatctg tccccagctt 1200
ggcccaactga ggccagcatg gctgggcccc ggatgcttgt ctctcagctc ccattcctgtg 1260
tacttccaca ttggtttaac cagaggaaaa ccgaaatcta caattgtcat aaacacattt 1320
aaatgtgcgt agaatcagcc atacaaattg tgaaacatac attgggctca tgggtattatt 1380
cactgtttgg tgggtggttag agtgactata gcaaacattt cttgaaagta gaaaataaga 1440
accagcacc ccttgagcta agtaatgtgc cctctgtcct caatattcct tcctgggctc 1500
cacattactg ccct 1514

```

<210> 101

<211> 1380

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 1639243CB1

<400> 101

```

tagtcatcga actgccactg ccctcctctg agcccatca tgctgggttg ttatggaatg 60
ggacagctct gtatctggga aagtcctcca gcctcacctt cttggctctt aagtgtaggc 120
tgttaccact tgcccagcct gggcctcctc tctccccacc ctttcacaag gcagctgcc 180
ttccgcactc actggcccat cccttcttct tcatcgctcc acccttccac ccccgctgcat 240
ggctgctgta ggtctggttt ttttgttttt gtttttttta agacggagtc tcaactctgcc 300
gccaggctgg agtgcagtgg cagaatcttg gctcactgca acctctgcct ccagggttca 360
agcgattctc ctgcctcagc ctcccagta gctgggacta caggcacctg ccaccatata 420
cagctaattt ttgtattttt agtagagatg gggtttcacc atgttgcca ggatctcttg 480
acctcgatgat ctgcccgcct cggcctccca aagtgcctgtg attacaggcg tgagccaccg 540
tgcctagcca ggtcttgttt tgaaaacctc actgtgggag attcaggcat cctccctaag 600
ccagctggcc gctgtgctaa agcctgttca gagttaataa taatcattag ctgaatggg 660
ctggggcctt tcagcttcag atctctaagc acttgaggc tgagtcagtc agccctcacc 720
ttccccctcc ttectgggct gcagagtgtg acagaatggg aaggcactgt ggggaaggaa 780
tcaggaatct tgctgctagc cagccttgc agtgacttct cgtctgggag tgggactga 840
gtcctctcag taaactaata agacttgac ctgacaaaagg tcaagatatg tagggaacac 900
agtgtatgct aggtgagac ctatgggtgt ggcagggtg gctgttgagc ctgaacttcc 960
agtactcctg cccttcttct tgtttacctg gcttggccta cagggggcac ccctggtctt 1020
gatgcctcaa gccagcatt tctgggtccc ctctgcaagc tcagagagca gggaggcttc 1080
tggtagtgtc cttgatgtct ctgtgtctgg ttggcacaaa gatcctgtgt aacatgaaat 1140
gaaaggtgca tcagcttggg ggctgggaaa cctgcagtat gggtttactc cgtccctatc 1200
actggtgtgg ctgtgggcaa accacttatt cctgcaccta cctcacaggg atgttgtag 1260
ggtttgatga gagaatgaat gttaatagga attggaaaat tcaaagcatt aaacacatgt 1320
aaacaggtgg taaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaagatcttt 1380

```

<210> 102

<211> 942

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 1335166CB1

<400> 102

```

gccttgccct tccaaagtgc cgggattaca ggcttgagcc actgtgcctg gcctattatt 60
gttatttaag atggagctt gctctgtcac cctggctgga gtgcagtggt gcaatcttgg 120
ctcactgcaa cctccacctc ctggattcaa gagattctcc tgcctcaacc tcttgagtag 180

```

```

ctgggattac aggcattgcac aaccacactg gctatttttt gtatctttaa cagagacagg 240
gtttccaccat gttggccagg ctggtctcga acttctgagc tcaagcgatc tgcctgcctt 300
ggcctcccaa agtgctggga ttacaggcgt gactcactgt gcccgaccag ggagactatt 360
atgatggcta tgagtggaca atgccaagaa actgaagtac actccttatac aggagtttca 420
acaaacaact agtcaaaatc aaataactca aagtccatgc aataaagata gttcaatagc 480
atttgagtcc aattgggtcag tctttgttca gggatgatgg tgattaaagg ccaaaccctc 540
cccattaaat gagagatcta cattagagag gtttcagaga cctccaccca gcaacagtgg 600
gcaagcagcg tttataaaca gaaaaaggaa gtggcattgg atataggctg attgggttaca 660
gctcaacctt tgtctaattt ggacacatct ggtcagtcgt cagtctgccc ctggctgaag 720
gctggctgct atgattggag aagactcctt tcttcttaca agaataact gcctgggtcaa 780
gttgtagttg gtttacataa taaataaggt tacatcccag tcaaatcctc cctacaaagt 840
atccctatt ctgagctcta ttacataga ctagtttttc tgctttaaataaatgggagtc 900
atgcaattaa cgaaacaga gagaacaaat gggcgccac tc 942

```

<210> 103

<211> 1815

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 166894CB1

<400> 103

```

gctcgagggt gaggaggaaa gccattactt caccaatgtg aatcaaggtc attcgtcacc 60
tcgggaagca cacagagctg cgacgttcag gaggagggca cgctgctccc aagccaaacc 120
acgatgcagc ctggcaccag tcccatgcag caggggagaag gggagtttct ggaagcgagg 180
atgagctggg cagccaggcc cagggccctg gccgtgcatg catcagcccc tctgagaaa 240
aagagctcaa ggactacatg gggcttacc cagcagtggtg gcaacgggtg agtttccagg 300
aggcgttggc aggagctgac cggccactga cgctggccgg ggtccctagg ggacagggac 360
tgagggttca gcggcctctg accgtccagc gtcatctgag gaaatcaggc caatcgtgag 420
ccaccagagt gtccctaccg agacggcctc tctccctagt ataacaagcc acatttgcgt 480
cctttctggg agagttagaa gcaagccacg tgggtgactt cttttcaaag tctctgcct 540
cagaggagga agccccatgg gtggaggcca ggcccgaggc catcctgctc tgggccagag 600
cgtgcctgtg tctgatgagc gaggacatgc ctctgtggc tgagccttcc atttcacacc 660
ggcagctcga accccccatc acggcttctt tttgggtccc cctcctcagg ggaccgagct 720
tgaaatgtgc ttgcccttca tccctaaggc ctgcgtccct tctcagaga ggactaagcc 780
cctctcccag tcttgatgtt ctccgaggcc cttcttattc acagaacctt tctggcttat 840
ttattcgctt gcttgcttct gatgtcttct cttactgaaa gcttgctgca aaggaccaca 900
cctgcttctc gaccaggaa cgtgggaaag ggaaaggctt ggcttgtttt ggtggagatg 960
gagatgctgg taacagtgga ggaatgccct ccttctgatt cacagtgggg aggtgctctg 1020
ggccccctgc actgcccagc gacttcagct tttggttgtc ctgctgagag gatgcggcat 1080
ttgagcagta gcttctggag cccagagtga agccccacc aggcacctg acaggaccac 1140
aggaatgctc ctggcacagc aagcaggact cttgagaagc tcagcctcca cgctccttgt 1200
agatgttcag ttcaaactcc acagtttgtg tgactcactg aagggttgg tttggctctc 1260
gctgacgagt ctgtcatcgg tgccaggagc tgagaacagc ccccttccct tttgaggctg 1320
gctgtccat gcgcaccctt ggccctcacat caatggagga tcacccatcc ctctcctgtg 1380
ctagaaaccc aatggctgta ttccataagc ctgcaggact cttgctcttc tcttatttta 1440
actacaccag cttaggagta gcctacatgc tccatcttca tttccttacc ccatctactc 1500
cacagtctac cattctcttg ctgaggttac ttacctggcc tctttcctct acactctttt 1560
ccacactcac ttgtccaggt gcacattaat acctacacct gactatccta cagataactc 1620
gaattcaaga tgggtggggga ataaactcac catcttactc tatatgtctg cctgtccttc 1680
agtatttctc atgtgggata aggggtctat gattcaccca cttgtccaag acagaacacc 1740
tcgatgtcat tcttttttta aatttttaat ttttttttga gacagagttt cgctgttata 1800
accaggctg gagtg 1815

```

<210> 104

<211> 1120

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 217969CB1

<400> 104

```
ctccgctgag ggaactggca ccttctccac acacctcctt cgccgagtca cagacaagct 60
atgcagagaa atctaccgga gccacggcaa ggctctggaa gaggacagag aagggagcag 120
cttcagagg ccacatcggt ttgtcgccat gacaacagag gggcggggccc gtgaccacct 180
gcccctggca caaggccccc aagtgcgaa gggctactga gtcttaagca aagagcaaga 240
ggcatcagcc tcatgggaca atccaccag agtcgcctaa ccacagacc caaaagtccc 300
ctagtggaca ccaccagtc caggaacaga tcaaattcaa gaatatgtaa acttgggtgg 360
gaaaaaaaaa cttttcctcc actagccagg aactaatatt tagcattttc attatgaaga 420
taggcagcat accacatgag cagcctctgt gtgtcggtca ccagtaaaaa tcataatatg 480
ttcatggcac atgatgggta ttgcagtttt gttttttgtt ttttttttga gacagagtct 540
gcctctgtca cccaacctgg agtgcagtgg taccatcaca gctcactgca acctcgacca 600
cccggtctcg agggatcctc ccacctcagc ctccaagttg ctccgactat aggcgtgtgc 660
caccacaccc agctaattct ttttagatgg ggtctcacta tgttgccctg gctagtctca 720
aacttcgggg cttaagcaat tctctacat cagcctccca aagtgtctgg gttacaggca 780
tgagccacca caccagcct gttgcagatg tcttgaaaga gtatttacc ttcattctca 840
ctgtgaaatt acggttggtt tcagacttgc cactattgaa tgcattatta aagtgttaca 900
tatatttcta catcgtaaat gtttataatt ggcattttga tatcagtatt tcaagagttc 960
tctccttctg aatccaaagt gtgttatatt ttaaatgtaa acattattct aagaaggggc 1020
tcaggatcaa ccctggagga ccagacgggc tacagcacac acaaaaaagc ttcagtacac 1080
tgtattagtg tgctgaggat gtcgcaacaa agtacttcag 1120
```

<210> 105

<211> 535

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 335237CB1

<400> 105

```
aaatgaatag ggagctttctc tagactgggt agccccaaaa ggtctgccgg aagagatgct 60
atttagtctg aagcctggat cgtgatgaca gttccccctg tgaagggtgca ggtagagtgt 120
tttgtaaaca tgtgaccatc acctgtgatc atatctttaa ccctcacact ccccatcct 180
aaccttcatg aaatgctgct catgctctga gctagtatct ctcatctaac ccctaccoca 240
cctccaggag acctgcctg ttacagtcta tgccgtattt agctgttgtc cttttctgca 300
gtagagtccc caggagtctt tcagggacag gcagtcagg acagctggtg ccagagcct 360
ccctggcttg cccactgggc agctccaggg acaacttgac ctgcccatac aaagccaaag 420
gtcagaacag gaggcagaac ctgccccggc cctcttcaaa ctccaaggga aagcctgttc 480
cttgatcct ttcagaaata aaaacaaaat gacatcaaaa aaaaaaaaaa aaaaa 535
```

<210> 106

<211> 1188

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 938306CB1

<220>

<221> unsure

<222> 1052

<223> a, t, c, g, or other

<400> 106

```
agtaactgtt ccttgccctc ccggttcaaa tgataccctg cctcagcccc ccagatagct 60
gggattacgg gcacatgcca ccacacctgg ctaacttttg aatttttagt agagacagag 120
ttcgccatgt tagccaggct ggtctcgaa tcttgacctc agatgataca cccacctcag 180
cctcccaacg tgctgggatt acaggtgtga gccactgcgc ccagctggct tatgctcttt 240
ttctaaaaat ttgtccactt aggaaagtgt cttgatttca ataaaattag catggattaa 300
```

```

attacttttt ttcttttttt tttttctttt ggtatttgat cttgctcgat gaatgtggac 360
atcagctatg ggtaggaat gctattcagg aacatctaata gtgtttgtga taatgtattt 420
gacttggtgc caaatggtac tcttgagaga tgtggttggt actttgacag attaaagcaa 480
agttaaattc tgatgaagtg caaaggaatt ttatcggtcc caggatggtt gccactgtg 540
ttgggcaaaa gagtgttttt ccaaaaaggt ccagagcaat ccgcatgtat tctgtcacct 600
ctacttccag tttcttcaaa ggccagccaa aagcttcatt ttcctacttc ctgtcatttc 660
caaaaccaca gcttaaacct gaaaaacaaa tgggaagctg tgttcctgcc tcttatgatt 720
gctgccactt acaaaccagc aagaacagag cacagcaagc agaggagagt tcagtcttgc 780
tgatgtcaca tgaagcagaa tgaggactct gcttggttgg ctcctgagtt taaaagataa 840
gtagacaatt tattgttggg aggttgagaa aattcaagaa cagagagaaa tctgacagtc 900
tggccataag aagaaatttt tgcatttatg ttcttatagc catacctgaa gtcaactaat 960
tacgtttgga gagaagtaga atgcttgctt taagaagaaa aaacaggcta gggcgcggtg 1020
gcacacacct gtaatccag cactttggga gncggaggcg ggcagatcat gaggtcagga 1080
gatcaagacc atcctgggcta gcatgtgaa accctgtctc tactaaaaat acaaaaaatt 1140
agccgcgcgt ggtggtgggc acctgtggtc ccagctgctc ggtggtgct 1188

```

<210> 107

<211> 638

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 1448129CB1

<400> 107

```

ctaagtgtga gaggaaggat tctgtggttc ccttaactgc aaaggattgt tcactatcgt 60
acatttagag ctgaagagaa ggcttgcgaa ggccatttga cttttttttc ctcccttctc 120
gcagcaggtc catcagcaat gtcacgaacc tctaaccaag acccctcaag caatccaatt 180
tctggcagca cttaagtgtc ttccatactc ctgatgagac atgatgctca tttctcaaca 240
aagatttatt gaatgtctgc tatgtttaat gcaccatggt ggagcctggg gaagatgcct 300
acgccatacc tgctgtcctt aatgaactca caggccagct tcgggcagac attccagcaa 360
gcattggagt ccaggctaata agtaaccaga gagaggtaca aactgggtga aagaaaggaa 420
cccttcctgg aggagtcagc atttgagcag ttccctgaagg tgctggtggg tagagggcat 480
tccaggcagg tgggtctctt cacagagtgg actgcagttt ggggtgcctg acaaagatga 540
gaagggttag gtttttcctt cctccctcct gcatgaaata atgggaaaga agccgtgata 600
gaaagattta gtttgagata ggataaaggt gctaagaa 638

```

<210> 108

<211> 648

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 1761049CB1

<400> 108

```

actgtaatgg gatactgata ctgcctcata gataatgaga ttaagtgtct cctaccttga 60
atgtctgtct cgggtgttggc ctggagatat ctttcccaag tgtagaggaa tccatctctg 120
tgcccgtggt aatggcttga caccaggaga gtttgagatc agcctgggca acatggcaag 180
aacctgtccc aaatttaata taataaattt taaataaaat tttaaaaata aaattaaata 240
aaaaattata caggagggcc atgtcagggt aattttaact gaagcaaac ttatttatc 300
ataacaatga agtttatatt actaggaaaa aatttcattc catgataaat gtgtggtatc 360
atgtttttct tcaaaatatt gaatttaaag aatgttcatt gcaatactgg caactaagtc 420
cagatttgct ttttaaccat ggagtaattt ctgagaaata tttgttttat tttattttat 480
tttattttat ttttttatg ttatttatgt tatttatgtt atgttatgtt atgttatgtt 540
atgttatgtt atgttatgtt atgttatgtt tttgaaacag agtccttgctc ttgttgccca 600
ggctggagtg caatggtgtg atctctgcct cctgggttca agcgattc 648

```

<210> 109

<211> 1181

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 1959587CB1

<400> 109

```
atttaaaatg tcttgttggc aaaatcatgt aacacctaaa gtcattcttt gtgcaaactct 60
catacttttg aaattactgt tccaggaggat attcatgtga aaggaaagct gtttgaacta 120
tatatatacc caagcctgta agcagaaatg actggacaga aatacaagcc tgtgagacag 180
agagatctag gagatcccta gagagggtcca tctagacctc caggatgaaa ttacctcatt 240
tagctcagtt tctcacctct ccgttagtgc tttgggtccac ggggggtcagt ggttctgcag 300
gatttcacca actggttcca cagtgggaat gtgaagagg acctggctgt gggaaaagct 360
gcttatcaaa gagaggcctg atagaaatgc tagggaaagt ggctgtgagc ctgcattatg 420
gacggggagca gagcggcagg gcatgttgct aggactaagc agtcgattca cgtgagtcctg 480
gacaccatgt ataaatatct tgaaaacttc ctgttacaat ttaatttact ccattgtgtt 540
agagggagtg cttccctgca catggtgttc tgtatatcta ttggttgatg tgcttttaggg 600
tatagtaatt atttatagca atagattgat acccagggtg cagagttgct ttagttatat 660
ctttatttcc ttttcttaga gattatagac ataatttgct tctgtcatct tttgaccaa 720
aatctaattg gaatgaagaa ttacacttga atgtttatct ttaagtaata agttctgtga 780
tagtgattat aggtgtctgc atttaaaaaa atgttttaaa aagtacagag gtcaaacaag 840
catctgtatg attgaacaaa tgttatttct tatatttctg tcaaatttta aaaaatggat 900
aaagaggccg ggcactgtgg ctcacgcctg taatcccagc actttgggag gccaaaggtg 960
gtggatcacc tgaggtcagg agttcaagac cagcctggcc aacatggtga aacctgtctc 1020
tactaaaaat ataaaaattg gccagatgtg gtgggtgggt cctgtaatcc cagccacccg 1080
ggaggctgag gttggagaat tgtttggacc caggagacgg aggttgctgt gagcccacat 1140
ggtgccaatg cactccagca atgggcgaca gagtggagaca c 1181
```

<210> 110

<211> 1291

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 2303463CB1

<400> 110

```
ggcctctcac ggagactttc ccaccagtgt aataaagagg agaaaacgtc acagcgggaag 60
ggcctgaccc tgtcgcaccc actaaggaaa cagctacggg gatgggaccc tggaaagctgc 120
tgtgggagcc tcatccaccg cttctctgac cccacccagg ctgcttccca ggcctcaggg 180
tcttagtggt gacctccggg ccgtgattaa tgcagttttt cagggttaagt actgcacgac 240
tactcgcttc tgaaactgat agacactgcc tcagctccgt gcagggcaga cgcacaagag 300
cagaatctcc gtgggacatc tctctggagc atcaatatta ctgcagtatt tggagaagaa 360
aaatttaaat aagtctctag gtaaagaatg gaacatttaa gacaagtctg gaaagtcatc 420
tgcctttaat aactgtcgtt tgtccttaac gtcagacttt ctccaagaca aaaactctaa 480
gaacttattt ccattcttac aaatagtaaa aatgataaat catatcaagt caattgaaag 540
tctgcctgct tgccttctca aatcacaata tggccttggt atgggtttat ttgtactttt 600
gtgggggatt cgtggatctt tagattaaaa aaaaaaatag gtttttgaac aatttttgca 660
agtttgcagc tgtagtttat tgatttattt ttgcaaccca tctaattctt ctgtctctct 720
cctctcctca ggctcaataa ttccatgggt ctctaaggct gtgtttattt tcttttaatt 780
ttgttgattt acttttatac tccatttgat tttttctatc tctcaccttt tctcagactc 840
agtcacccca caggctctgta aggctctgtt cattttcttt aaactttttt ttttctctct 900
ctcagattgg ataaattata ttgctatgtc tctgtttctg aactatagaa aagctcaaaa 960
ttactatttt tatttctcat tttttatatg gctatttttt ctctgctgat gtttcacatc 1020
tattcattta tgagaatatt ttcctttgac ctcagagcgg tgtttataat agctgccttc 1080
aaattcttgt ctgccgttta catcttgagc atcttgagga tggctactgc ctgcttttta 1140
tcttggtgat ttattacatt ttcacgtgtc ttcacgcacg tcttgaattg gaaattgtgc 1200
cctggagact gtatacaaga ctggattaaa aagactggat tctgaaaaaa aaaaaaaaaa 1260
aaaaaaaaaa aaaaaaaaaa aaaaaaaatt g 1291
```

<210> 111

<211> 594

<212> DNA
<213> Homo sapiens

<220>
<221> misc_feature
<223> Incyte ID No: 2512281CB1

<400> 111
tgatctccca acttgagtt tcaaggctgc ccttcccttc cctgtgctgt gggtttctcc 60
ccccgggct ccccgctcc tccctcagtg ctctgagagc acatggcagc tgccccagcc 120
ccgaagccaa gcctggcgcc tgtcctcggg cctctggaag tcctcccgcc ccctctacaa 180
gcccctacca gacgctctcc agggacggag tgtgcccctc cagccacagg aaaggggaga 240
ctgatccgcg tcaggtcccg ggatggaata gtaaccatga aatccagccg cagagcaatg 300
tgtctcaagc catcagtcac ccttcctaatt tctcaggaag ccaggcatgc ccttcaccca 360
gctgaacctt gaactccgcc ttggcctttg caaggaaatc agccagtaaa taagacccca 420
aaatcggagc aacctgagaa gacatgacca ggttacaagg agggagaact gtggccaagg 480
atggccacag gcccagatg gctcaccacg ctacacaaag aactggcctc ttggaagagc 540
ttcaacattt tcagagatac cagcgagggg gcggtcattg actgacttct ttcc 594

<210> 112
<211> 852
<212> DNA
<213> Homo sapiens

<220>
<221> misc_feature
<223> Incyte ID No: 2755924CB1

<400> 112
tgagtttatt tcaccacagt cgtaagcag gatgtgatgt ttattaaaat tcataattta 60
tttttctgta tctgtgtgct gccacacctg gccattagcg gttggagctg cccctctttg 120
ttgtcactgt catTTTTTaa acatagtata tgtattcttt ttcttttttt agtgacaggg 180
tttactatg ttgccacac tggccatgaa ctcttgagct caggatgatc tctaccta 240
gcctcccaag tagctgggac tacaggcaca tgccactgtg ccagcttgt cactgccaat 300
tttaactctg gcatgtttgt cccattgctg tattgccatg tgaaaaattt tgccaacagc 360
caggagacct ctgtttcctc tgtaaaagttg aatctttctt ccttgtaatg acttaatatg 420
gatttctatc aagtggagaa gaaatcttaa aaccgtacat actgttttgg gctctctggt 480
aggttttagg aggggagctc cccaggaggt aaatgggaat gccatccagc tttggcaaat 540
tcttattgaa actaatgagt ttaatagact gaccttatta ctctattta caacctcatt 600
acaaaattat ctgctcaatg tttgctaaat tctttatcgg tcttgagaag ctaagcccc 660
tgtttaattc ctgctaaaac cgtgaaaatt ttccccctt catatgaaaa gagtagcccc 720
aaaggccctt gagtaaaaat aaatagtggc tttggcaatt ttgacatcag gaataacct 780
gggaaatttg gcttaaagaa tttaaagcta aaaagtggtc tcctatacca cgtgaataaa 840
aaaaaaaaaa aa 852

<210> 113
<211> 1361
<212> DNA
<213> Homo sapiens

<220>
<221> misc_feature
<223> Incyte ID No: 2796369CB1

<400> 113
aactgtgtgt acacagtgag cgtaagtgc tcagaagact caaaagatat ctaacccatg 60
ctaagtggaa atttgggccc gcgtaaaaat ttcaccacgg gcaacagcta tgaccagga 120
ttcccccaag gtcataatac ggccccctat tgggcaaaact ggtaccccaa aaagtcccg 180
tacgggaata attcccgggg agacgagtc actagtcagg ccgctgccct ttttttttt 240
ttttcaatgt ttaagaaaat ttatagattt gtgttgggct gcattcaaaag ctgttccatg 300
ggccaacagt tggacaagtt tgccctagat catggggtct ttgtgctctg agaatggg 360
agtttgggag ggactcagct ttctgttggg tggccctgga tctggatctg gtgctgcccc 420
ctttctctgg agcacacaaa gggaacaaga aggccttgac ttgggcaagg aagccatcca 480

```

cagggctccc cagaagccag ggccaccagg agctcactgc tgtgcagagg caactcggct 540
tggttatttc ttgcctgagg caggggaacag ggaatgcagg gaagccaggc agcagcagga 600
ggctccaaac gctggtgtca gcaagccaga acccccacct gacttcactc ctgtctgccc 660
agctcacagt cgctgtcac tgggaggtcc ttggggcctg gacctgcctg acctgtggcc 720
ccagaagggg ctaagcccag agtcacatgg gatggagcct ggaatgcata ggccaagtgg 780
gttgtgcctg ggcagcaggc ctgggatatg aggggcctac tcctccatcc acagccccga 840
cacacattct ggaggggatg tcgtacctct gtgaccagcc ctccgcaccg tcacataggc 900
cagcaggggg caatgccagg tgccatccag agcccacccc catgtcctcc cggagggtcat 960
gagctctgcc ctacgtaccc cagccccaca atgggaagga ggcattggagc tggcagagca 1020
tgccgccagc cgtccctgtg tgtgtttctg aggaactcct gttcactctg agggcactgc 1080
ctaggtgccc gtccctggtc tgctgtctgt ggctctgatg aagctaacca gccttcttcc 1140
ccaccaagag ggtgtcccag cagcagctat ggctatgttc cggcgggggg aaggcgggga 1200
gccttctctc tctactggctg ttctattttt ttattttttt gaggcagagt ctctcgtcac 1260
ccaggctgga gtgcagcggc gtaattctct ctcgctgcaa cctccacctc ccaggttcaa 1320
gcgattctct tgcctcagcc ttccaagtag ctgggatcac a 1361

```

<210> 114

<211> 1650

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 3010920CB1

<400> 114

```

cttttttttt ttggtgaaac agccaagtga ataaaactat ctaccctagg tgagggtggc 60
tgtaatccca gcactttggg agggccaggc ggggggatca cctgaggtca ggagttcgag 120
accagccagg caaacatggt gaaacctcat ctctactaaa aatacaaaaa ttagccaggc 180
atggtaatcc caggtacgtg ggaggctgag gcaggagaat tgcttgaacc tgggaggcag 240
agggtgcagt gagccaaaat tatgccattg cactccagcc tgggcaacaa agcatgactc 300
tgtctccaga gggggaaaaa aaaacctgtc tacaaaactc tagtaagtta ttatttcat 360
catatggcac taagaaaatc atcatgtttg ccattaaaac taggtacact aataacttat 420
agcttaatat ttttggcatg gttcttactg aaatctgcta ctttcaacca agtcattatg 480
cctcgagagc tctgtcaaga cttaatatat gttcatctct atgataaata tttactgata 540
tttcagataa atagtttgtg ttgctgtaac acatatatc actacagaaa actgtagatt 600
tctatataag caagaaccaa ttattgtatg atcataagca ataccataag ggcaatatga 660
atatttacta gtattataca aaacagctag caccataat tgggatttca gtacagttct 720
tcctccagta cccacatcct tgagacgtat taggaggtaa aattcaccta agttttagaa 780
ctcaacaaac catatctcaa gtatttactt ctgcctttac catacgtgtg atcctgggca 840
agttagttaa cctccctcgg ccttagtttc ttcatattgt aaataataat acaaaccttg 900
ttaagttttt aacagaacaa tatattgtat gtaaagtacc taggaaaggg tctggcccat 960
agtaaatgct tgataatgac agtgattact gtaatcattg ttgaaattgc agagatcaaa 1020
caatttcctaa taatatagcc aaaaaagtga agtgctcagg accttcaaaa atatttcaaa 1080
gaatttgtaa tcaagatatt tttagacttg tatttctggg ttctacaaag tccagatggg 1140
atcttctgtc ctacacaagt taggctgtgt cgtaagactc ttctaccagg tactttcttg 1200
acatctgtga gaggcaagca ggtatttccc cagtacttaa acactaggtc tgctaattac 1260
tgagtcagat ttaacgaata cgtaaactac tctggctatg ctaaacactt agggaggctg 1320
catcagattt ttcaacaact ggtcagtatg tactgagtc actgagtacg cagctatgtt 1380
ctaaaaagtt gcattcacat caataatcaa acataatccc tgcttgagta gagtttacca 1440
tctagaggaa acaaaactaaa agaagtttaa aagaaaatgt agtatgtttt agtataatgg 1500
ctagctaaag ttgattttgc tgtgttttaa aaacgagaaa gaataataatt agatgagagg 1560
cagttataaa tataaattac aatttggatg ttgaatgtag caacagcgag aaagcacacg 1620
aacacaggat gaaacataag taatgctgac 1650

```

<210> 115

<211> 1845

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 3360955CB1

<400> 115

```

gaccgcagga attttttttt tttttttttg cattgtattt tcaactttat taaataagca 60
aaaaaaaaa actggataaa aagtaatttt gcaaagtcta ttaagctccc atttgtttgt 120
ggagggtgtg gctagtgtgc tagatccttg agtaactgac ctttttagtt tgtttttatg 180
ttgtttgttt tcagcttttg tccacaacag gcagtcactt cagaccagga agtctctaaa 240
agcaccgaga ctcttcggag gttaatgett tcggccaaga tcatggatgg ggaggacaca 300
gggctttacc atcagcactt ctcttggtac ctactatca acaggatgat ggcccatagg 360
agcaaaggca ccagttttca tgcactgccc agtttaccce ttttgcaaa tccttcctcc 420
tggcctcctg actatgacac caccagatg tccatathtt ctgccaggaa aagcctgctg 480
gggacaaaagc ttcttacttc atgtttatct tctctccatt tcagaaaatg tcccgtctta 540
cattgcaact tgttgaaagc aggaagtga ttttatagta agactaattt acattaaagt 600
gaaaatgaca ttattttgag ggggtaccct tcaagtaatt tgtttatatt ttaataatat 660
attgagagtc ttgggtgctc aggcctgttt tcctggcaca taaggaggca aagctgggtg 720
tttgagacca gcctggacaa tatagggaagg cctcatctat ataacaacaa caacaacaaa 780
aaagaatatg atagagacat tcattgagca ttacattttt tgagcattat atattctgtg 840
tcctgtgtct ccatccaaat tttagattca aaatatatta ttactgtaat gttaacccca 900
aaagaaaaat atttattggt aataatctaa gtactatgat atgtaaagaa gacctctcca 960
aatgctcaaa tcataccata ttgggaattg attactcaaa tagctctctc tatataccta 1020
tatctaaacc aggaccaaag actgaacagg ctcggaagat atctttgttt ttatttttgt 1080
tttagaaaac gggctctctc ttgttgccca gactagtctc gaactcctgg gctcaaaaaga 1140
tcctccccac tcaggctccc aaagtctctg gattacaggt gtgagccact gtgccagac 1200
tgaagttatc tttgtagggt gcttctttca catgggagat ggaacataca gatccaatac 1260
agatgggatt gttctataga gtaaaacgaa caagtgcata attaatccc atgtaagcca 1320
ttctctggtc aaaagtgttc caaaagtttc taccatttga tcctctatct ctgaagtga 1380
tacagaagca gtacaatgag gcagctatta gcagaggaag ggacctacat gggactcttg 1440
tagctcatgt ggatatggaa ctgcagaaga gagaggctga ggttgacgta aaataagacc 1500
ctgaaatcca aaggcaatca acgagatagc attggttaga gttcaaaacc tcagagatcc 1560
atatggcttt cttggggaaa aactcattta tcttgacctt aggcagaaag tgagaaggag 1620
gcagtgaaca ccaagacaga agaaaactta agcatttgaa gagggaaatt caatgggctc 1680
caggtgcctt ccatgtaaac atttcagaaa aactcccaga agtttgatag agctcatgct 1740
tgtcaatgaa actgtaaaat ccagtcaagt cgccatcctg gttggactca tagagggaac 1800
ggaggagcca agatggaaat gcagaaatca cccgtcttct gcgtc 1845

```

<210> 116

<211> 1061

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 3409459CB1

<400> 116

```

agctgacaga caagctcgag cacttagagc aagagaaaca tgaattgaga agacgatttg 60
agaaccgaga aggggagtg gaaggccgag tgtcagagct ggagagtgat gtgaagcagc 120
cgccctcct cctccaccca ggccccaaat caaaggctcc gcggcccggc cagccctcag 180
ctgctcaca cagattcagt ctccctccct ccctcacgtg gggaaagcac agcaggggatg 240
cgcggaaga atgtacctgt agatgtgtac ataccacagt gctgtaattt tgtatgtagc 300
aatcatgtaa atacatgtat ggattttata atatacatat ataaaaatct ataaaggcat 360
atttttagaa aaacagcaca ccactgcttc ttttgaaaat agtctgaata agaataaaat 420
gaatttctac agagcttctt gctcagctc ctgggtattt gcggggggcg ggtttggtct 480
gaatgcagct gacatgtcag actcaccact ggctgctgct ggagaggtag caatagtggg 540
tcccctgcac ccgggccacc tcagatgctg gtacttactt aaccaaggga tttggccagg 600
aagggcattc tcccctgccc ctcccgctg gcactgcccc ctgccagtgc tgcagcgtgc 660
catccgcaag gcaggcctgc caacgctgct tcccaggcca gcagggccct aaagcttggc 720
agaaacttca agccagaaag aaaccacgcc aggcaaaaggc ttccactgca tccttcagg 780
catgtggagg ataaacacgg agcttcccct ggcattctac cacatgtgct cccagttcaa 840
ctagggggag actcaaccct ggccacctg ttcatggaga aaatggacct ggcttctgga 900
agtgtcccaa gggcctcgac aggtcctgtt gcaagactta ggccaagaac aggggaagagc 960
aggacacggt ggggtgggaag aagccacggc cactcagcac atctctggag ggagagcaaa 1020
cagacctggt aagaaaattt ggcagcttga acctttattt t 1061

```

<210> 117

<211> 1085
<212> DNA
<213> Homo sapiens

<220>
<221> misc_feature
<223> Incyte ID No: 4102938CB1

<400> 117
ccttttcttga tctccgtaca cacgcatacc acccaccctc cctccctctc ttctcccaact 60
tcccctctct cctttctcac catctctcct cattatgcct tctctactcg atcaccctt 120
tgcagagaaa cctttctccc ttcttgctct tttccagttg aatttcctcg ccccggtgtc 180
ccaggtggct ggacatgcag cagaagggaa ctggggagac tctcgcaact ccaatcattt 240
ctcaaaactc aggttccagt ttgagaccg cctggccaac atgggtgaaac ctctgtctcta 300
caaaaaatac aaaaatttagc cgggcaaggt agcgcacctg tgggtcccagg tactcaggag 360
actgagggcat gagaatccct tgaaactggg aggcggaagt tgcagtgagc tgtgatcgtg 420
ccactgcact ccagcctggt tgacagagca acatgctatc tctaaaacaa acaaacaac 480
aaaaactcag gttccacac cctctaaacc ctgcctctc tcaggctaca gagacctctc 540
caggaggctg aagtgcctt acccgacca tctgaccagc caccgcccc tgcccgtgcc 600
ccaccgaggg cggaggtgct tctctgctct gttttatctc tggcctctga tcttgcatte 660
ttgtgccagg gcttagaccc aggcaactg ctccggtag tcaataaacc ggtaaaccag 720
caatagacat aagcggctat ttaacgaccc tgccctgaac cgacgaccgg gtcgaatttg 780
ctttcgaatt tctgcatte atccgcttat tatcacttat tcaggcgtag caaccaggcg 840
tttaagggca ccaataactg ccttaaaaaa attacgcccc gctgcaactc atcgcatgtac 900
tggtgtattc attagcattc tggcgacatg gaagccatca caaacggcat gtgaacctga 960
tcgccagcgg atcgactgcg gtgtgggtta atttttggcc tggggaacgg gggcaaaatg 1020
tccttttggc cgcttatcca ctggactccc ccggtggtgg ggaattttct acttgaaaag 1080
cggtta 1085

<210> 118
<211> 870
<212> DNA
<213> Homo sapiens

<220>
<221> misc_feature
<223> Incyte ID No: 4124601CB1

<400> 118
cccctttttt ttttttttga gacacagtat cgctctgttg cccagactgg agtgcgggtg 60
cgtgatctat gctcactgca acctctacct cccgggttca taccattctc ctgcctcagc 120
ctcccagagta gctgggacta caggcacccg ccaccacgcc cggctaattc tgtttttgta 180
tttttagtag agacgggggtt tcaccgtgtt agccaggatg gtctcaattt cctgacctcg 240
tgatctgccc gcctcggtct cccaaagtgc tgggattaca ggctattaca cgtcctttta 300
tactgactgg acttttgatc taaaaaacag tttggctgac ttggagtgat ggagttacta 360
gtctctgtgt cagggactgc aaatgctggg cctccagcag gggcagggcg gatccagcga 420
aagacagaag tgggtgggcc caaggggatg gagagctgca gaacacaagt cccgcctcaa 480
gggggcagca actgcccaga gtccgctgac ggccgctggg tgggactgca agcccagggt 540
tgccagatct gtcagctttt ttcaagataa gcttgaaatc agattttcac atggaattgt 600
cagctaattc aacaatttaa aacacagcat gggctgaatg caacatgcct atggttcaac 660
attggctgca ggccaccagt ttgtgcgttc ggatccagga ggattccatt aaggattgtg 720
cacctcagag ccacttgcaa agctgggtgca gacccttcat ttgcttctag aaaatcgggt 780
ttactccaca tgggttccgc aaaggatgag gacaaagcca gtgcctgag gtagaccgtg 840
ggagtcaccc acccactcat cattatttaa 870

<210> 119
<211> 3394
<212> DNA
<213> Homo sapiens

<220>
<221> misc_feature
<223> Incyte ID No: 4180577CB1

<400> 119

```

acgagccgga ccactagtta cggcgccagtg tgcctggaaag attaaagagg tcgctgttaa 60
atgtaaagaa aaaatgctcg tagctatttg cttcctggta ttggagcagt tcagttggtt 120
agtttatacc attggattca attcattgca ccatgggttg caaaagtgcc tgaggtcata 180
atggattgtt aaaataacta aattccagtg gttggaaact ctaggtttgt accatttttt 240
ctgctgtggg aaaaaacaac aacaacaagc atgatcaagg taacatcaca tttgatgtat 300
aatattatac tattaatgga atatcagtag acaactgtta acccattagt agcatgagta 360
taaacagtac acctgaataa attggagaca ttagccacta ggtttaacag tggaatcttg 420
atttgctag gtgacttctg ggattactgt ttgacaaata agagtaacat tttatttcat 480
ttcagaattt acgtcacttt tagctacaag agtaggaaga aggtaatcgg caaggcagaa 540
gagtatactc tttgccttag gatagcgtaa actcaggctg agacataccc ggcttataga 600
gttcttctag atgtgtagac tgtaaatgcc caaatcctct caactaaagt tttagtgtat 660
ccacaaagcc tctcatgtaa atttccagtg attccaccat tgcacttggt aatatgtatc 720
cttggttagta ccaggggatg tcctcgagca ccagttttat tttatctgcc attgcatctg 780
gattccatta cagcctctca gctgttactg cctgtggaca gttacttctg cttactgcct 840
gtagagagtt acctaacttc tcttctcagt tcttctcag gtccctggta ttttggcctc 900
agttgaaggg agtcttgctc tcatctctga gggttttaag tttgtttgat cccattgttg 960
tcttttctag ctttgagcat gtttttcagt attcatattt taacttactg agaacattaa 1020
agggaaatga taaactcgtag gtggggatat ggcagacagg tgcttggttg tttgagagaa 1080
gtagcagaag agataaaaata caaagtgcata tatgtttcag ctggagagga aagagagaga 1140
atattataga ttatatactt gtcccattgg ataccacgta tatgtttaaa tagggacaca 1200
tctccctatg ttttaactata cttataaaca actttgatac acattgcgtc ttttattctg 1260
tcacctgata ttttagtgta tctcaagtta cagattacat gtgtccttaa actatttctg 1320
aatttggact tagttccata tacagaaaga acttttagaaa attcattaat ttggatcttc 1380
tattgatagc cataaatatt atgtttatgt attctaaaac ctctttgttt agttagtact 1440
gttcatgaat gtaacaagct tcaatttctc atttgtgagt agtacatttg ctttttgttt 1500
gtttgtttgt ttgtttttga gatggagtct cacgctgtca ccaggctgga gtgcagtggc 1560
gcgatttcag ctccactgca cctccacctc ccagggtgcaa gtgatgccc tgcctcagcc 1620
tcccgatgag ctgggactac agacacccgc caccacacct ggctaatttt tgtattttta 1680
gtagagacgg ggtttcacca tgttggctag gctggtctca aactcttgac ctctgtatgt 1740
gccgcctctc gcctcccaa agtgctggga ttacaggcgt gagccaccac gccagccgt 1800
acatttactt tttaaagcag cagactaggt acactaatte tactcaaat attttcatgg 1860
gaatgtagtt atcaccaagt cctaaagtat ttttatgccc aaaaaaaatt tcattttaag 1920
gactacaaaa atgattctaa ttaaacattt tataatcaat agtaggttgg gtcttttagc 1980
attatatgtg tatatatata gacacatag tatacactta cattttgaca ggttcttcat 2040
tgagtcttga tgcactttaa acccagctgg ctaccagaga tgcgaagggt ggctctttga 2100
agattagcaa aatggagcgt tctgtcactt gagaaaagga aagttctttg cttttaaat 2160
acacagtttt catcatgccc acaatctata ttattggctg gttaaacagc actgccctat 2220
tagcaatggt aacaaaaatg aaattattta ttggcggtta tagattatct aattcaggaa 2280
atttctgagc tcaactttta cagcaactgt tatgccttct aatttagcaa ttgagttatg 2340
agtaagtttt gtgcttaact cctagaccct attgttgata accagatcaa atatagtctg 2400
tacagaggaa aacactggga acatttagta tttctaaagc ctcttttga gttactactg 2460
attgtaattt ggaactgata ataggtagag attgctaaca ctgttttttt tctctggatct 2520
tttttatgcc agaaattaaa caggttctgc taactctttt ttttctcttg gttatcacca 2580
gaatgaaaat atttaaagtg atgactctag aaaagccatc tgtgcctggg taacattgag 2640
tttgagtctc ttcaatatat attgatcatg tattgattaa tctttatttt ttcatatttt 2700
ggctagacaa attcagatct atataatgga atacccttc ttgagtgaac tatactacta 2760
atctacatga ttatatagta aggaaaaaag aagaaataac tgtaataggc atagtgtttg 2820
ttgttggttg tctgttcatt catgtgatac tactcatttc caaaattcac acaaaactac 2880
atgaggtgga ttatttgttt tgttcattat ttagtctcta tatgtttttt ctttagaaac 2940
agagtctcat tctgtcacc aggctggagt ccaatggggc ggtcatagtt cactgcagcc 3000
ttgaactctt tggctcatgt gatcctccca tctcagcctc ccacagcagg tgagactaca 3060
ggtcatgccc actgtgcctg actttttaat tttttttag agacgaggtt tcagtttgtt 3120
gcccaagctg atcttgaact cctggtttca agcgatcctc ccacctcgcc ctcccaaagt 3180
agtgggattt caggcatgac cacctggcct agttcctata cttttcttaa ttcttcagac 3240
ttctcacatt tagtatagt cattcatttc atcttctgtt ttattagcac cctttgtggc 3300
cagggaaata aaaggtggtg aaattcagtt ttcagtttag ttcttgaaag ctctgggaaa 3360
tggaagaact tccaagcaca gtgtcgataa tcaa 3394

```

<210> 120

<211> 2343

<212> DNA

<213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 5265807CB1

<400> 120
 gatgtggagg aagtgcgttc cctactctcg cagaaggaga acatcaatgt gctgggtgagt 60
 tgggagcagg gaggatggcg agaccggaat cctggctccc gtgttctgga cccctcccca 120
 ccggtggggc ggctccgtgt ctctctgct cagatgaggc cgcagttctt tccatcctgg 180
 agtgaaagcc tactctgcat ttggggatgg agagcaggtc aggcttgaag cccagatcac 240
 ttgggtgggg aatgccagg aacgtttgttc tcttctcaac ctccggagtc tgcagcgccc 300
 ccacccctcc tggaaaaagg agccggcttc tcagccatt ggccctgact ccacccctcc 360
 taggtgcttg cctttgactc ccaataccgg gacttaggac gctgcagaaa ctatgggaag 420
 aggttggggg gaagggcctg gagctctgag acttgggctg gggtccttga tagcccccct 480
 accccgcgcc catactccct gctctttttt gccttcccag gaccaagaga ggcgaactcc 540
 attgcctgct gctgcctacg taggcgatgt ccccatctc cagttgctac tgatgtcagg 600
 tgctaagtgc aatgctaagg acacactgtg gctgaccctt ctcatcgtg ctgctgcctc 660
 ccgaaacgag gtatgcaccc ccaccttcta ttaccctttc aaaaaggata agttgcttaa 720
 aggtggaggt taaggggtta ggagcctcag tctcagccc tacaagctgt tctaccccag 780
 gtagtgagga gatcaaatct tggcccttc ctctacccc tagaactggc cagtaagggg 840
 gttgagtgtc gccagctgat gtgactcagg aagagcttaa cactgggctg gcagactgcg 900
 ggggcccctg tgccaggcct gagtgactca tccctccctg gggggccacc ttgagctcga 960
 tccccctctg ggcactgaga tttcttttcc ttgtgatgga ggatgggtat agtaatctgt 1020
 tgcctccttg gccctcaagc agcttgaaag ttccctcag ccctgaagac agggaaaggg ggacctctc 1140
 cagttgggtc ctttatcccta tttccatcag cctgaagac agggaaaggg ggacctctc 1140
 ctacgttcat cacttccctg gctccagcca ccagggggag gggccgaggc cagtgggaat 1200
 gaggtgggtg gtgggttagga ggggcatttt agctcgggtt ttgcaaacga accagtatct 1260
 gtttgcttac tgagatactg aaaataatct ttgtgcctat gagaaggcta aagaggggtg 1320
 tcagggcaag atgtagcaag agcatttagt tttcctagaa gtatagggg gttgggatgt 1380
 gtggatctcc acatgtggat ttgtttggga atggatgtat gtgtgagtga agattttcag 1440
 ggtctaagtt ggcgagtcag gtggcttgaa gaagagcaaa gggtccttgt ttaatacagc 1500
 aaggtaagga agagcttaag ttctaagaag atccagaaga tggaaagtag gtattcagag 1560
 gctaggtatg ttggaaaaat gagaagagat tctgtgctgg ggaaggggct gtagatgcta 1620
 tgctgacctc tctctccttc catctctgtg tgttttacct cctgctocca ctctctctt 1680
 tgtccccctg cctcacaacag aaggtgctgg ggctgtgct ggcacattca gcagatgtga 1740
 atgcccggga caagctgtgg cagacaccac tgcattgtgg tgctgccaac cgggccacca 1800
 agtgtgtgga ggctctggca cccctgttga gcagcctcaa cgtggctgac aggagcgggc 1860
 gcagtgccct gcaccatgca gtgcatactg ggcattctga ggtgaggact gttcccatcc 1920
 aggccagct ggggcttttc cttttcttac cttcctactc acgtttccct gcttctgggc 1980
 cctcaagcct gaaggagaag cagcccggat gggtatataa acacttaagc tgaagcacta 2040
 aatgaatggt cagactagat taaaattctt aacttggtact tcaggagggt gataaaacct 2100
 tggattttgc gtggaaaata tttgcataca cgcctttctg aagagagtc atgtctttc 2160
 gcacatttgc aaagacttac aagagtttaa gaaccggccg ggcacggcgg ctcacgcctg 2220
 taatcccagc actttgggag gcgaaggcgg gtggatcatg agatcaggag atcgagatca 2280
 tcctggctaa catggtgaaa cctgtgtctc actaaaaata caaaaaatta gccgggtgtg 2340
 gtg 2343

<210> 121
 <211> 751
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 5405979CB1

<400> 121
 caccgtttga ctcaagtaac tttatttttc tctaaattat tttatctggc aacattttctg 60
 tggcttttaac tattaactgt aaacataaaa cagatgtgag gatgatctta ttccaaggag 120
 tggccgctgt gttccagccc cagggtgtct ggtactttcc ttcctcactc ctcacctccc 180
 acgtggcact atgcaaagct tcacattcta tcttgttctt ccttctccag tcgttcttgc 240
 tctcctgtt ctttctgcag ccgggcctgt cttctctttt cagccaagat cttcgcagcc 300
 tctcctgcat cagtgggtgc tgctgtgggc tttcccaagg ctgcgctgtt ttcggtttc 360
 cctcctgcgg cagcagcatg cttctcgtg gcatgcttgt ccactacatg cttctctagg 420

```

gcttcctctc cttgcggggc agcagcctgc tgagccaagg taccttccct ttccttggtg 480
ctcttctctt tctccgcttt cctcttgggc atgtcttggc cagaaagggc aggtaagcgg 540
tacttcacag gagaccctgg gtagggaggt ttcgttgtct ttggagactg tggataagct 600
ttagtagctg tcttggatat cacaggagaa gatggctctt taggaatgcc tccagaaaac 660
tcacatcttc tcagagggga cccgaagttc acaacaggaa gagaagttgc tgttcgttgc 720
ccccgcttca ccttctccac cagagaggcc t 751

```

<210> 122
 <211> 618
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 7481109CB1

```

<400> 122
atgatgagaa ctctcatcac cacacacca ctgcccctgc ttctattgcc gcagcagctg 60
ctgcagctgg tgcagtttca agaggtggat acagattttg atttcccaga agaagataaa 120
aaagaagaat ttgaagagtg tttggaaaaa ttttttagta cagggcccg cagaccacct 180
accaaagaaa aagtcaaaag acgtgtcctt attgaacctg gaatgccact aaatcatata 240
gagtactgta accatgaaat catgggaaaa aatgtttact acaaacaccg ttgggtggca 300
gaacattact tccttcttat gcaatatgac gagctccaaa aaatctgtta caacagattt 360
gtgccatgta agaattggaat taggaaatgt aacaggagca aaggtcttgt agaaggagtg 420
tattgtaatt taacagaagc atttgaaata ccagcgtgta aatacgaatc actttatagg 480
aagggctacg tccttatcac ttgttcatgg caaaatgaaa tgcaaaaacg tattcctcat 540
actataaatg atctcgtgga gccacctgaa cacagaagtt tcctcagtga ggatgggtgc 600
tttgtcatat cgccctag 618

```

<210> 123
 <211> 979
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 6247114CB1

```

<400> 123
gacctagagc aggcattgggt gggtcacagg ctttggagag cactctctgt cctgatcttt 60
tcagttgaga gacttcagct gtccattgct catttggact tagttcaaga attttgggtg 120
tcaccaggta aacagagccc tcagcatctg aatagaaact gaacaggaac agaagagatt 180
acactacatc tgagatggag acctttcctc tgctgctgct cagcctgggc ctggttcttg 240
cagaagcatc agaaagcaca atgaagataa ttaaagaaga atttacagac gaagagatgc 300
aatatgacat ggcaaaaagt ggccaagaaa aacagaccat tgagatatta atgaaccgga 360
tcctgttagt taaaaatacc agcctcagca tgtccaagga tgatatgtct tccacattac 420
tgacattcag aagtttacat tataatgacc ccaagggaaa cagttcgggt aatgacaaag 480
agtgttgcaa tgacatgaca gtctggagaa aagtttcaga agcaaacgga tcgtgcaagt 540
ggagcaataa cttcatccgc agctccacag aagtgatgcg caggggtccac agggcccca 600
gctgcaagtt tgtacagaat cctggcataa gctgctgtga gagcctagaa ctggaaaata 660
cagtgtgcca gttcactaca ggcaaaacaat tccccagggt ccaataccat agtggtacct 720
cattagagaa gatattgaca gtgctgacag gtcattctct gatgagctgg ttagtttgtg 780
gctctaagtt gtaaatccca cagagcttta ggactagggt cttactaagg aaggacctct 840
tcttgttcat tcttgtttaa acctttcctt aatatctact ctttagcact atagtgaact 900
cctgattatt tattctaact ggaggagtga aaaatccaaa attgtggata attcaattaa 960
aagttatgac tgataaaaa 979

```

<210> 124
 <211> 3012
 <212> DNA
 <213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 3243866CB1

<400> 124

```
gccctgcctc tgcccgctcc ccggggcgagg agccgcgggt ttcattggggc gattgcagcg 60
attccccac ccagagcgac ctgcccggcag cggcggcagc ggcaggagcc gcctttccga 120
ttccctacga tgcgggtgct gagctatggc aaagggcagc gaagtgcagc gcgagacccg 180
cgtaacgactg tgaagccac ctggagccac cttgcccggga ttgtacctgc aggcagaaag 240
tcttctacg accgtctttt cccttagagg caccagaatc cctgtaacca ttcattccagg 300
tggtgagaag atagttagca gccgagcacc catcttttga caccgtcctc tgaatcagc 360
tttgagatg ctttactct gtccgtcttc tgcagcagcc aggcagagtg cagactcctt 420
cacagccgtg aggaactctt caggctccag aagctcttaa acctgatcta caatggaaaa 480
aattcttttt tatctgtttc tcattggcat agcagtgaag gctcagatct gtccaaagcg 540
ttgtgtctgt cagattttgt ctctaactct tgcaccctt tgtgccaaaga aagggtcttt 600
attgtttcca ccaaacattg acagaagaac tgtggaactg cgggtggcag acaattttgt 660
tacaatatatt aaaaggaaag attttgccaa tatgaccagc ttggtggacc tgactctatc 720
caggaataca ataagtttta ttacacctca tgctttcgct gacctacgaa atttgagggc 780
tttgcatattg aatagcaaca gattgactaa aattacaaat gatattgtca gtggtctttc 840
caatcttcac catttgatac tgaacaacaa tcagctgact ttaatttcct ctacagcgtt 900
tgatgatgtc ttcgcccctg aggagctgga tctgtcctat aataatctag aaaccattcc 960
ttgggatgct gttgagaaga tggtagctt gcataccctt agtttggatc acaatatgat 1020
tgataacatt cctaagggga ccttctccca ttgcaacaag atgactcggg tagatgtgac 1080
atcaataaaa ttgcagaagc taccacctga ccctctcttt cagcagctc aggtactagc 1140
aacctcagga atcataagcc catctacttt tgcattaaat tttggtggaa accccttgca 1200
ttgcaattgt gaattgtgt ggttagggcg tctgtccaga gaagatgact tagagacctg 1260
tgcttctcct ccacttttaa ctggccgcta cttttggtca attcctgaag aagagttttt 1320
gtgtgagcct cctctcatta ctgctcatac acatgagatg agagtcctgg agggacaaa 1380
ggcaacactg aggtgcaaag ccaggggaga ccctgagcct gcaattcact ggatttctcc 1440
tgaagggaag cttatttcaa atgcaacaag atctctggtg tatgataacg gaacacttga 1500
cattcttacc acaactgtaa aggatacagg tgcttttacc tgcattgctt ccaatcctgc 1560
tggggaagca acacaaatag tggatcttca tataattaag ctccctcact tactaaatag 1620
tacaacccat atccatgagc ctgatcctgg ttcttcagat atctcaactt ctaccaagtc 1680
aggttctaata acaagcagta gtaattgtga tactaaattg agtcaagata aaattgtggt 1740
ggcagaagct acatcatcaa cggcactact taaatttaatt ttcaaagaa atatccctgg 1800
aatacgtatg tttcaaatec agtacaatgg tacttatgat gacacccttg tttacagaat 1860
gatacctcct acgagcaaaa cttttctggt caataatctg gctgctggaa ctatgtatga 1920
cttgtgtgtc ttggccatat atgatgatgg catcacttcc ctactgcca caagagtcgt 1980
gggttgcatc cagtttacta cgggaacagga ttatgtgctg tgccatttca tgcagtccca 2040
gtttttggga ggcacccatg ttattattat tgggtggaatc attgtagcat ctgtgctggt 2100
attcatcatt attctgatga tccggtataa ggtttgcaac aataatgggc aacacaaggt 2160
caccaagggt agcaatgttt attcccaaac taacggggct caaatacaag gctgtagtgt 2220
aacgctgccc cagtcctgtt ccaaacaaagc tgtgggacac gaagagaatg cccagtgttg 2280
taaagctacc agtgacaatg tgattcaatc ttcagaaact tgttcgagtc aggactcctc 2340
taccactacc tctgctttgc ctccctcctg gacttcaagc acttctgtgt cccaaaagca 2400
gaaaagaaaag actggcacaag agccaagtac agaaccacag aatgaagccg tcacaaatgt 2460
tgaatcccaa aacactaaca ggaacaactc aactgccttg cagttagcta gccgtcctcc 2520
cgattctgtc acagaggggc ccacgtctaa aagagcacat ataaagccaa atgcttttgt 2580
gactaatgtt gaccagattg tccaggaaac acagaggctg gagttaatct gaagagcacc 2640
acttctcctc tctctcctga aaaaatttgc cactgatatt tttactggat aaaattcaaa 2700
aatgtttcaa ttcacaaagg ctaattgttg aactgggtgc gtagaagaaa ttgtctacag 2760
gagccaagggt gaaagtctct gatgacggcg gaactggctc cattagacca tggttcatcc 2820
tcttttaaaa ccaaattttt ttttctctg gcctacaagt attttttttt taaaaaagaa 2880
aaaaagccta cattggcatc aagttctgta tcaatccatc ttacattgcc atccatgatt 2940
taacagactg tagaatcttg aataatctat atcactttaa caaataaatg ttttactatg 3000
acaaaaaaaa aa 3012
```

<210> 125

<211> 1600

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 7475633CB1

<400> 125

```
cgcgcgagga gcagccgctg ccgcccgtcg gggctcgctg gagccccaga ttcccagagc 60
ctcgggtcgc atggcagccg cttcggcgcc cggcccccg gccagctagg ggcggccccg 120
cgctccctca cggcccctcg gcggcgcccg tcggatcccg cctctctctg cggccccggg 180
cgcgccacct ccccgcgga ggtgtccacg cgtccggccg tccatccgtc cgtccctcct 240
ggggccggcg ctgaccatgc ccagcggctg ccgctgcctg catctcgtgt gcctgttgtg 300
cattctgggg gctcccggtc agcctgtccg agccgatgac tgcagctccc actgtgacct 360
ggcccacggc tgctgtgcac ctgacggctc ctgcagggtg gaccggggct gggaggggct 420
gcactgtgag cgctgtgtga ggatgcctgg ctgccagcac ggtacctgcc accagccatg 480
gcagtgcata tgccacagtg gctgggcagg caagttctgt gacaaagatg aacatatctg 540
taccacgcag tccccctgcc agaattggagg ccagtgcata tatgacgggg gcggtgagta 600
ccattgtgtg gctttaccag gcttccatgg gcgtgactgc gagcgcaagg ctggaccttg 660
tgaacaggca ggctcccat gccgcaatgg cgggcagtgc caggacgacc agggccttgc 720
tctcaacttc acgtgccgct gcttgggtgg ctttgtgggt gcccgctgtg aggtaaatgt 780
ggatgactgc ctgatcgggc cttgtgctaa cggtgccacc tgccttgacg gcataaaccg 840
cttctcctgc ctctgtcctg agggccttgc tggacgcttc tgcaccatca acctggatga 900
ctgtgccagc cgcccatagc agagaggggc ccgctgtccg gaccgtgtcc atgacttcga 960
ctgcctctgc ccagtggtc atggtggcaa gacttgtgag cttgtcttac ctgtcccaga 1020
cccccaacc acagtggaca cccctctagg gccaccta gctgtagtgg tacctgccac 1080
ggggccagcc cccacagcg caggggctgg tctgtgcgg atctcagtga aggaggtggt 1140
gcggaggcaa gaggctgggc taggtgagcc tagcttggtg gccctgggtg tgtttggggc 1200
cctcactgtc gccctgggtc tggctactgt ttgtctgacc ctgagggcct ggccgccccg 1260
tgtctgcccc cctggacctt gttgtacccc tgccccacac tatgtccag cgtgccagga 1320
ccaggagtgt cagggttagc tgctgccagc agggctcccc ctgccacgtg acttgcccc 1380
tgagcctgga aagaccacag cactgtgatg gaggtggggg ctttctggcc cccttctca 1440
cctcttcac ccctcagact ggagtgtcc gttctacca ccctcagct tgggtacaca 1500
cacagaggag acctcagcct cacaccagaa atattatttt ttaatacac agaattgtaag 1560
atggaatttt atcaaataaa actatgaaaa tccccaaaaa 1600
```

<210> 126

<211> 1001

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 1431268CB1

<400> 126

```
caatttttgt atcttttagta gagacggggt ttcaccatgt tggccaggct ggtcttgaac 60
tcttgacctc aagtgatctg cccgccccgg cctcccaaag tgctgggatt acaggcgtga 120
gccactgtgc cccgcccgga actcaggctc ttctgacca ggagcagcac ctgcttcagc 180
cactgtcttt gggctccctg ttggttgagt cacatctctc cctccatgtc taggctggag 240
tcctcagaag ctgcgtgcag ggctgtcccc tcagcctggc atactttcct cctgtcacc 300
ctttgtctcc tccttattca agtctgggccc caggggcctt ctctgcaggt cgtaactaaa 360
gtcgcacctc ctgccctaac ctccagcatg totgactctt tggatttcac caagcacttc 420
tcactttgca aagtcattga ttctgcaaat gttcatcgag gatgtactac gtgccaggct 480
ctggttaagg cacgggatgt agaacaagt tgctgtcgtt ttccagctca tgctctggct 540
ggagaggcgg tcagtgcaga gaataagcaa agaggcggag aggctgcgtc ctgcctcctc 600
agatgagcac tgtgtgtgcc agcacggtag ccgctggcca cagggtgagat tttcacttaa 660
gttgaatatg gccgggcgca gtggctcatc atgcctgtaa tcccagcact ttgggaggcc 720
gaggcaggcg gatcacgagg tcaggagatc gggaccatcc tggccaacac agtgaaactc 780
cgtctctact gccagggtgc tgggtgcagag gaggcattcc atcaattgaa ccttaaggaa 840
ctctggaggc aggggatatg aatgtttact aagtgtgtac atcacattaa gtacagaaat 900
cgtctggtgt cactacttca gtgtcaccgc tgcaacatcc agccgagggt gtcatgtcgt 960
ttgcagagca ggaaactcag cctcagaatg gttgccttgt c 1001
```

<210> 127

<211> 1424

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 2414185CB1

<400> 127

```

tcatgggggtg gaataaacac agaaaagact gaaaaagccg ggccaatggc tcaagcccgt 60
aattcccagc acttgggagg ccaaggtggg cggatcacct gaggtcagga gttcaagacc 120
agcctggcca agatggtgaa atcccatctc ttactaaaaa tacagaaatt agccaggcat 180
gggtggtgggt gcctgcaatt tcagctactc aggaggctga ggcaggagaa ccgctggaac 240
ctgggaggcg gaggttgcaag tgagccgaga tcccggccact gcgctccagc ctgagccaac 300
aacagcgaga ctccgtctaa aaaaaaaaaa agaagactga aaaaaaatat gttggagacg 360
actttcttac tgtggaaggg ctgagattcc atgatgggaa agcttagccc cacctttatt 420
ctgggcatttt gttgggtacc tgctggtttg ggataccagc agggaaagaa aacatggccc 480
ttgtcctcca gcccttaca cctgcaagac aagatgtacg ccttgagaa agctggagat 540
ccttccaagg caaggagcat gggccctcac aaatctccag aaaccagag agggcaacct 600
atggagatgt ctggcttaaa gggccagggtg acatccacag ccttacacac tctgcatttt 660
cctagaaggc ctccctctgg ctgtcagaca gatcaggctg gtgaccacga gccaggggga 720
cgctttctgg cccaaccaca gagacttagg gaactttcat tgatgatcag tcccctgcaa 780
cttcttcctt ttggtagccg ctgaaagtag gtcataaag ggcctagacc tcaggatgtc 840
tggggagaac ctcccttgag tgccagaaga ttcttgcatc cagggtcttg tggctaaggc 900
tcatgcttac tggcagtgct ggcctgatgg gttagaacc gcctggacca cctacgtggg 960
atattttgg gatgctgaga cgtttctcca aaataccttc tggaatggcc tgtatcagta 1020
gctgaagtca taggatcatt cctcccagct aaagaaatgc ctgctattaa atttttgtgg 1080
ataagaggaa gagacatagc ttgtggacgt gcagttgggt gaagtcatga acatactgac 1140
taatgatacc cagggtgaac tgctgactaa tttgtgtaaa cagagtggat ggtctctgat 1200
ggtagccct agggctctgt tccggcctcc ccacatgta tcaggaatgt tcatggaaat 1260
taaaaggtat actgataaaa tctatgggag atacaaaact aggagagatt gctggagtaa 1320
taaaattatg gaatgaactt tcagaaaaac tgctgtaggt tgaaataatg ggccaagct 1380
gtaacatatt agggatgaat agaaggtcct atatcaaagt tcat 1424

```

<210> 128

<211> 1282

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 5266594CB1

<220>

<221> unsure

<222> 9-10, 23

<223> a, t, c, g, or other

<400> 128

```

aacaaaaann cccaaaacca gtnaaaagcg tgcgctgacg gtttgccatt taaggattac 60
agcaaatgat tctatttgta atttctaccc ttcccattgc ctcccccca aaaaaaactg 120
tacctgagtt tacaaacata ttaacatata aataatgaga accgtcctgg tgggagcctc 180
ctccgttgct tctgctggag atgaacactg aggggcgctg taaccacaca gactgcctgt 240
gacatcgga gtctcacggc agctgtcctg ggcccgcagc tggctttttt gccacctcca 300
ggttcaacca ccagtctgtc tctgctgtgc ccagggtaga gcccgggggc tgtgagtatg 360
tgtggctccc ctgcccgta tgcctctggc tcaagctcat gctggaagg acgcttctc 420
tcccagacgt gcttcttggt ggcaggccag tccttctgct ggcactggga gccgcagtac 480
cgggccacct ggcagcgccc acagatgttg aactcccga gctgcttctc aatcactgtg 540
caggagggt agtggcactc atagtaggtg caagagttct cctcctcttc cactacatcc 600
ccgttggcat tataataccg ggtcacattt aggacaggaa actgttcttc tatagggtct 660
gtaggcagct ctgaattgc aagccaatac aagcttctgc ccctctgctt gctggaggtc 720
tcttcagcct tgggccgcca gcccacggc accagttgca ggttgctccag gcgattgtcc 780
acggtcacag cgttgagggtg caccacctgg aagcccggg ccacgcccc ccggtgccgc 840
tcccacagca gctcatgaag gagtctccca gageccctc ctcggttctt gtcaaaggca 900
taggcaaata tcttagcacc atttccatct gcattcactt ccattcgggc ctcaaaggag 960
tagctctcca ccagcgggat gtcctgtcca tcgatcagcg tgtatttggg cttcccggcc 1020
acccggccga gccgcacgat acccaatttg aagtcggtca tggccggggc tgcgcttcgc 1080

```

```

cggcagcccg ctccctcggg aggcgcgcgac ggggcccggg cgagccgcgg cgcccccgtta 1140
cagacggggac cgagccgggt cgggtacacc agcgggctcc gggcggacga gctggccggc 1200
tcggcctccc gccgcctgcg ccgcgcgctt cagaactgcg ccgtcgagat gattaaagcc 1260
atgccctata tatattacgc gc                                     1282

```

<210> 129

<211> 642

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 7610617CB1

<400> 129

```

cggctacagc tgcaagtcaa agaggtcaga ggccttctga tgttggcgct ggaagcctct 60
cctttcttgc cccacaggtt atgggggctt ctccactccc cgctcttact tgcctgcttc 120
tccagggatc cctagaagag cagtgggttag caagcatatg ctatagccat caaatacgag 180
tctacagaaa tgtaggcttc cccttggggc tgagttaggg cctctgcttc tgaaatgagc 240
aacctctggt tgctgggtgg ggctagggtt tgctcccttt ctctctcac atactctttc 300
ctgggtgacc tcatcccctc ccattgcctt aaacatcttc caggtagctg tgtcatccac 360
ttgtgtagtt ccagttcaga aatcccgtca gctccattca ttcatttggt cattcattca 420
gcaaacatat gtggcatctc tgtcccaggt actgctctac aacctgggtg tacaataggg 480
acccaaacag acactccctt ccccatgcac tccctgctca cagacacacc tgcgtggcag 540
tgcttgggtg tcttcacagc tccctgatgc cgagctgtga tcccaccctt ctgcctccct 600
ggccccctc agcgtgctgc caccattcac cttgaggctg ag                                     642

```

<210> 130

<211> 1326

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 1902436CB1

<400> 130

```

ctcacctcag cctcctaagt agctgggact acagggtgcct gacaccatgc ctggctgatt 60
tctgtttgtt tgtttgtttt tttttgtaga gatggggtct tactttgttg cccaggctgg 120
tctcagaaat gactttttata tatacttttta ttttaagttt ttttcttcaa ctatgtttgt 180
cttttatgaa gttaatcttg ctcatcagca acacaaatgc tgtatctttt atactacaca 240
ggccctgtac cctttgctct gatttctact ccataatttg tatgctgtta actgtctctg 300
tgaatttctt gtcatttttg aataattttc aaactattct tacttgggct gatctgtttt 360
caatgctttt ggcatacgaa tatagattta caagactctt ctcagtgtca cccacactt 420
ctgtcatgtt atgttttttaa aaattctctg attgagctat aactgcttaa agatgtgaaa 480
tgagtatttt caatagattt agtttttttca gggggtaatg acagacttgt taaaaaaaaa 540
aggttataaa tacaaaaagt ataaagaagg aaaaacctat gcaaaatctg agcagagtta 600
actttctggt gaatgtcatt cccgtctctt agttatgtgt gtataattat gcataataga 660
gattctattg tatccaagga ttttttaaac ctttttgagg tgagatctgg ctgtgttgcc 720
caggctggag tgcaagggtt gttcataggt aagatgatgg tgcactatag ccaaggactc 780
ctggactcag gtgatactcc tacctcagcc tccaagtaa ctgggactgc tggcacacac 840
cactgcacct ggcttaagtg ttttttccaa gaagtcaaag gctaattgatt ctaaaagaaag 900
ttagtgtaca gaccttata catgagtatt atgtgcatct tctttagtgg cagaattctg 960
ttttaagtat tataattgga aaaacaactg aagagacatt taaggaacac cactatttca 1020
gcagacagtt gatagaagat accttttaaa agaagtgggt gaatgtcaca catttgaaac 1080
ttaaaaaaag ataaacccaa aacgtcagta gaagggatgc cagagggaat aatcttgatg 1140
gcaagcagat agatttgata agccaatggg atttccact tctctttgct atttttctat 1200
tttgagccc tacttgattt aggcaagtat catgagtgga tgctaaaagt acttagtaaa 1260
agtttgatga ggaatagaat gtttattcaa tcccaaagta tactcatatt atcaccaaaa 1320
aaaaaa                                     1326

```

<210> 131

<211> 1486

<212> DNA
<213> Homo sapiens

<220>
<221> misc_feature
<223> Incyte ID No: 2310369CB1

<400> 131
cccaaacaga gcgttttctg gtcgcgaac tcttcccggc tccaccgca cctacagcct 60
ccaaccaaacc cccaccaagt ctcgaccctc tccccaggc ctcgaccctc ccgctcgccg 120
ctgcgttggg accgcgcacc cggtaactgc gcacggctgg cctcccggag tctgtcgccg 180
tcgctcctcc tggaccggg ttcccaggac aacgcctccc ggaacgcagg gagcaggccg 240
aggccgcgcg gtgggctcgc agcgcctcgc cgctctctct tccccaggcc cgcactccca 300
cttgggctcc agggccccag ggctgacgtt cccccagct tagaccctga gtcgttttcc 360
cccgtttccc ggctgaatta ggttcttctt ctccacaggt gtgtgcagt gcctcaggga 420
tccggaaagt ctaggactga acttctccta acatccagta atggggacct ggaacctggg 480
cgtactagag tgccgcgcgt agggctccag gtcgctggct tctgcgtttt cttcctctcc 540
aaagttagag atctcctatc tgtgtctca tacatactgc cgcctgagggt gccatggccc 600
caaagccggg ggcgagtggt agcacagccc tgtcccatct ggtgctggga gtggtgtctc 660
tgcacgcagc cgtgagcaca gccgaggcaa gtcgaggggc tgctgctggc ttcctgctcc 720
aggtcttggc tgccaccacc acgctggccc cagggtgtag cacacatgaa gactgccttg 780
ctggagcctg ggtggccacc gtcatcgccc tcccccttct ggccttcgat ttccactggg 840
tgaatgggga ccgctcctct gccaacctgc tcttgggagg aggcattggt ctggcagtggt 900
ctggcgccca cctcgccctt gaggccgct ctgtggctgg tcaggcaatg ctggtggtgg 960
tcgcagtgc cactctcatt gtagctgtct tcacggccaa cacttatggg atgtgggggg 1020
ggcgatgct ggtgtgga gccctcctga gccggctgga ggaggacagg ctgctgctgc 1080
taccgaagga ggatgtctgt cgctgggcct tggtgttagg cagctgggct tactgccggg 1140
ccctgcatac acagcgccct cagtgggagt gacagttgga tacagccagg cagggtttct 1200
gccctgcgca acactttccc tcccacctgc ctgctcctgg cgccttctcc ctaggggtag 1260
actcttctgc ctactgaagt ggggttgcgt cacattgact ggtcaggggc agagtctggg 1320
tgctgtcctt tggccacgtg tggggacttg tctagaccag aatgaaaggg acagggtccc 1380
agacacgttt ggggtctctg attctgggct ggacacggtt gtggatccag agaagaggcc 1440
tagtctccaa taaatcttag gaattttgca ggaaaaaaa aaaaaa 1486

<210> 132
<211> 1523
<212> DNA
<213> Homo sapiens

<220>
<221> misc_feature
<223> Incyte ID No: 6180576CB1

<400> 132
ggcgcggtgg cgtgtcagga ggccgggcag ggggtgcggg ggaccgacgg acgcacgggc 60
ggcgcgccgg gagccatgga gcgcggccct ggggcccggg ggcgcgccgg ggggtgggct 120
tcccacggca cgacatggag acctgtggtt gcgaggctcc ctggggctcg gcttggaccg 180
cgatggggct ggccctggc ctctaaccg ggctgtgtgc tggggcggtg gctggggggg 240
cgctctcccc cctgcccgcg actcggagca cccccacccc tcccctgccg ggccaggccg 300
ggcgcggttg ttggcggggg ccccggtgga ggcccgccct ggcgcgccgg cgcctgaat 360
gggctgtcgc tgagtgaagt ctgctgcctc ttctgtgcc cgcctgccc cggccgcac 420
gctgccaagc tcgccttctt gccgcgggag gccacctact cctggtgcc tgagcccag 480
ccggggcctg gtggggccgg ggccgcccc ttggggaccc tgagagcctc ctggggcgca 540
ccggggcgtt ggaagctgca cctgacggag cgtgccgact tccagtacag ccagcgcgag 600
ctggacacca tcgaggtctt cccacccaag agcgcccgcg gcaaccgct ctctgcatg 660
tatgttcgct gcgtgcctgg tgccaggtag acggtcctct tctcgcagg caatgccgtg 720
gacctgggccc agatgagcag cttctacatt ggccctgggt cccgcctcca ctgcaacatc 780
ttctcctacg actactccgg ctacgggtgcc agctcgggca ggccttccga gaggaacctc 840
tatgccgaca tcgacgccgc ctggcaggcc ctgcgcacca ggtacggcat cagcccgag 900
agcatcatcc tgtacgggca gagcatcgcc acgggtgccca ccgtggacct ggccctcgcc 960
tacgagtgtg ccgcgggtgt gctgcactcg ccgctcacct cgggcatgcg cgtcgcttc 1020
cccacacca agaagacct ctgcttcgac gccttcccta acatcgagaa ggtgtccaag 1080
atcacgtctc ccgtgctcat catccacggc acggaggacg aggtgatcga cttctcgcac 1140

```

gggctggcgc tctacgagcg ctgccccaaag gcggtggagc cgctgtgggt ggagggcgcc 1200
gggcacaacg acatcgagct ctacagccag tacctggagc gcctgcgtcg cttcatctcc 1260
caggagctgc ccagccagcg cgcctagcgg cggcccaac cggccggacc tcagcaataa 1320
ggcgggcccc ggacctcacc ccgcgcgggc cccacccag gggctgcatg tggaccccc 1380
gggcggccca ggggaccccg ccccgaccca ggggtgtggt acgatgtaca ggcaacagag 1440
ctacgcactc ctttcctttt ggaagcaaga agaaaatacg tgaaaacgga aattaaagat 1500
ttaaaatttt aaaaaaaaaa aaa 1523

```

<210> 133

<211> 848

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 2274523CB1

<400> 133

```

tagttctaga actcctgacc tcttgatctg cccacctcag cctcccaaag tgttgggatt 60
acaggggtga gccaccacgc ccagcccagc agtggcttct gtgtgggaag aagggtcagg 120
ctgaaggacc ctgggaggat gccccacca acatgtcttc tctcatcggt gggtaattgc 180
ccatgtttgc ccagccattc tcacctatca gggcttccaa aagaatggcc aaagtaagtt 240
ctaacaactt tgcttctttg ccaaggcaag caccaatgct acttttctgc cctctctgga 300
tgcccggtgac ttccgttcca caggaagcca agctcctgag gcagctgaaa ttcagtcagg 360
ggactggggt ctgtgtactg atctacacc cacttcatac ctacttcttc aaactgtccc 420
caacactggg aacacctgtc cttgaatatc ctacgaatt taaggggaaa aaaagggtga 480
agcagaaaaga ctttttctct ccaaaattgt gcctgcttgc ttgggggcct aggcattgag 540
acttgaaaat aaatcaagct tgggtggggc atggtggctc ccgctgtgta tcccagcact 600
ttggaaggcc gaggcaggcc aggcagggtg agggaggatca caaggtcagg agatcgagac 660
catcctgggt aacacggtga aaccctgtct ctactaaaaa taaaaaaca aaattagcca 720
gggtgggtggc ggggtgcctgt agtcccagct gctctggagg ctgaggcagg agaatggcat 780
gaaccaggga ggcaaagcta gcagtgcgcg gagatcgcac cactgcactc aagcctgaca 840
taattgcg 848

```

<210> 134

<211> 2758

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 1801820CB1

<400> 134

```

cggcgggggc ggcgcctgac tggaggggat gggctgccgg gcgcgtaggg gccatgccgc 60
ccgggacccg ggcctgccgc gtccgcgcgc ccggccgcgg cgccccacgt ccgcgcggg 120
atggtgaacc tggcgcccat ggtgtggcgc cggcttctgc ggaagagggt ggtgctcgcc 180
ctggtcttcg ggctgtcgct cgtctacttc ctcagcagca ccttcaagca ggaggagagg 240
gcagtgcgag ataggaatct cctccagggt catgaccata atcagcccat cccgtggaaa 300
gtgcagttta acttgggcaa tagcagtcgt ccgagcaatc agtgccgcaa ctccattcaa 360
gggaagcacc tcatcacgga tgaactcggc tacgtttgcg agaggaagga tttgctggta 420
aatggctgct gtaatgtcaa cgtccctagc acgaagcagt actgctgtga tggctgctgg 480
cccaacgggt gctgcagcgc ctatgagtac tgtgtctcct gctgcctgca gccaacaag 540
caacttctcc tggagctctt cctcaaccgg gcagccgtgg cattccagaa cctcttcatg 600
gcagtgcgag atcactttga gttgtgcctg gccaaatgca ggacctcatc tcagagcggtg 660
cagcatgaga acacctaccg ggaccccata gcaaagtatt gctatggaga aagcccgccc 720
gagctcttcc ccgcttgacg ggtgcagcgg acttgctcca gcctgggtga ggaggccccg 780
ctgaagaact cgcctcctgg gacccagctt cagccatcgg gccaggctgc aggaagaaga 840
caaaggcagc gtgaggaaac cttggctttg accccttctc gtgttgtcat ctttggcttc 900
gctcaccacc cgggcttacc agatggaaact cttctgtaaa gcagcttggc ccctccagcc 960
agtccattc gggaaagatg aaaccggagg ccgggctcac ggtggtgggt gagttcttgg 1020
atgactcagc cctgggacct gcacgggac ctgtgacttg tgttcatcgg gggccggtgt 1080
cacttccagt tttgatccag gctctttcac tgtaaaatta tttattggat tcctttggag 1140

```



```

taatgggaac attctaattgt tttatgtagg aaaatgcctt gccattctag ttgaatatgt 1200
tcaaggaaat tatttttgtt gttgttctgt gttctcgagt ttcaggagtt aaatcattct 1260
tccacccaga tacaacattt tctcttttag gacgtgaata ttctctctag gcagttattt 1320
ttgtttgtat ttcgacagta tcaagcatag gccctgaacg tgacctgtta gccatatact 1380
gacgtgtaaa attatctaaa aactcagaca ctcttccatt ctaatctaac tgcacgattt 1440
ctaacagtgg gcaccatgtg cctgccctca ggttattttc cagtggtagt cgaatgtgct 1500
catataccct atggagagca ctgttttagc agaaatctaa ttctcttcc tggaggaaat 1560
tgttctcatt tcttttgcca cttaaaatta actgtgggct actcagccag ggtacagtgg 1620
gagcctcagg aaggtcagag gcaacctcct cccctgttct atcaatagaa acccaacgtt 1680
gaggcaattc ctaaacagac gcacctcgta gcttgctgta tgtgtttatt ctttattgct 1740
ttcagctttg gggctgtaac aggtacaaat atttggtttc cctatgattt atagagaaga 1800
agaagaaacc cagctttcta tcagagcact gcaagagaag agtcttacac ctgccctcag 1860
tgggagatga gaatggtcat tatgacttag agaatgctac acgtgtaggt tgctggtgtg 1920
tctgaatcc acaggcataa agcactcccc atttttctta ctgtaatgca gattctccgg 1980
ctcaaggtct agaataattg atcctaagat caagacatca tgcccttcga atagtactgc 2040
tctttgtttt caggagtcac gtgaacacac aactctccta tttctctcac gaacctcagg 2100
attgagcaag gtctttgtaa tttttttggt tctactttatt gacctgggag caaggtgcta 2160
attctgtggt cagtattcaa tgttttttct agtggagctt tttctttggg ccatatttgc 2220
cttctaatac attcctgcaa tatgtagtgg tgatttcct tagcttcctc ctactacctc 2280
ttatactcat ccccccaat tatttgcctc ccttaataaa gttttcctag aaggtaagct 2340
ggtcaggcaa tttgaaaaat attagatccc aagaaatcta ttccgtttgc attggacttc 2400
tcggattcca tgtgtttgca gcaggactac atcgaaactc gatgtgccgg attgtggcag 2460
gtctgcattg ctcattccatc tattgttttt ggtaactcag tttggaattt cagtgtctgt 2520
cttccctggg ttgacattgg aatcagcctc tcttttgagc ttattttaac tcttgagcaa 2580
cataacatag atttaatgtg aacagtttat accaaagggc agcctgtgcc tgtttatgga 2640
tcctctctgc ctttgtactt gaagagcgca ttttacattt ccagtccttt cacagacagg 2700
agctccaacc ttacgatgga gaattaaact tgcttgattt tccactttgt ggatgagg 2758

```

<210> 135

<211> 1650

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 3010920CB1.comp

<400> 135

```

gtcagcatta cttatgtttc atcctgtggt cgtgtgcttt ctgctgttg ctacattcaa 60
catccaaatt gtaatttata tttataactg cctctcatct aattatattc tttctcgttt 120
ttaaaccaca gcaaaatcaa ctttagctag ccattatact aaaacatact acattttctt 180
ttaaacttct tttagtttgt ttctcttaga tggtaaaact tactcaggca gggattatgt 240
ttgattattg atgtgaatgc aactttttag aacatagctg cgtactcagt ggactcagta 300
catactgacc agttgttgaa aaatctgatg cagcctccct aagtgtttag catagccaga 360
gtagtttacg tattcgttaa atctgactca gtaattagca gacctagtgt ttaagtactg 420
gggaaatacc tgcttgccct tcacagatgt caagaaagta cctggtagaa gagtcttacg 480
acacagccta acttgtgtag gacagaagat accatctgga cttttagtaa cccagaaata 540
caagtctaaa aatatcttga taacaaattc tttgaaatat ttttgaaggc cctgagcact 600
tcactttttt ggctatatta tttaggaattg tttgatctct gcaatttcaa caatgattac 660
agtaatcact gtcattatca agcatttact atggggccaga ccttttctta ggtactttac 720
atacaatata ttgttctggt aaaaacttaa caaggtttgt attattattt tacaaatgaa 780
gaaactaagg ccgaggaggg ttaactaact tgcccaggat cacacgtatg gtaaaggcag 840
aagtaaatac ttgagatatg gtttgttgag ttctaaaact taggtgaatt ttacctccta 900
atacgtctca aggatgtggg tactggagga agaactgtac tgaaatccaa atatttggtg 960
ctagctgttt tgtataatac tagtaaatat tcatattgcc cttatggtat tgcttatgat 1020
catacaataa ttggttcttg cttatataga aatctacagt tttctgtagt gaatatatgt 1080
gttacagcaa ccacaactat ttatctgaaa tatcagtaaa tatttatcat aggaatgaac 1140
atatattaag tcttgacaga gctctcgagg cataatgact tggttgaaag tagcagattt 1200
cagtaagaac catgccaaaa atattaagct ataagttatt agtgtacctt gttttaatgg 1260
caaacatgat gattttctta gtgccatat atgaataaaa taacttacta gaagtttgta 1320
gacaggtttt tttttccccc tctggagaca gctcatgctt ttgttgccca ggctggagtg 1380
caatggcata attttggctc actgcaacct ctgcctccca ggttcaagca attctcctgc 1440
ctcagcctcc cacgtacctg ggattaccat gcctggccta tttttgtatt tttagtagag 1500

```

```

atgaggtttc accatgtttg cctggctggt ctggaactcc tgacctcagg tgatccccc 1560
gcctgggcct cccaaagtgc tgggattaca ggccacctca cctaggctag atagttttat 1620
tcacttggtc gtttcaccaa aaaaaaaaaag 1650

```

<210> 136

<211> 1061

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 3409459CB1.comp

<400> 136

```

aaaataaagg ttcaagctgc caaatTTTTt tccagggtct gtttgctctc cctccagaga 60
tgtgctgagt ggccgtggct tcttcccacc caccgtgtcc tgctcttccc tgttcttggc 120
ctaagtcttg cacaggagcc tgtcgaggcc cttgggacac ttccagaagc caggtccatt 180
ttctccatga acaggtgggc cagggttgag tctcccccta gttggactgg gagcacatgt 240
ggtaagatgc caggggaagc tccgtgttta tcctccacat gcctggaagg atgcagtga 300
agcctttgcc tggcgtggtt tctttctggc ttgaagtttc tgccaagctt tagggccctg 360
ctggcctggg aagcagcgtt ggcaggcctg ccttgcgat ggacgctgc agcactggca 420
ggggggcagt ccaggcgga ggggcagggg aggatgccct tcctggccaa atcccttgg 480
taagtaagta ccagcatctg aggtggcccg ggtgcagggg aaccactatt gctacctctc 540
cagcagcagc cagtgggtgag tctgacatgt cagctgcatt cagaccaaac ccgcccccg 600
caaataccca gagactgagg caggaagctc tgtagaaatt cattttattc ttattcagac 660
tattttcaaa agaagcagtg gtgtgctgtt tttctaaaaa tatgccttta tagattttta 720
tatatgtata ttataaaatc catacatgta tttacatgat tgctacatac aaaattacag 780
cactgtggta tgtacacatc tacaggtaca ttcttgccgc gcacccctgc tgtgctttcc 840
ccacgtgagg gagggaggga gactgaatcg gttgtgagca gctgagggct ggccggggccg 900
cggagccttt gatttggggc ctgggtggag gaggaggggc ggctgcttca catcactctc 960
cagctctgac actcggcctt cccactcccc ttctcggttc tcaaatcgtc ttctcaattc 1020
atgtttctct tgctctaagt gctcgagctt gtctgtcagc t 1061

```

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
4 April 2002 (04.04.2002)

PCT

(10) International Publication Number
WO 02/026982 A3

(51) International Patent Classification⁷: **C12N 15/12**,
C12Q 1/68, C07K 14/47, 16/18, A01K 67/027, A61K
38/17, 39/395, B01J 19/00, G01N 33/50, 33/53

(21) International Application Number: PCT/US01/30042

(22) International Filing Date:
25 September 2001 (25.09.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/236,869 29 September 2000 (29.09.2000) US
60/239,812 11 October 2000 (11.10.2000) US
60/240,108 12 October 2000 (12.10.2000) US
60/241,282 17 October 2000 (17.10.2000) US
60/242,218 20 October 2000 (20.10.2000) US

(71) Applicant (for all designated States except US): **INCYTE GENOMICS, INC.** [US/US]; 3160 Porter Drive, Palo Alto, CA 94087 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **YUE, Henry** [US/US]; 826 Lois Avenue, Sunnyvale, CA 94087 (US). **TANG, Y., Tom** [US/US]; 4230 Ranwick Court, San Jose, CA 95118 (US). **NGUYEN, Daniel, B.** [US/US]; 1403 Ridgewood Drive, San Jose, CA 95118 (US). **YAO, Monique, G.** [US/US]; 1189 Woodgate Drive, Carmel, IN 40633 (US). **XU, Yuming** [US/US]; 1739 Walnut Drive, Mountain View, CA 94040 (US). **TRIBOULEY, Catherine, M.** [FR/US]; 1121 Tennessee Street #5, San Francisco, CA 94107 (US). **SANJANWALA, Madhu, S.** [US/US]; 210 Sylvia Court, Los Altos, CA 94024 (US). **CHAWLA, Narinder, K.** [US/US]; 33 Union Square, #712, Union City, CA 945877 (US). **BAUGHN, Mariah, R.** [US/US]; 14244 Santiago Road, San Leandro, CA 94577 (US). **SAPPERSTEIN, Stephanie, K.** [US/US]; 3531 Highland Avenue, Redwood City, CA 94062 (US). **LAL, Preeti** [IN/US]; P.O. Box 5142, Santa Clara, CA 95056 (US). **THORNTON, Michael** [US/US]; 9 Medway Road, Woodside, CA 94062 (US). **GANDHI, Ameena, R.** [US/US]; 837 Roble Avenue #1, Menlo Park, CA 94025 (US). **RAMKUMAR, Jayalaxmi** [US/US]; 34359 Maybird Circle, Fremont, CA 94555 (US). **ELLIOTT, Vicki, S.** [US/US]; 3770 Polten Place Way, San Jose, CA

95121 (US). **ARVIZU, Chandra** [US/US]; 490 Sherwood Way #1, Menlo Park, CA 94025 (US). **THANGAVELU, Kavitha** [IN/US]; 1950 Montecito Avenue #23, Mountain View, CA 94043 (US). **GIETZEN, Kimberly, J.** [US/US]; 691 Los Huecos Drive, San Jose, CA 95123 (US). **DING, Li** [CH/US]; 3353 Alma Street #146, Palo Alto, CA 94506 (US). **AU-YOUNG, Janice** [US/US]; 233 Golden Eagle Lane, Brisbane, CA 94005 (US). **TRAN, Bao** [US/US]; 750 Salberg Avenue, Santa Clara, CA 95051 (US). **POLICKY, Jennifer, L.** [US/US]; 1511 Jarvis Court, San Jose, CA 95118 (US). **LEE, Sally** [US/US]; 825 East Evelyn #425, Sunnyvale, CA 94086 (US). **LU, Dyung, Aina, M.** [US/US]; 233 Coy Drive, San Jose, CA 95123 (US). **BURFORD, Neil** [GB/US]; 105 Wildwood Circle, Durham, CT 06422 (US). **WARREN, Bridget, A.** [US/US]; 10130 Parwood Drive #2, Cupertino, CA 95014 (US). **GURURAJAN, Rajagopal** [IN/US]; 5591 Dent Avenue, San Jose, CA 95118 (US). **DUGGAN, Brendan, M.** [AU/US]; 243 Buena Vista Avenue #306, Sunnyvale, CA 94086 (US). **HONCHELL, Cynthia, D.** [US/US]; 400 Laurel Street #203, San Carlos, CA 94070 (US). **HAFALIA, April, J., A.** [US/US]; 2227 Calle de Primavera, Santa Clara, CA 95054 (US).

(74) Agents: **HAMLET-COX, Diana et al.**; Incyte Genomics, Inc., 3160 Porter Drive, Palo Alto, CA 94304 (US).

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report

(88) Date of publication of the international search report:
25 September 2003

[Continued on next page]

(54) Title: SECRETED HUMAN PROTEINS

(57) Abstract: The invention provides human secreted proteins (SECP) and polynucleotides which identify and encode SECP. The invention also provides expression vectors, host cells, antibodies, agonists, and antagonists. The invention also provides methods for diagnosing, treating, or preventing disorders associated with aberrant expression of SECP.



WO 02/026982 A3



For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 01/30042

A. CLASSIFICATION OF SUBJECT MATTER IPC 7 C12N15/12 C12Q1/68 C07K14/47 C07K16/18 A01K67/027 A61K38/17 A61K39/395 B01J19/00 G01N33/50 G01N33/53		
According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) IPC 7 C12N C12Q C07K A01K B01J G01N		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the International search (name of data base and, where practical, search terms used) BIOSIS, EMBL, SEQUENCE SEARCH		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P, X	DATABASE EMBL 'Online! EBI; 23 May 2001 (2001-05-23) OTA, T. ET AL.: "Human membrane or secretory protein clone PSEC0241" Database accession no. AAB88452 XP002222315 abstract	1-20, 23, 26-47, 56, 123
P, X	- & EP 1 067 182 A (HELIX RES INST) 10 January 2001 (2001-01-10) *SEQ ID NO:272* the whole document	1-20, 23, 26-47, 56, 123
<div style="text-align: center;"> <div>---</div> <div>---/---</div> </div>		
<input checked="" type="checkbox"/> Further documents are listed in the continuation of box C. <input checked="" type="checkbox"/> Patent family members are listed in annex.		
* Special categories of cited documents : <div style="display: flex; justify-content: space-between;"> <div> *A* document defining the general state of the art which is not considered to be of particular relevance *E* earlier document but published on or after the international filing date *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) *O* document referring to an oral disclosure, use, exhibition or other means *P* document published prior to the international filing date but later than the priority date claimed </div> <div> *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art. *&* document member of the same patent family </div> </div>		
Date of the actual completion of the international search		Date of mailing of the international search report
25 November 2002		0 8. 05. 03
Name and mailing address of the ISA European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo nl, Fax: (+31-70) 340-3016		Authorized officer Hillenbrand, G

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 01/30042

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	<p>DATABASE EMBL 'Online! EBI; 23 May 2001 (2001-05-23) OTA, T. ET AL.: "Human membrane or secretory protein clone PSEC0241" Database accession no. AAF93879 XP002223091 abstract</p>	1-20,23, 26-47, 56,123
P,X	<p>-& EP 1 067 182 A (HELIX RES INST) 10 January 2001 (2001-01-10)</p> <p>*SEQ ID NO:271* the whole document</p>	1-20,23, 26-47, 56,123
T	<p>DATABASE EMBL 'Online! EBI; 6 November 2001 (2001-11-06) OTA, T. ET AL.: "Human polypeptide and full-length cDNA" Database accession no. AAM93427 XP002222316 abstract</p>	
P,X	<p>-& EP 1 130 094 A (HELIX RES INST) 5 September 2001 (2001-09-05)</p> <p>*SEQ ID NO.:3054* the whole document</p>	1-20,23, 26-47, 56,123
T	<p>DATABASE EMBL 'Online! EBI; 6 November 2001 (2001-11-06) OTA, T. ET AL.: "Human polypeptide and full-length cDNA" Database accession no. AAK94348 XP002223092 abstract</p>	
P,X	<p>-& EP 1 130 094 A (HELIX RES INST) 5 September 2001 (2001-09-05)</p> <p>*SEQ ID NO.:3053* the whole document</p>	1-20,23, 26-47, 56,123
P,X	<p>DATABASE EMBL 'Online! EBI; 8 February 2001 (2001-02-08) SHIMKETS, RA. AND LEACH, M.: "Human secreted Human ORFX ORF1664 polypeptide sequence and polynucleotide sequence" Database accession no. AAB41900 XP002222317 abstract</p>	1-20,23, 26-47, 56,123
P,X	<p>-& WO 00 58473 A (CURAGEN CORP ; LEACH MARTIN (US); SHIMKETS RICHARD A (US)) 5 October 2000 (2000-10-05) *SEQ ID NO.:3328* the whole document</p>	1-20,23, 26-47, 56,123

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 01/30042

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	DATABASE EMBL 'Online! EBI; 8 February 2001 (2001-02-08) SHIMKETS, RA. AND LEACH, M.: "Human secreted Human ORFX ORF1664 polypeptide sequence and polynucleotide sequence" Database accession no. AAC76109 XP002223093 abstract	1-20,23, 26-47, 56,123
P,X	-& WO 00 58473 A (CURAGEN CORP ; LEACH MARTIN (US); SHIMKETS RICHARD A (US)) 5 October 2000 (2000-10-05) *SEQ ID NO.:3327* the whole document	1-20,23, 26-47, 56,123
A	WO 99 40100 A (HUMAN GENOME SCIENCES INC ;ROSEN CRAIG A (US); KYAW HLA (US); LAFL) 12 August 1999 (1999-08-12) the whole document	1

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US 01/30042

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
see FURTHER INFORMATION sheet PCT/ISA/210
2. ☒ Claims Nos.: 21-22, 24-25, 48-55
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
see FURTHER INFORMATION sheet PCT/ISA/210
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this International application, as follows:

see additional sheet

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:
1-20, 23, 26-47 (partially), 56, 123

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.1

Although claim 19 is directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.

Although claims 33 and 35 are directed to a diagnostic method practised on the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.

Continuation of Box I.2

Claims Nos.: 21-22, 24-25, 48-55

Present claims 21-22 and 24-25 relate to an extremely large number of possible agonists or antagonists and methods for use of this agonists or antagonists. Claims 48-55 are extremely broadly and imprecisely drafted product claims directed to an array comprising different nucleotide molecules. In fact, the claims contain so many options and variables that a lack of clarity and conciseness within the meaning of Article 6 PCT arises to such an extent as to render a meaningful search of the claims impossible.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. Claims: Claims 1-20, 23, 26-47 (partially), 56, 123

Invention 1:

An isolated polypeptide of SEQ ID NO: 1 and polynucleotide SEQ ID NO: 68 encoding such a polypeptide (and fragments thereof), a promoter linked to such a polynucleotide, a cell or transgenic organism transformed with such a polynucleotide, a method for producing such a polynucleotide, an antibody (and method of its preparation) that specifically binds to such a polypeptide, a method for detecting a target polynucleotide in a sample, a method of treating a disease comprising administering to a patient such a polypeptide, a screening method for agonists or antagonists or compounds that bind to such a polypeptide, a method for screening for a compound that modulates the activity of such a polypeptide, a method of screening a compound for effectiveness in altering expression, a method of assessing toxicity of a test compound, a diagnostic test comprising said antibody, a microarray comprising such a polynucleotide, and a method of generating a transcript image by using such a microarray.

Inventions 2-67

An isolated polypeptide of SEQ ID NO: 2-67 and polynucleotide SEQ ID NO: 69-134 encoding such a polypeptide (and fragments thereof), a promoter linked to such a polynucleotide, a cell or transgenic organism transformed with such a polynucleotide, a method for producing such a polynucleotide, an antibody (and method of its preparation) that specifically binds to such a polypeptide, a method for detecting a target polynucleotide in a sample, a method of treating a disease comprising administering to a patient such a polypeptide, a screening method for agonists or antagonists or compounds that bind to such a polypeptide, a method for screening for a compound that modulates the activity of such a polypeptide, a method of screening a compound for effectiveness in altering expression, a method of assessing toxicity of a test compound, a diagnostic test comprising said antibody, a microarray comprising such a polynucleotide, and a method of generating a transcript image by using such a microarray.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 01/30042

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
EP 1067182	A	10-01-2001	AU 5849300 A	30-01-2001
			AU 5849400 A	30-01-2001
			AU 5849500 A	30-01-2001
			AU 5850500 A	30-01-2001
			EP 1067182 A2	10-01-2001
			EP 1197554 A1	17-04-2002
			EP 1203816 A1	08-05-2002
			EP 1201754 A1	02-05-2002
			WO 0104312 A1	18-01-2001
			WO 0104299 A1	18-01-2001
			WO 0104300 A1	18-01-2001
			WO 0104301 A1	18-01-2001
			JP 2002017376 A	22-01-2002
EP 1130094	A	05-09-2001	EP 1130094 A2	05-09-2001
			JP 2002017375 A	22-01-2002
WO 0058473	A	05-10-2000	AU 3774500 A	16-10-2000
			EP 1165784 A2	02-01-2002
			WO 0058473 A2	05-10-2000
WO 9940100	A	12-08-1999	AU 2577799 A	23-08-1999
			CA 2320625 A1	12-08-1999
			EP 1053245 A1	22-11-2000
			JP 2002502589 T	29-01-2002
			WO 9940100 A1	12-08-1999